

# Existing Use Evaluation for Zones 3, 4, & 5 of the Delaware Estuary Based on Spawning and Rearing of Resident and Anadromous Fishes

September 30, 2015



## Summary

Severe pollution problems have affected the Delaware Estuary for over 100 years. A key component to the estuary's restoration was the establishment of water quality criteria by the Delaware River Basin Commission in 1967. Faced with technical and financial challenges, these 1967 standards included a compromise position for 38 miles of the 133 mile long estuary: a limited "use" for Zones 3, 4, and upper Zone 5 that did not include "propagation" along with lower dissolved oxygen water quality criteria. Yet the 1967 goals were ambitious nonetheless, and the recovery in both dissolved oxygen and the living resources in these urban zones of the estuary have been tremendous. Recent data suggest that part of this recovery has been an exceedance of the 1967 goals for the "use" of the estuary, with substantial recovery in "propagation" even in Zones 3, 4, and 5. Because of Clean Water Act regulations requiring reconciliation of "existing uses" and "designated uses", these recent data warranted an evaluation to determine the extent to which a "propagation" use had been achieved for Zones 3, 4, and 5.

To evaluate the "propagation" use, DRBC reviewed available data for resident and anadromous fishes collected since 2000 that quantified spawning and/or rearing of early life stages of these estuarine species. This evaluation attempted to both assess whether any evidence of successful reproduction existed for each species in each estuary zone, and the degree to which successful reproduction had been restored for that species. Detailed reviews were possible for 9 fish species: Atlantic Sturgeon, American Shad, Striped Bass, White Perch, Bay Anchovy, Atlantic Silverside, Alewife, Blueback Herring, and Atlantic Menhaden. For all species evaluated, successful reproduction was clearly demonstrated in one or more of the compromised estuary zones. In addition, moderate to strong reproduction was demonstrated for multiple species in each zone indicating substantial recovery in the "propagation" use for Zones 3, 4, and upper Zone 5. Weak and inconsistent spawning by Atlantic Sturgeon, and limited spatial recovery in spawning and rearing by American Shad and Striped Bass, suggested that full restoration of the "propagation" use is not supported by the current available data. Based on this evaluation, DRBC staff recommends that partial restoration of the "propagation" use be recognized as the existing use for Zones 3, 4, and upper Zone 5 of the Delaware Estuary.

## **Introduction**

Water quality problems have plagued the tidal Delaware River for decades, with early evidence of severe hypoxia extending at least to 1912 (Philadelphia 1914). The USGS began daily recordings of dissolved oxygen at the Ben Franklin Bridge and at Chester, PA, in the 1960s and these records provide the most comprehensive documentation on the timing and magnitude of the estuary's historic hypoxia. The dissolved oxygen conditions for 1967 (see Figure 1) are largely representative of this severe hypoxia, and show hypoxic conditions beginning in May and extending into November.

By the 1950s, the various water quality problems (including dissolved oxygen) and their effects on both human populations and the Delaware Estuary ecosystem warranted federal investment to circumscribe the problem and chart a course for recovery (FWPCA 1966). The U.S. Public Health Service initiated the Delaware Estuary Comprehensive Study (DECS) in 1961, focusing on a computer model for the effects of municipal and industrial wastewater inputs to the Delaware Estuary, and how reductions in those wastewater loadings could lead to recovery of the dissolved oxygen within the estuary.

Near the conclusion of this study, the 1965 revisions to the Federal Water Pollution Control Act authorized interstate water quality standards to protect both human uses and ecological uses (i.e., "propagation of fish and wildlife", PL 89-234) of such interstate waters. Combined with the recent formation of the Delaware River Basin Commission in 1961 to address both interstate water supply and interstate water quality problems, the 1965 revisions to the Federal Water Pollution Control Act established the requisite authorities and framework for DRBC to adopt water quality standards for the Delaware River in 1967.

Controversy surrounded the initial establishment of dissolved oxygen criteria for the Delaware Estuary because of the significant cost-benefit tradeoffs in terms of ecological restoration versus wastewater treatment costs (FWPCA 1966, Wright and Porges 1971). Dissolved oxygen goals as high as 4.5 mg/L and as low as 1.0 mg/L were evaluated in the policy making process of the 1960s, with a compromise being reached where a dissolved oxygen criterion of 3.5 mg/L would be selected for Zones 3, 4, and upper Zone 5 of the estuary<sup>1</sup>. This 3.5 mg/L daily average dissolved oxygen criterion was then established through the 1967 adoption of DRBC's first Water Quality Regulations. It is important to note that this dissolved oxygen criterion has remained unchanged since 1967 (DRBC 2013).

In setting a dissolved oxygen criterion at the compromise value of 3.5 mg/L, and thus below the recognized requirements for fish and other aquatic life (e.g., FWPCA 1968), the DRBC, its member

---

<sup>1</sup> Zone 3 extends from River Mile (RM) 95 to RM 108.4; Zone 4 extends from RM 78.8 to RM 95; and the upper portion of Zone where the designated use and the dissolved oxygen criterion are consistent with Zone 3 and Zone 4 extends from RM 70 to RM 78.8. For an explanation of the River Mileage system, see DRBC's website at [www.nj.gov/drbc/basin/river/](http://www.nj.gov/drbc/basin/river/)

states, and the federal government set a goal for estuary restoration in these zones that protected only migratory fish passage and “low level maintenance of resident fish in the critical reaches of the estuary” (Wright and Porges 1971, pg. 1-3/9). In terms of current water quality standards language, two levels of aquatic life use protection were established for the estuary in 1967. For Zone 2 upstream of the hypoxia, and in lower Zone 5 and throughout Zone 6 below the hypoxia, full aquatic life use protections were established with the designated use of “wildlife, maintenance and propagation of fish and other aquatic life, passage of anadromous fish” or similar language (DRBC 1967; see Appendix A). In the estuary’s hypoxia regions of Zones 3, 4, and upper Zone 5, the “propagation” component to this use was removed and the aquatic life designated use (and thus the restoration goal) in these estuary zones was set to “wildlife, maintenance of resident fish and other aquatic life, passage of anadromous fish”.

Despite these compromises and the recognition in 1967 that full restoration of dissolved oxygen would be difficult, the recovery of both dissolved oxygen and most fish stocks in the Delaware Estuary have been remarkable. Striped Bass runs currently rival those of other major estuaries (e.g., Kahn et al. 2015), American Shad now freely migrate through, and spawn in, the Delaware Estuary (DRBFWMC 2011), and broader surveys of the fish assemblage show high catch rates and remarkable diversity (e.g., Pyle 2015). Sustained funding by DRBC and the USGS for the dissolved oxygen sensors at the Ben Franklin Bridge and Chester, PA, have documented the remarkable recovery in dissolved oxygen concentrations (see [waterdata.usgs.gov](http://waterdata.usgs.gov)). Figure 2 presents the full daily mean distribution of July dissolved oxygen at the Ben Franklin Bridge since the 1960s, showing the steady rise of summer dissolved oxygen beginning around 1980 and extending to ca. 2000, although dissolved oxygen has been quite variable since 2000 (see detailed summaries of 2000 to 2014 data in Appendix B).

As part of the recovery in dissolved oxygen and fish stocks, data began emerging that some degree of “propagation”<sup>2</sup> was occurring in Zones 3, 4, and upper Zone 5 where such fishery restoration was not anticipated in 1967. Because Clean Water Act regulations require any improvement in conditions to be captured and recognized, in part to prevent “backsliding” or loss of improvements gained through restoration efforts, data demonstrating successful reproduction in Zones 3, 4 and upper Zone 5 would serve as the initial basis for a re-evaluation of the designated uses for these estuary zones. Specifically, 40 CFR § 131.10(i) states “*Where existing water quality standards specify designated uses less than those which are presently being attained, the State shall revise its standards to reflect the uses actually being attained.*” In other words, if the “Existing Use” (i.e, the use presently being attained) is higher than the “Designated Use” currently adopted in regulations, reconciliation is needed under federal rules.

---

<sup>2</sup> The term “propagation” is not defined in DRBC’s Water Quality Regulations. Often used in the context of artificial rearing (e.g., fish hatcheries), other regulatory agencies have defined propagation as “reproduction of fish, aquatic life and wildlife within their natural environment.” (Delaware 2014). Based on the historical context of the aquatic life uses established in 1967, this report recognizes propagation somewhat more broadly as both the reproduction and the critical early life stage rearing necessary to recruit aquatic life into the population. For the fish species examined in detail in this report, this definition of propagation would include protection of spawning, eggs, larvae, and early stage juveniles.

The DRBC's Water Quality Advisory Committee began deliberations on this issue in 2009 as part of a broader evaluation of nutrient and dissolved oxygen conditions for the Delaware Estuary (see meeting minutes at [www.drbc.net](http://www.drbc.net)). A petition by three non-profit organizations in 2013 emphasized the need to proceed in reconciling the Existing Use with the Designated Use based on their evaluation of fishery data from the Delaware Estuary. Finally, the Water Quality Advisory Committee discussions culminated in a September 2013 resolution recommending the following:

*"The WQAC recommends that within one year the DRBC staff will prepare for the Committee a description of the existing aquatic life uses of the Delaware Estuary with citation to source materials which will be used to craft a formal Finding of Existing Uses that will be presented by the Committee to the Commissioners."*

In this report, DRBC conducts an evaluation of available data for resident and anadromous fishes to directly address the question as to whether, and to what degree, a broader "propagation" use has occurred in Zones 3, 4, and upper Zone 5 of the Delaware Estuary. The data sources and the methods for evaluating these data are first described. The results of the evaluation are then provided for each species in each of the three downgraded zones of the estuary. Finally, the individual evaluations are synthesized into a zone-specific assessment of the "propagation" use.

In addition, DRBC documents the dissolved oxygen conditions in the Delaware Estuary during a matching data window as an appendix to this report (see Appendix B). Although many factors interact to determine the population sizes of estuarine fish, the historic efforts to restore dissolved oxygen in the estuary as well as guidance from EPA on documenting existing uses together suggested that a summary of contemporary dissolved oxygen data would be a valuable component to a more comprehensive report.

---

## **Data Sources**

This report uses the results from three separate data collection efforts for resident and migratory fish species inhabiting the Delaware Estuary in order to evaluate the extent and consistency of “propagation” in Zones 3, 4, and 5. These data sets are described in detail below, but the decision to focus on these three data sets for the current evaluation is first discussed.

The evaluation of “propagation” in the broadest sense could include an evaluation of the reproductive success of many aquatic assemblages in the tidal Delaware Estuary, from planktonic invertebrates to freshwater mussels to anadromous fish. All are aerobic organisms utilizing the tidal Delaware River for reproductive output and subsequent nursery of juvenile organisms. Yet two factors recommend the use of the resident and migratory fish assemblages above other aquatic organisms. First, from a historical context, the restoration of dissolved oxygen in the Delaware Estuary centered on the restoration of fishery resources such as American Shad and Striped Bass. In order to assess the efficacy of the past 40+ years of dissolved oxygen restoration, its origins in fishery restoration argue for an evaluation of that fishery resource and the extent of successful “propagation” in this resource. Second, and more importantly, is the pragmatic reality that little or no data currently exist on the extent of successful reproduction for most organisms in the Delaware Estuary, and even less data exist on the spatial and temporal patterns in that reproductive success. Only the intensive monitoring of fishery resources for both regulatory and management objectives produces quantitative data on the location and timing of successful spawning as well as the later rearing stages. Therefore, the current evaluation of “Existing Use” for Zones 3, 4, and 5 of the Delaware Estuary relies exclusively on previously collected fishery data for eggs, larvae, and juveniles of resident and migratory fishes utilizing these zones of the Delaware Estuary.

Fishery surveys in the Delaware Estuary have been (and continue to be) conducted to fulfill many objectives, from permit compliance to fishery management obligations to basic academic research. Much of this effort is directed at assessment of adult populations, and such surveys provide limited utility for the current assessment of “propagation” in the Existing Use context because of their focus on older life stages. Other surveys do capture and quantify early life stages for many fish species, but have limited spatial and/or temporal resolution and thus only provide presence/absence answers to questions that (as shown later) require evaluations of strength and consistency. Impingement and entrainment reports at a single facility in an isolated year are examples of such valuable fishery data which have not been incorporated into this evaluation because they shed little additional light on the core questions beyond what is already established with more comprehensive data sources.

Three data collection efforts, however, directly target early life stages of one or more fish species and provide data that directly answer questions about the relative strength of spawning and early life stage survival in the Delaware Estuary across a broader spatial and temporal extent. In particular, these three data sets share three common features that make them particularly well-

suited for evaluating the “Existing Use” questions relative to the full “maintenance & propagation” use within DRBC’s water quality regulations:

- i. Quantitative data on the success of early life stages in Zones 3, 4, and/or 5 of the Delaware Estuary;
- ii. Multiple years of data at known and consistent spatial locations;
- iii. Data collection during the period from ca. 2000 to 2014 when dissolved oxygen conditions exhibited no long-term trends (see Figures 3 and 4).

The first of these three data sets was identified through discussions at the Delaware River Basin Fish & Wildlife Management Cooperative and comes from regulatory monitoring for the PSEG Salem Nuclear Generation Station, a large nuclear power plant located near the lower boundary of DRBC Zone 5 within the Delaware Estuary (~RM 51; see Figure 3). In advance of PSEG’s submission of its renewal application in 2006, ichthyoplankton surveys were required throughout the tidal Delaware Estuary (river and bay) during 2002, 2003, and 2004. Detailed descriptions of the methodology can be found in the respective Biological Monitoring Program annual reports (PSEG 2002, PSEG 2003, PSEG 2004). In summary, the entire estuary from Trenton to the Atlantic Ocean was divided into 14 “zones” (see Figure 4; henceforth referred to as “reaches” to avoid confusion with DRBC water quality management zones), with multiple grids within each reach. Surveys were conducted twice monthly from April through July resulting in 8 surveys each year. During any survey, station locations were randomly selected within each reach from among the grids, with multiple stations (ranging from 2 to 14) within each reach and somewhat even effort per unit area across the estuary. Sampling consisted of a 1 m diameter conical plankton net with 500 µm mesh netting and 500 µm mesh capture bucket. Depending on depth, single integrated tows proceeded at successively deeper vertical positions for a total trawl time of 4 to 6 minutes. Equal amounts of time were sampled at each depth, with no surveys at station locations with depths 10 ft or less, and increasing numbers of strata for increasing depths: for 10-20 ft depth, sampling at 5 ft and 15 ft strata; for 20-30 ft depths, sampling at 5 ft, 15 ft, and 25 ft strata; etc. Samples were preserved in the field with 10% formalin and stained with Rose Bengal. In the lab, a complex subsampling procedure was used, and identifications were made to species level, if possible, for each life stage. However, only “target species” were enumerated for all samples. These target species were Blueback Herring, Alewife, American Shad, Atlantic Menhaden, Bay Anchovy, Atlantic Silversides, White Perch, Striped Bass, Bluefish, Weakfish, Spot, and Atlantic Croaker along with two invertebrate groups (*Neomysis americana*, *Gammarus* spp.).

Data from the PSEG Ichthyoplankton surveys are provided as event summaries for each of PSEG’s reaches, with results broken out by species for the following life stages: eggs, yolk sac larvae, post-yolk sac larvae, juveniles, adult, and undetermined larvae (see Appendix D). Thus, data from multiple survey stations are pooled to represent each of the PSEG reaches in the estuary for each survey date. The PSEG Ichthyoplankton results provide the most direct measure of spawning activity for key species such as American Shad and Striped Bass, although the high variability in catch rates for eggs and yolk sac larvae are important to keep in mind given their episodic occurrence and short life-stage duration. In addition, the PSEG Ichthyoplankton data provide some measure of the success by tracking each fish species through each of their early life stages, a key component in the overall assessment of successful “propagation” for the target species

included in the PSEG surveys. As shown in Figure 4, however, the PSEG reaches cross the boundaries for DRBC zones, leading to ichthyoplankton data that apply to more than a single DRBC zone.

The second data set identified for inclusion in this evaluation originated from a member of the Delaware River Basin Fish & Wildlife Management Cooperative. The NJ Division of Fish & Wildlife, Bureau of Marine Fisheries, conducts Delaware Estuary beach seine surveys from June through November each year, with initial sampling for these surveys beginning in 1980 and continuing every year through the present. These surveys constitute a key fishery independent monitoring project that is required under the Atlantic State Marine Fisheries Commission's "Fishery Management Plan for Striped Bass." Designed primarily for monitoring juvenile Striped Bass, the survey also currently records the abundance of all species captured. Full survey details are contained in annual reports posted on the Bureau of Marine Fisheries website (e.g., 2014 report at [www.njfishandwildlife.com/artdelstudy15.htm](http://www.njfishandwildlife.com/artdelstudy15.htm)), with key survey details summarized here. Seining occurs during daylight hours at beach locations within the tidal Delaware River with a 100 ft long, 6 ft deep bag-containing seine net with ¼ inch mesh. In addition to counts by species, lengths are measured for target species (see [web](#) report for list) and the range of sizes is recorded for non-target species. Sampling currently occurs at 32 fixed stations beginning upstream at RM 126 (Newbold Island) in DRBC Zone 2 and extending downstream to RM 53 (Augustine Beach) near the lower end of DRBC Zone 5 (Figure 3). Sampling stations and effort have remained highly consistent for the 2000-2013 period used in the current evaluation of Existing Use. Greater sampling effort occurs in Zones 2, 4, and 5 than in Zone 3 because fewer suitable "beach" seining locations are available in Zone 3. Fish species are typically identified in the field, with nearly 100 species identified through the duration of the survey.

It is important to distinguish the nature of data collected in the New Jersey Seine Survey from the data collected in the PSEG Ichthyoplankton survey, and to acknowledge the limitations with respect to the Existing Use question evaluated in this report. The New Jersey Seine Survey begins to capture fish of different species typically when they reach between 20 mm and 30 mm in length. Depending on the species, this could represent nearly mature fish (e.g., Bay Anchovy) or it could represent early young-of-year fish recently spawned and reared near an individual beach seine location (e.g., Striped Bass). In addition, because of varying fidelity to the estuary itself beyond the early nursery stage, some species can be more clearly identified as young-of-year juveniles spawned and/or reared within the estuary (e.g., American Shad) while other species use the estuary for all or a larger portion of their lifecycle, thus mixing young-of-year juveniles with older juvenile and adult fish (e.g., White Perch). As a result, the New Jersey Seine Survey provides a measure of the combined success of both spawning and rearing through the minimum size threshold for those species where young-of-year fish can be clearly identified (American Shad, Alewife, Blueback Herring, Striped Bass [note: although Striped Bass utilize the estuary for multiple years, the NJ Bureau of Marine Fisheries uses a suite of criteria to accurately distinguish young-of-year from older fish]). For species where the distinction between young-of-year and older fish cannot be made (e.g., Atlantic Silversides, Bay Anchovy, Hogchoker, White Perch), these data provide more of a combined population estimate and an indicator of whether successful reproduction and rearing had occurred, but not a more narrow measures of

“propagation” alone. In addition, and in contrast to the PSEG Ichthyoplankton surveys, the New Jersey Seine Survey samples and identifies all fish species collected in their survey. For the current Existing Use assessment, the New Jersey Seine Survey data are therefore more important for species whose young-of-year could be clearly distinguished.

The New Jersey Seine Survey thus provides important data for assessing propagation and, in particular, provides a historical context upon which to assess the current results. The survey’s origins in the early 1980s cover the period when dissolved oxygen was only beginning to recover from decades of chronic hypoxia and when some migratory fishes were initiating their recovery within the Delaware Estuary. Such origins, combined with an unbroken time series, provide an insightful perspective on current spawning efforts. Those early years of the survey typically showed very low young-of-year abundance, especially in Zones 3 & 4, and these low catches delineate the lower end of the “propagation” scale when fish spawning and recruitment was either low or failing. Although the lower end of the “propagation” scale is therefore clearly delineated, the upper end of the recruitment scale (e.g., pre-European numbers) that would establish a benchmark for highly successful propagation remains uncertain making it difficult to assess how complete the recovery of spawning and rearing has been without that historical upper range clearly delineated. Nevertheless, the temporal extent of the New Jersey Seine Survey at a minimum provides a clear measurement scale for both poor recruitment as well as moderate to strong recruitment for many estuarine species, yielding a scale that is well-positioned to answer questions about the success or failure of spawning and recruitment of estuarine fish species.

The third and final data set used for this report involves multiple targeted collections and tracking of early stage juvenile Atlantic Sturgeon (also known as river-resident juveniles; see species results below). Beginning in 2009, Delaware’s Division of Fish & Wildlife shifted their sampling methods for juvenile Atlantic Sturgeon to better capture spring spawned young-of-year fish from the Delaware Estuary as well as both age-1 and age-2 fish which had overwintered in the Delaware Estuary. These targeted surveys for early stage, river-resident juvenile Atlantic Sturgeon were completed in 2009, 2010, 2011, 2012, and 2014. In addition, Brundage and O’Herron conducted separate surveys in order to tag and track juvenile Atlantic and Shortnose Sturgeon (Brundage and O’Herron 2010). For both research groups, surveys primarily involved small-mesh gill net deployment in targeted locations, with supplemental data collected through active and passive tracking of tagged individuals collected in the gill net surveys. Tracking also provided spatial data to improve the placement of gill net surveys. Gill net surveys focused on areas between Carney’s Point, NJ (RM 71) and Marcus Hook, PA (RM 81) covering the upper portion of Zone 5 and the lower portion of Zone 4, although additional surveys further upstream and downstream were also included. Passive and active tracking extended throughout the Delaware Estuary from Zone 2 through Zone 6 (Delaware Bay). Detailed methods and results can be found in the various annual reports (Brundage & O’Herron 2010, Fisher 2010, Fisher 2011a, Fisher 2011b, Fisher 2012, Fisher 2013, Brundage & O’Herron 2014).



## **Evaluation Methodology**

The recovery of fish spawning and rearing in the Delaware Estuary involves a gradual process of increasing success as water quality, habitat, and overall stocks improve toward conditions existing prior to severe human alteration. Initially, some limited spawning and rearing may occur in the estuary, particularly for fish species tolerant of heavy human influences such as the hypoxia historically present in the Delaware Estuary. With continued recovery of impaired conditions, further increases in the spawning and rearing would be expected as more species of fish and more individuals of each species find increasingly suitable conditions for growth and development. With substantial restoration of conditions and with the resulting increases in population sizes, the utilization of the estuary may approach levels seen historically for many, if not most, fish species.

Recognizing the continuum in this recovery spectrum, the current report seeks to go beyond a simple yes/no (or binary) answer to the question about fish “propagation” in the estuary. To achieve a more continuous representation in the restoration of fish “propagation” for the Delaware Estuary, and to seek a relatively objective answer to these questions which undoubtedly contain many subjective elements, the evaluation of data for each fish species was broken into a series of yes/no questions that represent increasing levels of success in restoring each species’ use of the Delaware Estuary. An example is shown in Table 1. These yes/no questions were evaluated separately for each fish species and for each of the three DRBC water quality zones where the “propagation” use is not currently designated in the DRBC Water Quality Regulations (Zones 3, 4, & 5; note that Zone 5 has different “uses” in the upper and lower portion; these two sections of Zone 5 were evaluated separately<sup>3</sup>). And to acknowledge the lack of clear objective thresholds for decisions on each question, a middle category between “yes” and “no” was allowed and used in numerous settings when the data were considered equivocal.

Initial stages in “propagation” success can be demonstrated with some evidence of eggs, larvae, and/or early juvenile stages in a specific estuary zones. Even if such data are not consistent year after year, the presence of measureable numbers of early life stage individuals allows for an affirmative answer to the question, “Is there any evidence of successful reproduction?” This is the first question addressed along the continuum.

Further restoration would then be demonstrated with increased numbers of early life stage individuals and/or increased consistency among years in catch data for these early life stages. Such moderate levels of consistency would permit an affirmative answer to the question, “Is there regular evidence of successful reproduction for this species?” This is the second and middle category for evaluating the degree of successful reproduction.

More complete restoration is then demonstrated with a high level of consistency and substantial numbers of these early life stage individuals in most years. Of course, estimates of population sizes and spawning success are notoriously uncertain or variable for many fish species, so a high

---

<sup>3</sup> Upper Zone 5 has “maintenance” only while lower Zone 5 has the full “maintenance and propagation” use

degree of consistency does not necessarily require equal numbers year after year without any breaks in the string of successful year classes. Instead, this high variability is acknowledged and helps establish expectations for the third question, “Are there strong numbers and high consistency of successful reproduction?”

Table 1. Example of the Evaluation Matrix. (Overall structure is explained in text. The first three color-coded columns in the table indicate priorities in different studies. The first shows whether the 1979 Task Force (Nadeau et al. 1979) identified the species as a key “migratory or marine/estuarine” (m), “resident (freshwater/brackish water)” (f). The second shows a red box for those species included as part of PSEG Salem’s targeted species list. The third column shows in blue species averaging more than 10,000 fish caught per year and in green species averaging more than 10 fish caught per year in the NJ Seine Survey.

<b>Species X</b>							
1979 Task Force PSEG target (12 spp) NJ: 10 K/yr and 10/yr groups	Common name	Scientific name		Any evidence of successful reproduction?	Regular evidence of successful reproduction?	Strong numbers & high consistency of successful reproduction?	"Nursery" with juveniles
	m	Species X	<i>Genus species</i>				
			<b>Zone 3</b> <i>RM 95-108.4</i>	Y	Y	?	Y
			<b>Zone 4</b> <i>RM 78.8-95</i>	Y	?	N	Y
			<b>Zone 5 (upper)</b> <i>RM 70-78.8</i>	Y	N	N	Y
			<b>Zone 5 (lower)</b> <i>RM 48.2-70</i>	N	N	N	?

This third and highest level of “propagation” restoration used in this report proved to be the most problematic and ambiguous of the three categories for evaluating the fishery data. Two issues arose. First, it was not always clear how consistent and how high catch data needed to be to qualify for a “yes” in this category, with discussions of a more rigorous threshold. Although a number of candidate thresholds were considered, a clear choice for a threshold was not apparent and none was employed in the current evaluation. The second limitation with this highest category involves the degree to which it represents full successful restoration of “propagation” for any particular species. Although some managed stocks (e.g., Striped Bass) have officially been designated as “restored” by management agencies, quantitative historical data are typically sparse and involve comparisons across different sampling methods. Thus, even for populations who have seen substantial recovery, it is difficult or impossible to infer whether population sizes have recovered to levels seen prior to major alteration of the estuarine ecosystem (water quality, habitat, etc.). In the absence of such comparative data, this study contends that an affirmative

answer to this highest category of “propagation” still leads to ambiguity on the question of whether the “propagation” use has been fully restored or whether continued improvement toward full restoration remains possible; both levels of restoration may be implied by a “yes” to this question.

Because subjectivity persisted even with these attempts to minimize its influence, initial evaluations for key species were presented to fishery biologists both at a Delaware River Basin Fish & Wildlife Cooperative technical committee meeting, and in smaller group settings with fishery biologists from the states of Delaware, New Jersey, and Pennsylvania. Data from multiple sources were reviewed for the key species, and the decisions for each question were evaluated. Typically, a consensus could be reached on the appropriate answer to each question, but complete agreement was not always possible.

Two additional and important clarifications are needed to understand the current data evaluation, with the two issues being related. The first relates to different evaluation standards for different species. The second relates to different expectations for fish species in different zones of the estuary.

This report did not typically apply different standards in the data evaluation matrix for different species. As described above, the data that would qualify for an affirmative or negative evaluation to the “any” vs “regular” vs “strong” questions were largely the same for all species. One important distinction was made to account for the salinity regime in which spawning occurs for each species. For obligate freshwater spawning fish, affirmative answers to the questions in freshwater zones (i.e., Zones 3 and 4) required increasing demonstration of eggs and larval presence and abundance. However, a number of marine and brackish water fish species use different salinity regimes for different stages in their spawning and larval rearing phases (e.g., Bay Anchovy, Atlantic Menhaden). For these more marine or brackish water species, the demonstration of low to moderate levels of “use” for an estuary zone could be demonstrated without substantial data on eggs, in particular. Thus, the use of upper Zone 5 for Bay Anchovy was assessed as affirmative for the “regular” level because substantial and moderately consistent numbers of larval and juvenile fish were observed even though eggs were not similarly strong. In contrast, the lack of consistent egg numbers in Zone 4 for American Shad, despite substantial evidence of larval and juvenile use in this zone, led to an ambiguous evaluation for whether American Shad use qualified as “regular evidence of successful reproduction”. This approach constituted the only incorporation of life history and habitat use into the evaluation matrices, and the only change in the standards by which different species were evaluated.

More broadly, each fish species evaluated in this report does not have the same expected use in all zones of the estuary. As mentioned above and in the species descriptions below, different salinity requirements and preferences among species leads to substantially different expectations for which fish should be “propagating” in the brackish lower Zone 5 than in the purely freshwater Zone 3. In addition to salinity, each species has different habitat preferences that make even similar salinity zones (e.g., Zones 2, 3, and 4) more or less suitable for each species. These differences in the basic ecology for each fish leads to different expectations for

what “successful propagation” looks across multiple species for each zone of the estuary. A group of primarily fisheries biologists attempted to define these different expectations for many of the key species (although Atlantic Sturgeon were omitted) in 1979 when evaluating the fishery and D.O. requirements in the estuary (Nadeau et al. 1979; see Appendix C for Table 1 of their evaluation). Given the history of gross pollution in the estuary and the sparse availability of reliable historic data, any such attempt at setting the precise expectations for fish recovery in the estuary is subject to debate, and the evaluation in Nadeau et al. would benefit from an updated review of these expectation in light of the observed recovery seen since 1979. Unfortunately, a thorough review of the expectations for each species based on both historical data and current biological requirements was beyond the scope of the current study. Nevertheless, the Table 1 expectations in Appendix C begin to distinguish the varying expectations for “successful propagation” among the estuary’s zones. In particular, the 1979 evaluation shows, for instance, that Bay Anchovy were not expected to use Zone 4 in appreciable numbers for spawning or nursery, while American Shad were not expected to use Zone 5 in appreciable numbers for spawning and nursery activities. These broader expectations for each zone are therefore used in the final evaluation for each zone of the estuary (see Conclusions section), while acknowledging some uncertainty for the exact expectation in each zone.

## Results – Species Level Evaluations for Successful “Propagation”

Each species’ evaluation is presented separately below beginning with fish whose data most directly addresses the questions of “propagation” for the estuary

### ATLANTIC STURGEON



Atlantic Sturgeon (*Acipenser oxyrinchus*) are large (to 14 ft / 4 m), long-lived (60 years), and late-maturing (5 to 22 years) anadromous fish that use tidal estuaries for spawning (ASSRT 2007 and references therein). Spawning occurs in a zone beginning at the salt front and extending upstream to the fall line, although exact locations are not known for each estuary. In the Delaware Estuary, this zone extends from upper Zone 5 through Zone 2. Spawning occurs in spring and early summer, perhaps as late as mid to late June in the Delaware River (Simpson and Fox 2007). Individuals are not believed to spawn every year but instead may return every 1 to 5 years to their spawning grounds. Juvenile sturgeon become tolerant to brackish water and reside in both freshwater and brackish water areas within their natal estuary for one or more years (river-resident juveniles). Juveniles migrate out of their natal estuaries beginning at age 2 and will mix with Atlantic Sturgeon from other estuaries in coastal and estuarine habitats, but will then return to their natal estuary to spawn (McCord et al. 2007). It is important to note that laboratory work has shown that young juvenile fish sustain high mortality under depressed dissolved oxygen conditions (e.g., Secor and Gunderson 1998).

Recent surveys have confirmed successful reproduction and recruitment of Atlantic Sturgeon in the Delaware Estuary. Although isolated young-of-year catches have been noted for decades, the first documentation of at least a small to moderate year class occurred with the capture of over 60 young-of-year sturgeon beginning in the fall of 2009 (Fisher 2010, Brundage & O’Herron 2010). However, subsequent surveys have documented only sporadic success for Atlantic Sturgeon spawning and rearing. The following year in 2010, no young-of-year Atlantic Sturgeon were captured when fishing in a manner consistent with the 2009 efforts but at reduced efforts. In 2011, another successful year class was documented when nearly 50 young-of-year were captured (Fisher 2012). In 2012, only two young-of-year Atlantic Sturgeon were captured during targeted surveys (Fisher 2013). Limited surveys were conducted in 2013, so no comparable data


exist. Then in 2014, another successful year class was documented with over 200 river-resident juveniles captured during the targeted surveys, although size ranges prevented clear indications for the number of young-of-year fish (I. Park *personal communication*, H. Brundage *personal communication*). It is important to note that survey effort was not consistent among years, so variations in catches may partly be attributable to variable effort.

Gill net captures of young-of-year Atlantic Sturgeon were all made in lower Zone 4 and upper Zone 5 (above RM 70) within the Delaware Estuary. Tagging of select individuals subsequently showed extensive use of all zones in the Delaware Estuary for juvenile rearing (Brundage & O’Herron 2010, Fisher 2011b, Brundage & O’Herron 2014). Although the location of Atlantic Sturgeon spawning grounds within the Delaware River remains unknown and somewhat controversial (Simpson 2008, Breece et al. 2013), the initial collection of young-of-year fish in both upper Zone 5 and lower Zone 4 leads to an affirmative answer within these two zones for the question, “Is there any evidence of successful reproduction?” For Zone 3 and lower Zone 5, the movement of tagged age-0 fish into and through these zones, combined with the possibility of suitable spawning habitat in both Zone 2 and Zone 3, leads to the uncertain and inconclusive answer of “?” for the question, “Is there any evidence of successful reproduction?” for Zone 3 and for lower Zone 5. Furthermore, the initial positive results in 2009 combined with the failure or near failure to capture young-of-year fish in both 2010 and 2012 leads to a “no” decision for all zones with respect to both of the two higher classifications: “Is there regular evidence of successful reproduction for this species?” and “Are there strong numbers and high consistency of successful reproduction?”

Table 2. Evaluation Matrix for Atlantic Sturgeon in Zones 3, 4, and 5 of the Delaware Estuary

<b>Atlantic Sturgeon</b>						
1979 Task Force PSEG target (12 spp) NJ: 10 K/yr and 10/yr groups	Common name	Scientific name	Any evidence of successful reproduction?	Regular evidence of successful reproduction?	Strong numbers & high consistency of successful reproduction?	"Nursery" with juveniles
		Atlantic Sturgeon	<i>Acipenser oxyrinchus</i>			
		<b>Zone 3</b> RM 95-108.4	?	N	N	Y
		<b>Zone 4</b> RM 78.8-95	Y	N	N	Y
		<b>Zone 5 (upper)</b> RM 70-78.8	Y	N	N	Y
		<b>Zone 5 (lower)</b> RM 48.2-70	?	N	N	Y

## AMERICAN SHAD



Maryland DNR

American Shad (*Alosa sapidissima*) are an anadromous fish of the western Atlantic Ocean, with spawning runs extending from the Saint Johns River, Florida, to the Saint Lawrence River, Canada. Spawning occurs in both tidal and non-tidal freshwater regions of coastal rivers and streams during spring and early summer. Juvenile shad spend their first summer within their natal river and migrate to the Atlantic Ocean in late summer and fall of that same year. American Shad have been the focus of fisheries and water quality management programs for over 100 years in the Delaware River (e.g., Gay 1892, Hardy 2009) and were a key component to the dissolved oxygen recovery programs of the 1960s and 1970s (FWPCA 1966, Wright and Porges 1971). American Shad have been shown to be sensitive to low dissolved oxygen levels (Stier and Crance 1985).

Summary results from PSEG Ichthyoplankton data are presented in Figure 5. American Shad eggs were primarily caught in lower Zone 2 and upper Zone 3, with broader distributions for both yolk sac larvae and post-yolk sac larvae. The broader distribution of post-yolk sac larva likely represents passive movement via river currents as well as increased catchability for this life stage due to its longer persistence time in the estuary. Overall, these data show successful “propagation” for American Shad in Zones 2, 3, and 4 of the estuary, with some utilization of upper Zone 5, as well. The strength and consistency, however, varies by zone. In Zone 3, relatively high and consistent larval fish numbers were collected in the three survey years, but egg numbers varied markedly within the zone in all three years, with higher egg densities collected in the lower areas of Zone 2 and the upper areas of Zone 3 compared to the lower areas of Zone 3. In Zone 4, moderately strong densities of larvae were collected in all three years, but American Shad eggs were rarely collected.

Figure 6 presents the young-of-year catches for American Shad in Zone 3, 4, and 5 for the New Jersey Seine Survey. The main pattern in these results is the restoration of the nursery in the Delaware Estuary beginning in the late 1980s. In addition, even with fewer stations and thus lower sampling effort in Zone 3, these results show high and relatively consistent use of Zone 3 when compared to Zones 4 and 5 downstream. Zone 4 results are similar to the Zone 3 results, although greater frequency of low catches is seen in Zone 4 than in Zone 3 for any given year. Finally, young-of-year American Shad show substantially less use of Zone 5 than of the upstream zones

Based on the combined results from the two surveys, American Shad propagation has been documented in Zones 3 and 4 since 2000, with some sparse evidence for larval fish use and stronger evidence for juvenile fish use in upper Zone 5. An affirmative answer is indicated in all three zones for this first question, “Is there any evidence of successful reproduction?” while a

negative response is indicated for lower Zone 5. For Zone 3, the high egg catches in the upper areas of the zone and the strong larval fish numbers in all years further indicate an affirmative answer to the second question, “Is there regular evidence of successful reproduction for this species?” However, the inconsistent egg densities in all three years within Zone 3 indicate some decrease in the utilization of lower sections of Zone 3, and an ambiguous “?” is thus provided for the highest level of use and the question, “Are there strong numbers and high consistency of successful reproduction?”

In Zone 4, the sparse egg catches and the tapering numbers of larval fish provide ambiguity to the question, “Is there regular evidence of successful reproduction for this species?” Certainly some regular use of Zone 4 is indicated, but for later stages of immature and juvenile American Shad, and an ambiguous “?” answer is provided for Zone 4 at this second level. Furthermore, these same patterns and the lower catches of young-of-year American Shad in Zone 4 together indicate a “no” answer to the highest utilization question, “Are there strong numbers and high consistency of successful reproduction?”

Finally in Zone 5, only sparse numbers of larval American Shad have been collected in the upper section above RM 70, although increased use by young-of-year fish is noted. This indicates substantial use by later stages of juvenile American Shad more consistent with a nursery function. Thus, a negative response is indicated for both of the higher use questions: “Is there regular evidence of successful reproduction for this species?” and “Are there strong numbers and high consistency of successful reproduction?” For lower Zone 5, the absence of any eggs or larvae leads to negative responses for all questions.

Table 3. Evaluation Matrix for American Shad in Zones 3, 4, and 5 of the Delaware Estuary

		<b>American Shad</b>						
1979 Task Force PSEG target (12 spp) NJ: 10 K/yr and 10/yr groups		<b>Common name</b>	<b>Scientific name</b>		<b>Any evidence of successful reproduction?</b>	<b>Regular evidence of successful reproduction?</b>	<b>Strong numbers &amp; high consistency of successful reproduction?</b>	<b>"Nursery" with juveniles</b>
	m	American Shad	<i>Alosa sapidissima</i>					
				<b>Zone 3</b>	Y	Y	?	Y
				RM 95-108.4				
				<b>Zone 4</b>	Y	?	N	Y
				RM 78.8-95				
			<b>Zone 5 (upper)</b>	Y	N	N	Y	
			RM 70-78.8					
			<b>Zone 5 (lower)</b>	N	N	N	?	
			RM 48.2-70					



## STRIPED BASS



Striped Bass (*Morone saxatilis*) are an anadromous species native to the western Atlantic Ocean from Florida to Canada. They are long-lived (up to 30 years) with males maturing relatively early (100% at 3 years) while females mature later at a much larger size (50% at 6 years; ASMFC 2013). The Delaware Estuary is one of the four primary spawning grounds for Striped Bass including the Hudson River and the Chesapeake Bay and its tributaries (Able and Fahay 1998). Spawning occurs in tidal freshwater to oligohaline zones (less than 3 ppt salinity; Wang and Kernehan 1979). For the Delaware Estuary, spawning begins in early April and can extend into June and July, although peak activity occurs in late April or early May (Wang and Kernehan 1979). Juvenile fish spend 2 or more years in coastal estuaries (and may move upstream into non-tidal areas) before joining coastal migratory populations. Striped Bass population management and recovery has been a model for other species' management, with Striped Bass populations largely recovering from historic lows by the 1990s (e.g., ASMFC 1995).

Summary results from the PSEG Ichthyoplankton surveys are presented in Figure 7. Consistent patterns are seen across the three years of surveys, with egg and larval catches extending from Zone 2 through Zone 5 each year and peak abundances in Zone 4 and, particularly, Zone 5. Reduced numbers are apparent in Zone 3. Overall, these ichthyoplankton data largely match adult Striped Bass spawning survey patterns, with consistent documentation of Striped Bass spawning beginning in Zone 2 and peaking in Zones 4 and 5 (M. Kaufman *personal communication*, G. Murphy *personal communication*).

Results from the New Jersey Seine Survey are presented in Figure 8. It is worth emphasizing that the New Jersey Seine Survey is primarily designed and implemented to accurately represent Striped Bass young-of-year recruitment. Among the many strengths in these data, New Jersey fisheries biologists use highly developed length criteria that vary through the season to separate young-of-year fish from age-1 and older fish. The most striking pattern in these data, like with the American Shad, is the overall recovery of Striped Bass spawning in the Delaware Estuary since the 1980s that is related to both the restoration of dissolved oxygen within the Delaware Estuary and the broader restoration of the Striped Bass stock along the Atlantic coast. Within the more recent past, Figure 8 largely corroborates the patterns seen in the PSEG Ichthyoplankton results, with Zone 4 and Zone 5 demonstrating notably similar data distributions for young-of-year fish. In the seine data, however, there is even greater similarity in the Zone 4 and Zone 5 results indicating both the upstream shift in Striped Bass towards Zone 4 as summer progresses (and the salt line within the Delaware Estuary moves increasingly upstream), as well as an increased use of Zone 4 for young-of-year nursery grounds. Like the ichthyoplankton results, decreased catches


in Zone 3 further highlight the more limited use of Zone 3 by Striped Bass early life stages compared to both Zone 4 and Zone 5 downstream.

The combined and largely consistent patterns from the ichthyoplankton surveys and the seine surveys clearly demonstrate successful reproduction by Striped Bass in all zones of the Delaware Estuary. Within the evaluation matrix, an affirmative answer to the question, “Is there any evidence of successful reproduction?” is thus indicated for Zones 3, 4, and 5. In addition, the regularity in even the modest ichthyoplankton and young-of-year catches for Zone 3, combined with the stronger catches in Zones 4 and 5, indicate an affirmative answer for all three zones to the questions, “Is there regular evidence of successful reproduction for this species?” Finally, the consistently high catches of eggs and larvae in Zone 5 provides a clear indication for “strong numbers and high consistency of successful reproduction” while the somewhat reduced egg and larval catches in Zone 4 create an ambiguous response to this question. In Zone 3, where numbers are clearly reduced for all early life stages, a negative response is indicated for this third question within the evaluation matrix.

Table 4. Evaluation Matrix for Striped Bass in Zones 3, 4, and 5 of the Delaware Estuary

<b>Striped Bass</b>							
1979 Task Force PSEG target (12 spp) NJ: 10 K/yr and 10/yr groups	Common name	Scientific name		Any evidence of successful reproduction?	Regular evidence of successful reproduction?	Strong numbers & high consistency of successful reproduction?	"Nursery" with juveniles
	m	Striped Bass	<i>Morone saxatilis</i>				
			<b>Zone 3</b> RM 95-108.4	Y	Y	N	Y
			<b>Zone 4</b> RM 78.8-95	Y	Y	?	Y
			<b>Zone 5 (upper)</b> RM 70-78.8	Y	Y	Y	Y
			<b>Zone 5 (lower)</b> RM 48.2-70	Y	Y	Y	Y

## WHITE PERCH



White Perch (*Morone americana*) is the second of two species in the genus *Morone* native to the Delaware Estuary, and is one of the most abundant fish species in the Delaware Estuary. In contrast to Striped Bass, White Perch are semi-anadromous and typically remain resident within a river system while conducting seasonal migrations from brackish waters to tidal freshwaters as well as migrating above the head-of-tide into non-tidal rivers and streams. Spawning occurs in oligohaline tidal areas (less than 3 ppt salinity) as well as both tidal and non-tidal freshwater zones (Wang and Kernehan 1979). For the Delaware Estuary, spawning begins in late March and can extend into June (Wang and Kernehan 1979).

White Perch results from the PSEG Ichthyoplankton surveys are presented in Figure 9. These results demonstrate a broad utilization across most of the Delaware Estuary by White Perch, with high egg catches beginning in the upper reaches of Zone 2 and extending into Zone 5. Likewise, larval fish densities persisted in all three years at high levels through Zones 2, 3, 4, and much of Zone 5. Although Zone 5 shows some tapering in the distribution, it is noteworthy that one or more reaches in Zone 5 meet or exceed the catches for larval White Perch demonstrated in the upstream estuary zones.

White Perch catches in the New Jersey Seine Survey are presented in Figure 10. It is important to recognize that the results in Figure 10, unlike the seine results from previously reviewed species, do not represent young-of-year fish alone. Instead, because of the resident behavior of White Perch and because of overlapping size distributions among age classes, the results in Figure 10 indicate the combined abundance of both young-of-year fish and mostly younger age classes of resident White Perch. This blending of age classes limits the ability to infer the degree of successful “propagation” for White Perch based on these data. These results (and additional length distribution data provided by NJ Bureau of Marine Fisheries) primarily assist in demonstrating broad utilization by young-of-year and older age classes throughout the estuary, and confirm that young-of-year fish are indeed using Zones 3, 4, and 5 as early life stage rearing habitat.


As mentioned, the assessment of reproductive success and the extent of “propagation” for White Perch can mainly be inferred through the PSEG Ichthyoplankton surveys. The New Jersey Seine Surveys clearly demonstrate that young-of-year juveniles currently use Zones 2, 3, 4, and 5, helping to support an affirmative answer to the question, “Is there any evidence of successful reproduction?” The PSEG Ichthyoplankton provide even greater insight into the strength of this utilization by White Perch, demonstrating strong and relatively consistent egg and larval catches

in Zones 2, 3, 4, and upper Zone 5. Combined, these data reveal a broad and highly successful utilization of the Delaware Estuary and demonstrate an affirmative assessment to the two additional matrix questions of “Is there regular evidence of successful reproduction for this species?” and “Are there strong numbers and high consistency of successful reproduction?” for all zones except lower Zone 5. Indeed, the success of White Perch across such a broad extent of the Delaware Estuary provides a scale upon which to evaluate the other estuarine fish species. Only in lower Zone 5 did the data show decreasing use by White Perch, with low egg catches and inconsistent larval densities within the zone and among years. These results provide some ambiguity to the second question, “Is there regular evidence of successful reproduction for this species?” within lower Zone 5, and a negative response to the question, “Are there strong numbers and high consistency of successful reproduction?”

Table 5. Evaluation Matrix for White Perch in Zones 3, 4, and 5 of the Delaware Estuary

		<b>White Perch</b>						
1979 Task Force PSEG target (12 spp) NJ: 10 K/yr and 10/yr groups		Common name	Scientific name		Any evidence of successful reproduction?	Regular evidence of successful reproduction?	Strong numbers & high consistency of successful reproduction?	"Nursery" with juveniles
	<b>f</b>		White Perch	<i>Morone americana</i>				
				<b>Zone 3</b> RM 95-108.4	Y	Y	Y	Y
				<b>Zone 4</b> RM 78.8-95	Y	Y	Y	Y
				<b>Zone 5 (upper)</b> RM 70-78.8	Y	Y	Y	Y
				<b>Zone 5 (lower)</b> RM 48.2-70	Y	?	N	Y

## BAY ANCHOVY



*Diana Rome Peebles*

Bay Anchovy (*Anchoa mitchilli*) is perhaps the most abundant fish of the Delaware Estuary (Wang and Kernehan 1979). They inhabit a broad range of estuarine and coastal marine habitats, and their distribution spans all salinity regimes of the Delaware Estuary. They are short-lived (typically 3 years or less) and repetitively spawn throughout late spring and summer in moderate to high salinity waters (typically 5 ppt or higher). Larval fish appear to have some ability to migrate up-estuary, with both viable larvae and juvenile fish collected at higher densities in more freshwater areas than eggs (Wang and Kernehan 1979, Able and Fahay 1998).

Bay Anchovy results from the PSEG Ichthyoplankton surveys are presented in Figure 11. It is important to note that even higher densities of eggs and larvae (peaks annually exceeding 100,000 and thus natural log scale of 12 on Figure 11) were collected further downstream in Delaware Bay; these higher Delaware Bay catches are not shown in the figure. Given the Bay Anchovy's natural history, the pattern in Figure 11 primarily demonstrates a preference for higher salinities in spawning. However, substantial and somewhat consistent catches of eggs are seen lower Zone 5, with sizable catches of larvae in upper Zone 5 and even into Zone 4. Only in 2004 were catches of Bay Anchovy larvae still appreciable upstream in Zone 3. This suggests upstream transport of Bay Anchovy larvae from principal spawning areas further downstream.

The results from New Jersey Seine Surveys are presented in Figure 12 for Bay Anchovy. Two important notes are warranted about this figure. First, because catches in Zone 3 were typically very low with many zero catches, data are presented for Zone 4, upper Zone 5 (RM 70 to 78.8), and lower Zone 5 (RM 48.2 to 70). Second, seine surveys for Bay Anchovy will necessarily capture relatively late stages of juveniles given their size range, and all catches will represent a mixture of juveniles and mature adults, some of whom are also young-of-year. Thus, the results presented in Figure 12 do not narrowly represent "propagation" success by focusing on early life stages. Supplemental length distribution data provided by NJ Bureau of Marine Fisheries help to demonstrate that juvenile fish were included in the Bay Anchovy catch, and that some nursery use exists for these early life stages.


The combined results from the Bay Anchovy ichthyoplankton and seine surveys primarily confirms the more intense and complete use of higher salinity areas of the Delaware Estuary for spawning and rearing activities. Nevertheless, the capture of eggs in lower Zone 5 and the capture of larvae relatively consistently in both Zone 4 and Zone 5 suggests a broader utilization of the Delaware Estuary for early life stages of the Bay Anchovy than simply the higher salinity waters of Delaware Bay (Zone 6). For Zone 3 of the estuary, the presence of larval fish in 2004 provides some ambiguity for the question, "Is there any evidence of successful reproduction?"

while it is clear that a negative answer is warranted in Zone 3 for the two higher demonstrations of early life stages. For Zone 4, the increased regularity of both larval fish catches and capture in the seine surveys leads to an affirmative response to the question, “Is there any evidence of successful reproduction?” Yet there remains uncertainty about how regular this use is, and it does not include the earlier spawning and egg stages of reproduction, leading to an ambiguous answer for the question, “Is there regular evidence of successful reproduction for this species?” and a negative response to the question, “Are there strong numbers and high consistency of successful reproduction?” In upper Zone 5, more regular larval catches indicate an affirmative answer to the questions, “Is there any evidence of successful reproduction?” and “Is there regular evidence of successful reproduction for this species?”, while the lack of eggs and the much higher catches downstream indicate a negative response to the question, “Are there strong numbers and high consistency of successful reproduction?” Finally in lower Zone 5, the regular presence of eggs and larvae combined with juvenile catches in the seine surveys demonstrates an affirmative answer to both of the first two “use” questions. However, comparisons to data from Delaware Bay also demonstrate that much stronger and more complete utilization occurs further down in the estuary and thus a negative response is warranted for the final question, “Are there strong numbers and high consistency of successful reproduction?”

Table 6. Evaluation Matrix for Bay Anchovy in Zones 3, 4, and 5 of the Delaware Estuary

		<b>Bay Anchovy</b>						
1979 Task Force PSEG target (12 spp) NJ: 10 K/yr and 10/yr groups	m	Common name	Scientific name	Any evidence of successful reproduction?	Regular evidence of successful reproduction?	Strong numbers & high consistency of successful reproduction?	"Nursery" with juveniles	
			Bay Anchovy	<i>Anchoa mitchilli</i>				
				<b>Zone 3</b>	?	N	N	Y
				<i>RM 95-108.4</i>				
				<b>Zone 4</b>	Y	?	N	Y
				<i>RM 78.8-95</i>				
				<b>Zone 5 (upper)</b>	Y	Y	N	Y
				<i>RM 70-78.8</i>				
				<b>Zone 5 (lower)</b>	Y	Y	N	Y
				<i>RM 48.2-70</i>				

## ATLANTIC SILVERSIDE



Atlantic Silverside (*Menidia menidia*) is also among the most abundant species in the Delaware Estuary (Wang and Kerhehan 1979). Atlantic Silverside are short-lived fish, with typical lifespans of 2 yrs or less (Fay et al. 1983). Although found from freshwater to saltwater zones, they are most common in the higher salinity waters of Delaware Bay and its associated tidal wetlands. Spawning occurs in intertidal areas in spring and summer, typically at salinities greater than 15 ppt, with repeat spawning occurring through the spawning season. For the current data evaluation, it is worth emphasizing that the intertidal spawning and adhesive eggs likely lead to poor representation of the eggs stage with open-water ichthyoplankton surveys such as the PSEG surveys of 2002, 2003, and 2004. Despite this limitation for eggs, larval Atlantic Silverside can be a dominant catch in ichthyoplankton samples (Able and Fahay 1998).

Atlantic Silverside results from the PSEG Ichthyoplankton surveys are presented in Figure 13 showing larvae in these three years largely confined to the lower portions of Zone 5 (RM 48.2 to RM 70), and eggs poorly represented in these open-water samples. Not shown in Figure 13 are the results further downstream in Delaware Bay, where larval abundances were typically as high or higher than those seen in lower Zone 5. As mentioned above, the lack of eggs in the ichthyoplankton collections is expected given spawning behaviors and egg characteristics. Overall, these results demonstrate nearly exclusive use of lower Zone 5 and seaward for reproduction and early life stages (where propagation is already a designated use), with little or no use of Zones 3 or 4 by these life stages.

New Jersey Seine Survey results are presented in Figure 14. Like with Bay Anchovy, catches of Atlantic Silverside were typically very low in Zone 3 and thus Figure 14 focuses on the patterns in Zone 4, upper Zone 5, and lower Zone 5. Also like Bay Anchovy, seine catches represent multiple overlapping year classes and stages of maturity, and cannot narrowly quantify the numbers of immature young-of-year fish in each sample. As a result, the results in Figure 14 do not capture patterns in “propagation” alone but more broadly assess habitat use by multiple life stages. Additional length distribution data from the NJ Bureau of Marine Fisheries, however, establishes that some young-of-year fish are using all zones of the Delaware Estuary, including upstream in Zone 2. Yet the seine data for Atlantic Silverside primarily indicate use of zones in the estuary with moderate to high salinities and sparse use of the freshwater and oligohaline areas.

The patterns from the ichthyoplankton and seine surveys indicate that the “propagation” use for Atlantic Silverside resides mainly in the high salinity zones of the estuary. For Zone 3 and Zone 4, no eggs or appreciable larval numbers indicate a negative response to all three matrix questions. For upper Zone 5, the sporadic larval fish catches leave ambiguous the response to even the basic question, “Is there any evidence of successful reproduction?” Thus, in upper Zone 5, the higher two matrix questions are negative with no “regular” nor “consistent” or “strong”


indications of early life stage use. Instead, these zones primarily are utilized as nursery areas by some small fraction of the juveniles as well as small numbers of adult fish. Only in lower Zone 5 do the data indicate an affirmative answer to the propagation questions, with both of the first use questions, “Is there any evidence of successful reproduction?” and “Is there regular evidence of successful reproduction for this species?” being supported with an affirmative answer from the regular larval fish catches. However, these catches in lower Zone 5 were at times less consistent and weaker than Delaware Bay catches, leading to an ambiguous assessment for the final question, “Are there strong numbers and high consistency of successful reproduction?”

Table 7. Evaluation Matrix for Atlantic Silverside in Zones 3, 4, and 5 of the Delaware Estuary

		<b>Atlantic Silverside</b>						
1979 Task Force PSEG target (12 spp) NJ: 10 K/yr and 10/yr groups		Common name	Scientific name		Any evidence of successful reproduction?	Regular evidence of successful reproduction?	Strong numbers & high consistency of successful reproduction?	"Nursery" with juveniles
	<b>m</b>		Atlantic Silverside	<i>Menidia menidia</i>				
				<b>Zone 3</b> RM 95-108.4	N	N	N	N
				<b>Zone 4</b> RM 78.8-95	N	N	N	?
				<b>Zone 5 (upper)</b> RM 70-78.8	?	N	N	?
				<b>Zone 5 (lower)</b> RM 48.2-70	Y	Y	?	Y



**ALEWIFE  
&  
BLUEBACK HERRING**



Alewife and Blueback Herring (*Alosa pseudoharengus* and *Alosa aestivalis*, respectively) are regularly considered and managed together as a pair of species referred to as River Herring. These two species can be somewhat difficult to distinguish in the field, and their behaviors and life histories broadly overlap. Like American Shad, the River Herring are anadromous fish which return to freshwater habitats from the ocean to spawn, although the exact preferences and localities for spawning of the two species continue to be researched and evaluated across their distributional range. For the current data review and assessment, these two species are likewise considered together because of the mixed and confounding patterns within the data sets that suggest possible confusion of the species or poor delineation of the timing and location of reproductive behaviors.

Results from the PSEG Ichthyoplankton surveys are presented in three figures (Figures 15, 16, and 17). These data show Alewife to be the dominant species spawning and rearing across many estuary zones, with Blueback Herring rarely detected in sampling, and the undetermined eggs/larvae of “*Alosa* spp.” mirroring the Alewife patterns. Results from the New Jersey Seine Survey are presented in Figures 18 and 19 for Alewife and Blueback Herring, respectively. These data represent primarily young-of-year production in the two species (some limited numbers of adults and age-1 fish are also caught), and are thus expected to provide some correspondence to the earlier life stages monitored via the ichthyoplankton surveys. Yet the patterns from the seine surveys reveal opposite dominance patterns, with Blueback Herring catches orders of magnitude higher than the Alewife catches. Moreover, field observations in the Delaware Estuary more closely match the results from the seine survey, with Blueback Herring frequently encountered while Alewife remain rare (M. Kaufman *personal communication*).

Multiple explanations could individually or collectively reconcile the apparent conflicting patterns from the two data sets. First, incorrect identifications could have reversed the species identity in either the ichthyoplankton or the seine surveys. Second, each sampling technique could be biased toward sampling one species preferentially over the other species. For instance, the daytime, near-shore seine sampling is well-suited for collecting young-of-year Striped Bass but likely under-samples schooling pelagic fish such as Alewife. Third, the spatial and temporal extents of each survey may lend themselves to increased efficiency in sampling one species over the other. These and other explanations highlight the challenge in distinguishing successful reproduction and rearing for each of these species, and underscore the decision by many

agencies and organizations to simply pool the two species as the broader River Herring complex despite important distinctions in their life histories.

Regardless of the reasons, the conflicting patterns in the ichthyoplankton and seine surveys severely limit the current assessment of “propagation” for each species. At the very least, eggs, larvae, and juveniles of both species have been identified in Zones 3 and 4 as well as the upper portion of Zone 5, leading to an affirmative response to the question, “Is there any evidence of successful reproduction?” The ichthyoplankton data further indicate moderate to strong spawning success in multiple years across Zones 2, 3, 4, and upper 5. However, it is not clear which of the two species (or perhaps both) are broadly utilizing the estuary for spawning and rearing. The assessment of the two higher “use” questions results is the ambiguous “?” for both species in Zones 3 and 4 because of these unresolved discrepancies between the two data sets, For upper Zone 5, weaker results regardless of species ID leads to an ambiguous answer to the question, “Is there regular evidence of successful reproduction for this species?” and a negative answer to the highest question, “Are there strong numbers and high consistency of successful reproduction?” For lower Zone 5, this initial question of “Is there any evidence of successful reproduction?” is answered in the affirmative because of moderate post-yolk sac larval catches in multiple years, while the remaining categories are not supported given the lack of eggs and inconsistencies in larval catches for either species.

Table 8. Evaluation Matrix for Alewife & Blueback Herring in Zones 3, 4, and 5 of the Delaware Estuary

1979 Task Force PSEG target (12 spp) NI: 10 K/yr and 10/yr groups		<b>Alewife &amp; Blueback Herring</b>		Any evidence of successful reproduction?	Regular evidence of successful reproduction?	Strong numbers & high consistency of successful reproduction?	"Nursery" with juveniles
		Common name	Scientific name				
m	Alewife	<i>Alosa pseudoharengus</i>	Zone 3	Y	?	?	Y
			RM 95-108.4				
			Zone 4	Y	?	?	Y
			RM 78.8-95				
			Zone 5 (upper)	Y	?	N	Y
			RM 70-78.8				
			Zone 5 (lower)	Y	N	N	Y
			RM 48.2-70				
m	Blueback Herring	<i>Alosa aestivalis</i>	Zone 3	Y	?	?	Y
			RM 95-108.4				
			Zone 4	Y	?	?	Y
			RM 78.8-95				
			Zone 5 (upper)	Y	?	N	Y
			RM 70-78.8				
			Zone 5 (lower)	Y	N	N	Y
			RM 48.2-70				

## ATLANTIC MENHADEN



Maryland DNR

Atlantic Menhaden (*Brevoortia tyrannus*) is another member of the Clupeidae family along with American Shad, Alewife, and Blueback Herring. Atlantic Menhaden adults are migratory coastal fish, with estuaries utilized as both nursery and seasonal adult habitats. Spawning primarily occurs in near-shore oceanic settings and can occur during both northward and southward migrations along the coast. Larval and juvenile Atlantic Menhaden then utilize estuaries as nursery grounds, with “ingress” into the Delaware Estuary peaking from December through May (Wang and Kernehan 1979). Juvenile fish continue to move further up-estuary into oligohaline and the lower freshwater portions of estuaries. Thus, both larval and juvenile Atlantic Menhaden (but not eggs/spawning) may use multiple zones within the Delaware Estuary.

For the current data evaluation, a limitation needs to be acknowledged for Atlantic Menhaden since the timing of both PSEG Ichthyoplankton and NJ Seine surveys only partially overlap with the period of larval and juvenile use of the Delaware Estuary. The current data sets can, therefore, only partly describe to use of the Delaware Estuary for “propagation” by Atlantic Menhaden.

Results from the PSEG Ichthyoplankton surveys are presented in Figure 20 for Atlantic Menhaden. Strong differences were seen among the three years surveyed, with 2002 showing a broad distribution of post-yolk sac larvae in many estuarine zones, while 2003 showed very low catches everywhere (low catches of larvae were also seen in Delaware Bay in 2003). In 2004, lower catches were observed than in 2002 but some utilization of Zone 5 was seen in addition to moderate catches in Delaware Bay (Zone 6).

Results from the New Jersey Seine Survey are shown in Figure 21 (overall catches very low in Zone 3; results from Zone 4 are presented along with separate analyses for upper and lower Zone 5). Juvenile Atlantic Menhaden were captured in all estuary zones, but only sporadically in Zones 2 and 3. In Zone 4, Atlantic Menhaden were not captured on a typical survey (i.e., median is zero for most years) but moderate to high catches are nevertheless somewhat common. In both upper and lower Zone 5, Atlantic Menhaden are more frequently encountered, although occasional very large catches (e.g., more than 1000 juvenile fish in a single seine haul) are seen at a frequency similar to Zone 4.

The combined results from the two surveys suggest some utilization of Zones 4 and 5 by larval and juvenile Atlantic Menhaden. Ichthyoplankton results were more variable, suggesting strong among-year difference in use (perhaps reflecting broader stock status or variable larval transport within the estuary). For Zone 3, low and inconsistent catches of larvae and juveniles provides only an ambiguous “?” answer to the first question, “Is there any evidence of successful reproduction?” and clear “no” answers for higher use questions. For Zone 4, increased

consistency and use across both larval and juveniles indicates “yes” for the initial question, “Is there any evidence of successful reproduction?” However, although juvenile catches were seen occasionally each year, moderate larval catches in Zone 4 only occurred in 2002, leading to an ambiguous answer to the question, “Is there regular evidence of successful reproduction for this species?” and a negative answer to the question “Are there strong numbers and high consistency of successful reproduction?” Finally, little or no distinction can be made between the results from upper Zone 5 and lower Zone 5, both in the ichthyoplankton surveys and the seine surveys. In addition, the larval catches in Zone 5 were among the highest seen throughout the estuary (including Zone 6) for both 2002 and 2004, even if large differences were seen among years. These results indicate that both upper and lower Zone 5 show regular Atlantic Menhaden use, and thus a “yes” to the questions, “Is there any evidence of successful reproduction?” and “Is there regular evidence of successful reproduction for this species?” However, both the inconsistency among years and the complex use of the estuary year-round without sufficient data during all seasons provides an ambiguous answer for the final question, “Are there strong numbers and high consistency of successful reproduction?”

Table 9. Evaluation Matrix for Atlantic Menhaden in Zones 3, 4, and 5 of the Delaware Estuary

<b>Atlantic Menhaden</b>							
1979 Task Force PSEG target (12 spp) NI: 10 K/yr and 10/yr groups		Common name	Scientific name	Any evidence of successful reproduction?	Regular evidence of successful reproduction?	Strong numbers & high consistency of successful reproduction?	"Nursery" with juveniles
	m		Atlantic Menhaden	<i>Brevoortia tyrannus</i>			
			<b>Zone 3</b> RM 95-108.4	?	N	N	?
			<b>Zone 4</b> RM 78.8-95	Y	?	N	Y
			<b>Zone 5 (upper)</b> RM 70-78.8	Y	Y	N	Y
			<b>Zone 5 (lower)</b> RM 48.2-70	Y	Y	N	Y

### ADDITIONAL SPECIES EVALUATED

A number of additional fish species use Zones 3, 4, and 5 as important parts of their spawning and/or nursery habitat. Data were evaluated for these additional species, but either the data were too sparse to provide any specific insight or the results were ambiguous and did not establish whether a given zone was being used for successful reproductive effort. For four species (Spot, Atlantic Croaker, Weakfish, Bluefish), the primary use centered on Delaware Bay (Zone 6) and catches in the upper zones of the estuary were so sparse that no clear indication of successful reproduction could be established. For Hogchoker, larval and juvenile use is expected throughout the estuary based on life history information, and Hogchokers were captured across all zones of the estuary. However data were not collected on larval fish, and juveniles could not be reliably identified in the seine surveys. As a result, the extent of use by Hogchokers in each zone could not be established. For Inland Silverside, a species similar to Atlantic Silverside but with a lower salinity preference, substantial use of Zones 3, 4, and 5 may be possible for spawning and rearing but no larval fish data were available and seine data could not separate mature from juvenile fish. Finally, Gizzard Shad were not directly assessed in the PSEG Ichthyoplankton surveys but may have been recorded in some of the broader taxonomic categories like “Clupeidae” undetermined. The NJ Seine Survey data also provided little specificity on the location and extent of juvenile use of the different estuary zones by Gizzard Shad.

Many more species of fish were captured and recorded as part of the New Jersey Seine Surveys. Shortnose Sturgeon data, for instance, were too sparse in any data set to assess this endangered fish species within this report. In addition, distribution and abundance data exist for species such as freshwater mussels and benthic invertebrates, but these data sets typically provide little specific information on the question of “successful reproduction”. As a result, although some data were available for many additional fish and non-fish species, these additional data sources were not used in the current evaluation of the Existing Use with respect to questions of “propagation”.

## **Conclusions**

Successful “propagation” in Zones 3, 4, and upper Zone 5 of the Delaware Estuary is clearly supported by the three primary data sources evaluated in this report. Thus, the goals for fishery restoration established in 1967 through DRBC’s designated uses have been exceeded, at least in part by the successful restoration of dissolved oxygen to 3.5 mg/L as a daily average concentration. The current designated use in Zone 3, 4, and the upper 8.8 miles of Zone 5 includes only “maintenance” of fish and other aquatic life; the “Existing Use” based on data collected since 2000 indicates that at least some degree of “propagation” (i.e., spawning and/or rearing of early stage larvae and juveniles) has been achieved in these 38 miles of the 133 mile long Delaware Estuary.

The current data evaluation also seeks to determine the strength of that “propagation” use across the suite of species for which specific relevant data were available. Because of different expectations for each species in the three estuary zones, each of these zones will be summarized separately.

In Zone 3 (Table 10), the evaluations for Atlantic Sturgeon, American Shad, and Striped Bass all fall short of the highest possible expectations for their recovery (e.g., see Appendix C). In particular, Atlantic Sturgeon catch rates and tracking data show sparse use of Zone 3. Both American Shad and Striped Bass exhibit moderate numbers and some consistency in eggs, larvae, and juveniles, but numbers are not as strong or as consistent as in other zones of the estuary. White Perch, on the other hand, shows strong numbers and high consistency in eggs, larvae, and juvenile use of Zone 3. Unfortunately, although some sizeable numbers exist for either Alewife or Blueback Herring, the ambiguity in these data sets precludes a more specific evaluation on the strength of these two species’ recovery in Zone 3. Finally, the three marine or brackish water species (Atlantic Menhaden, Bay Anchovy, Atlantic Silverside) are not expected to show extensive use of this freshwater zone of the estuary, and their weaker evaluation for Zone 3 does not reflect a failure in the “propagation” use. Overall, then, some moderate to strong recovery in “propagation” has been seen in Zone 3, but full “propagation” is not established.

The evaluation of Zone 4 (Table 11) is similar to Zone 3. Atlantic Sturgeon young-of-year have been caught in recent years in this zone, but overall numbers are small and year-to-year catches are highly sporadic. American Shad numbers also show substantial weakness in Zone 4, with a near absence of eggs in the 2002, 2003, and 2004 PSEG Ichthyoplankton surveys despite sizeable larval catches and similar habitats as upstream in Zone 3. Striped Bass numbers are moderately strong across eggs, larvae, and juveniles, but somewhat higher and more consistent data are seen further downstream in Zone 5 for eggs and larvae, indicating somewhat less use of Zone 4 than Zone 5 by Striped Bass for spawning and rearing. White Perch continue to show strong numbers in Zone 4 indicating a high degree of use of this zone, while both Alewife and Blueback Herring continue to be uncertain for Zone 4 because of ambiguity in the data sets. For the marine or brackish species, both Atlantic Menhaden and Bay Anchovy begin to show appreciable use of

Zone 4, which is not surprising given the beginning of the salinity transition in lower Zone 4 depending on the time of year and freshwater inflows. Together, these data (like in Zone 3) demonstrate moderate to strong recovery in “propagation” for Zone 4, but again full “propagation” has not been established.

In upper Zone 5 (RM70 to RM78.8; Table 12), some level of Atlantic Sturgeon spawning has been demonstrated, but the numbers and consistency again are weak and fall far short of attaining the full expectations for a successful population. By contrast, both White Perch and Striped Bass demonstrate strong numbers and high consistency across all early life stages examined. For both species, nearly complete or complete attainment of “propagation” is seen. For the three freshwater Clupeidae (American Shad, Alewife, and Blueback Herring), numbers begin to or strongly taper off in this zone where the transition to brackish water is more evident than upstream. These reductions in use for the three species may be a natural attenuation and therefore may not be a failure to attain an expected use in these 8.8 miles of the estuary, although appreciable ambiguity exists, particularly for Alewife and Blueback Herring (see Appendix C). Finally, the increased use by larval and juvenile Atlantic Menhaden and Bay Anchovy matches the life history expectations for these species. Across all species, then, the data for upper Zone 5 largely reflect moderate to strong attainment of expectations, with Atlantic Sturgeon numbers representing the weakest recovery and poorest demonstration of successful reproduction by species expected to use upper Zone 5. Thus, full “propagation” has not been established for upper Zone 5.

Finally, in lower Zone 5 (RM48.2 to RM70), both strength and weakness are seen in Table 13. However, in this lower portion of Zone 5, the designated use for these 22 miles of the estuary currently includes both “maintenance” and “propagation” and, indeed, the data evaluation reflects the more complete attainment of expected “propagation” for this estuary zone. Specifically, this reach of estuary represents the more distinct salinity transition, with only Striped Bass clearly expected to use lower Zone 5 as a core part of its spawning during early spring when freshwater inflows are typically higher and the salt line is shifted down-estuary. As Table 13 indicates, Striped Bass demonstrate strong and consistent evidence of successful reproduction here in lower Zone 5. Other species show less extensive use of lower Zone 5, but all of these shifts are expected based on their life history. The marine and brackish species primarily spawn either further down in Delaware Bay or the Atlantic Ocean, so the “regular” use of lower Zone 5 demonstrates a high degree of expected use. For the freshwater species, the tapering down of spawning and rearing in this salinity transition zone likewise matches the life history expectations for these species. Together, these data demonstrate a more complete use of lower Zone 5 for spawning and rearing in a manner consistent with each species’ life history.

A cautionary note is needed at this point. The evaluation of “propagation” in each estuary zone, and particularly the current attempt to identify the strength of any demonstrated spawning and rearing for these fish species, must not be confused with an evaluation of the causes for any failure to completely attain the expected patterns of spawning and rearing of these fish species. This report does not demonstrate, for instance, that water quality (e.g., dissolved oxygen) is the reason for the weaker-than-expected spawning and rearing numbers of any species, although

improvements to water quality have been a key component in the documented recovery of these fish populations (see Appendix B for a summary of contemporary dissolved oxygen data). Many factors, both natural and anthropogenic, continue to impact the populations of all fish species evaluated in this report (e.g., Iannuzzi et al. 2009). Most notable are the current population sizes for some species (e.g., Atlantic Sturgeon) that likely are too low to saturate the spawning and rearing grounds in any given year. But additional factors, such as habitat conditions and climate, likely play an important role in the strength of spawning and rearing for all species. The failure to demonstrate full attainment of the “propagation” use in Zones 3, 4, and upper Zone 5 may, therefore, reflect both limitations in water quality conditions as well as limitations in many other facets of each species’ biology. It is beyond the scope of this report to attempt to attribute the limitations in “propagation” to any one or any suite of causative agents.

The combined data sets evaluated for this report nevertheless indicate that the “Existing Use” attained within the Delaware Estuary in the period between 2000 and 2014 includes “propagation” for Zones 3, 4, and the upper 8.8 miles of Zone 5. In each of these zones, the evidence is moderate or even strong for one or more species, indicating substantial progress in attaining the full Clean Water Act aquatic life goal of “maintenance and propagation” for the species evaluated. The “Existing Use” for Zones 3, 4, and upper Zone 5 therefore substantially exceeds the “designated use” of only “maintenance”. But the data evaluated in this report also indicates that weakness in successful reproduction exists for one or more species for all three zones. Thus, full attainment of a “maintenance and propagation” use has not been demonstrated at this time based on the data available and examined for this existing use evaluation.



Table 10. Final Evaluation Matrix for Zone 3 across all species with sufficient data

<b>Zone 3 (RM 95 to RM 108.4)</b>							
1979 Task Force PSEG target (12 spp) NJ: 10 K/yr and 10/yr groups	Common name		Scientific name		Any evidence of successful reproduction?	Regular evidence of successful reproduction?	Strong numbers & high consistency of successful reproduction?
			Atlantic Sturgeon	<i>Acipenser oxyrinchus</i>		?	N
B			Striped Bass	<i>Morone saxatilis</i>	Y	Y	N
f			White Perch	<i>Morone americana</i>	Y	Y	Y
B			Alewife	<i>Alosa pseudoharengus</i>	Y	?	?
B			Blueback Herring	<i>Alosa aestivalis</i>	Y	?	?
B			American Shad	<i>Alosa sapidissima</i>	Y	Y	?
B			Atlantic Menhaden	<i>Brevoortia tyrannus</i>	?	N	N
B			Bay Anchovy	<i>Anchoa mitchilli</i>	?	N	N
B			Atlantic Silverside	<i>Menidia menidia</i>	N	N	N

Table 11. Final Evaluation Matrix for Zone 4 across all species with sufficient data

<b>Zone 4 (RM 78.8 to RM 95)</b>							
1979 Task Force PSEG target (12 spp) NJ: 10 K/yr and 10/yr groups	Common name		Scientific name		Any evidence of successful reproduction?	Regular evidence of successful reproduction?	Strong numbers & high consistency of successful reproduction?
			Atlantic Sturgeon	<i>Acipenser oxyrinchus</i>		Y	N
m			Striped Bass	<i>Morone saxatilis</i>	Y	Y	?
f			White Perch	<i>Morone americana</i>	Y	Y	Y
m			Alewife	<i>Alosa pseudoharengus</i>	Y	?	?
m			Blueback Herring	<i>Alosa aestivalis</i>	Y	?	?
m			American Shad	<i>Alosa sapidissima</i>	Y	?	N
m			Atlantic Menhaden	<i>Brevoortia tyrannus</i>	Y	?	N
m			Bay Anchovy	<i>Anchoa mitchilli</i>	Y	?	N
m			Atlantic Silverside	<i>Menidia menidia</i>	N	N	N

Table 12. Final Evaluation Matrix for upper Zone 5 across all species with sufficient data

<b>Zone 5 upper (RM 70 to RM 78.8)</b>								
1979 Task Force	PSEG target (12 spp)	NJ: 10 K/yr	and 10/yr groups	Common name	Scientific name	Any evidence of successful reproduction?	Regular evidence of successful reproduction?	Strong numbers & high consistency of successful reproduction?
								Atlantic Sturgeon
m				Striped Bass	<i>Morone saxatilis</i>	Y	Y	Y
f				White Perch	<i>Morone americana</i>	Y	Y	Y
m				Alewife	<i>Alosa pseudoharengus</i>	Y	?	N
m				Blueback Herring	<i>Alosa aestivalis</i>	Y	?	N
m				American Shad	<i>Alosa sapidissima</i>	Y	N	N
m				Atlantic Menhaden	<i>Brevoortia tyrannus</i>	Y	Y	N
m				Bay Anchovy	<i>Anchoa mitchilli</i>	Y	Y	N
m				Atlantic Silverside	<i>Menidia menidia</i>	?	N	N

Table 13. Final Evaluation Matrix for lower Zone 5 across all species with sufficient data

<b>Zone 5 lower (RM 48.2 to RM70)</b>								
1979 Task Force	PSEG target (12 spp)	NJ: 10 K/yr	and 10/yr groups	Common name	Scientific name	Any evidence of successful reproduction?	Regular evidence of successful reproduction?	Strong numbers & high consistency of successful reproduction?
								Atlantic Sturgeon
m				Striped Bass	<i>Morone saxatilis</i>	Y	Y	Y
f				White Perch	<i>Morone americana</i>	Y	?	N
m				Alewife	<i>Alosa pseudoharengus</i>	Y	N	N
m				Blueback Herring	<i>Alosa aestivalis</i>	Y	N	N
m				American Shad	<i>Alosa sapidissima</i>	N	N	N
m				Atlantic Menhaden	<i>Brevoortia tyrannus</i>	Y	Y	N
m				Bay Anchovy	<i>Anchoa mitchilli</i>	Y	Y	N
m				Atlantic Silverside	<i>Menidia menidia</i>	Y	Y	?

## **Acknowledgements**

This document was prepared by Delaware River Basin Commission staff: Steven J. Tambini, Executive Director. Erik L. Silldorff was the principal author. Reviews and improvements were provided by Tom Fikslin, John Yagecic, and Ron MacGillivray of the Delaware River Basin Commission. Great appreciation is extended to Russ Allen and Jen Pyle at the NJ Div. Fish & Wildlife, Bureau of Marine Fisheries, for sharing and clarifying the extraordinary data collected since the 1980s through the NJ Seine Survey. Ken Strait of PSEG likewise provided substantial assistance and insight for the PSEG Ichthyoplankton surveys. The Delaware River Basin Fish & Wildlife Management Cooperative technical committee assisted at many points during this effort and these colleagues never failed to provide guidance and support. The detailed consideration by Russ Allen, Hugh Carberry, Jen Pyle, and Heather Corbett (NJ Div. Fish & Wildlife, Bureau of Marine Fisheries); Mike Kauffman and Greg Murphy (PA Fish & Boat Commission); and Ed Hale and Ian Park (DE Fish & Wildlife) was invaluable to this data evaluation. Finally, discussions with Dewayne Fox (Del. State Univ.), Hal Brundage (Envir. Res. & Consult., Inc.), Ian Park (DE Fish & Wildlife), and Matt Fisher (formerly DE Fish & Wildlife) regarding Atlantic Sturgeon populations greatly improved the evaluation of Atlantic Sturgeon data.

## **References Cited**

- Able, K.W. and M.P. Fahay. 1998. *The First Year in the Life of Estuarine Fishes in the Middle Atlantic Bight*. Rutgers University Press, New Brunswick, NJ; 342 pp.
- ASMFC (Atlantic States Marine Fisheries Commission). 1995. Amendment #5 to the Interstate Fishery Management Plan for Atlantic Striped Bass. Fisheries Management Report No. 24 of the Atlantic States Marine Fishery Commission. 54 pp.
- ASMFC (Atlantic States Marine Fisheries Commission). 2013. Species Profile: Atlantic Striped Bass. Fisheries Focus 22(5): pgs. 4, 5, 13, 16.
- ASSRT (Atlantic Sturgeon Status Review Team). 2007. Status review of Atlantic sturgeon (*Acipenser oxyrinchus oxyrinchus*). Prepared by the ASSRT for the National Marine Fisheries Service, National Oceanic and Atmospheric Administration, July 27, 2007; 188 pp.
- Breece, M.W., M.J. Oliver, M.A. Cimino, and D.A. Fox. 2013. Shifting distributions of adult Atlantic Sturgeon amidst post-industrialization and future impacts in the Delaware River: a maximum entropy approach. PLoS ONE 8(11): e81321. doi: 10.1371 / journal.pone.0081321
- Brundage, H.M. and J.C. O'Herron. 2010. Acoustic telemetry studies of the distribution and movement of juvenile sturgeons in the Delaware River and Estuary. *Chapter 1 in* L. Calvo et al. Effects of flow dynamics, salinity, and water quality on the eastern oyster, the Atlantic sturgeon, and the shortnose sturgeon in the oligohaline zone of the Delaware Estuary. Report submitted to the United States Army Corps of Engineers. Seaboard Fisheries Institute. Penns Grove, NJ.
- Brundage, H.M. and J.C. O'Herron. 2014. Juvenile Atlantic Sturgeon movement and habitat use in the Delaware Estuary during 2009-2011 *Chapter 2 in* Bochenek et al. Effects of flow dynamics, salinity, and water quality on the eastern oyster, the Atlantic sturgeon, and the shortnose sturgeon in the oligohaline zone of the Delaware Estuary. Final Report Project Year 2011-2012 submitted to the United States Army Corps of Engineers. Seaboard Fisheries Institute. Penns Grove, NJ.
- Delaware 2014. Title 7 of Delaware Administrative Code: Section 7401 Surface Water Quality Standards. (revised October 11, 2014) {available at <http://regulations.delaware.gov/AdminCode/title7/7000/7400/7401.shtml>}
- DRBC (Delaware River Basin Commission). 1967. Section X of the Comprehensive Plan: Quality Standards for the Interstate Waters of the Delaware River Basin. 7 pp.

DRBC (Delaware River Basin Commission). 2013. Administrative Manual – Part III: Water Quality Regulations (with Amendments through December 4, 2013). Delaware River Basin Commission, West Trenton, New Jersey. 146 pp.

DRBFWMC (Delaware River Basin Fish & Wildlife Management Cooperative). 2011. Delaware River Sustainable Fishing Plan for American Shad. *prepared by* DE Div. Fish & Wildlife, NJ Div. Fish & Wildlife, PA Fish & Boat Comm., NYS Div. Fish & Wildlife, U.S. Fish & Wild. Serv., Nat. Mar. Fish. Serv. *for* Atlantic States Marine Fisheries Commission Shad and River Herring Management Board. 67 pp.

Fay, C.W., R.J. Neves, and G.B. Pardue. 1983. Atlantic Silverside - species profile: life histories and environmental requirements of coastal fishes and invertebrates (mid-Atlantic). U.S. Fish & Wildlife Service publication in cooperation with U.S. Army Corps of Engineers. 24 pp.

Fisher, M.T. 2010. State of Delaware Annual Compliance Report for Atlantic Sturgeon. Delaware Div. of Fish & Wild. submission to the Atlantic States Marine Fisheries Commission, Atlantic Sturgeon Plan Review Team; September 2010; 25 pp.

Fisher, M.T. 2011a. State of Delaware Annual Compliance Report for Atlantic Sturgeon. Delaware Div. of Fish & Wild. submission to the Atlantic States Marine Fisheries Commission, Atlantic Sturgeon Plan Review Team; September 2011; 23 pp.

Fisher, M.T. 2011b. Atlantic Sturgeon Final Report for State Wildlife Grant Project T-4-1. Delaware Div. Fish & Wild; January 2011; 44 pp.

Fisher, M.T. 2012. State of Delaware Annual Compliance Report for Atlantic Sturgeon. Delaware Div. of Fish & Wild. submission to the Atlantic States Marine Fisheries Commission, Atlantic Sturgeon Plan Review Team; September 2012; 30 pp.

Fisher, M.T. 2013. State of Delaware Annual Compliance Report for Atlantic Sturgeon. Delaware Div. of Fish & Wild. submission to the Atlantic States Marine Fisheries Commission, Atlantic Sturgeon Plan Review Team; September 2013; 26 pp.

FWPCA (Federal Water Pollution Control Administration) 1966. Delaware Estuary Comprehensive Study: Preliminary Report and Findings. U.S. Dept. of Interior, Federal Water Pollution Control Administration; Phila, PA. 124 pp.

FWPCA (Federal Water Pollution Control Administration) 1968. Water Quality Criteria. Report of the National Technical Advisory Committee to the Secretary of the Interior. U.S. Dept. of the Interior, Washington D.C. 215 pp. (*frequently referred to as the "Green Book" based on the cover binding color*)

- Gay, J. 1892. The shad streams of Pennsylvania. Rep. Penn. State Comm. Fish. 1889-90: 151-187.
- Hardy, C.A. 2009. Fish or foul: A history of the Delaware River basin through the perspective of the American Shad, 1682 to the Present. Penn. History 66(4): 506-534.
- Iannuzzi, T.J., J.L. Durda, D.V. Preziosi, D.F. Ludwig, R.G. Stahl Jr, A.A. DeSantis, and R.A. Hoke. 2009. Development of a preliminary relative risk model for evaluating regional ecological conditions in the Delaware River Estuary, USA. Int. Envir. Assess. Mgmt. 6(1): 164-179.
- Kahn, D.M., P. Himchak, C.A. Shirey, M. Fisher, T. Baum, R. Allen, G. Murphy, M. Kaufmann, H.E. Corbett, A. Linn, and R.W. Miller. 2015. Restoration of the Delaware River Spawning Stock of Striped Bass. *in M.Armstrong, G.Shepherd and P.Perra, (eds)*. Northeast Atlantic Coast Striped Bass Fisheries Management: Past, Present, and Future. American Fisheries Society, Bethesda, MD.
- McCord, J.W, M.R. Collins, W.C. Post, T.I.J. Smith. 2007. Attempts to develop an index of abundance for age-1 Atlantic Sturgeon in South Carolina, USA. American Fisheries Society Symposium #56: 397-403.
- Nadeau, R., R. Hemmett, J. Miller, R.W. Miller, D.M. Clark, D. Jacangelo, L. Skinner, R.W. Marshall, T.A. Strekal, C. Emery, S.D. Selzer, D.P. Pollison, R.C. Albert, R.C. Kausch, and A.A. Anderson. 1979. Dissolved Oxygen Requirements of a "Fishable" Delaware River Estuary. Report to the Delaware River Basin Commission by the Ad Hoc Task Force to Evaluate Dissolved Oxygen Requirements of Indigenous Estuary Fish. *{available online at [http://www.state.nj.us/drbc/library/documents/del-estuary DOrequirements 1979report.pdf](http://www.state.nj.us/drbc/library/documents/del-estuary_DOrequirements_1979report.pdf)}*
- Philadelphia 1914. Report on the collection and treatment of the sewage of the City of Philadelphia. City of Philadelphia (Pennsylvania, USA), Dept. of Public Works, Bureau of Surveys; G.S. Webster (chief eng.), G.E. Datesman (prin. asst. eng.), W.L. Stevenson (asst. eng.). 170 pp.
- PSEG (Public Service Enterprise Group). 2002. Biological Monitoring Program Annual Report-2002. 754 pp.
- PSEG (Public Service Enterprise Group). 2003. Biological Monitoring Program Annual Report-2003. 740 pp.
- PSEG (Public Service Enterprise Group). 2004. Biological Monitoring Program Annual Report-2004. 762 pp.

- Pyle, J. 2015. Studying the Delaware River – 2014 Report. NJ Div. Fish & Wildlife online report available at [www.nj.gov/dep/fgw/artdelstudy15.htm](http://www.nj.gov/dep/fgw/artdelstudy15.htm) {accessed 30-September-2015}
- Secor, D.H. and T.E. Gunderson. 1998. Effects of hypoxia and temperature on survival, growth, and respiration of juvenile Atlantic sturgeon, *Acipenser oxyrinchus*. Fishery Bulletin 96: 603-613.
- Simpson, P. C., and D. A. Fox. 2007. Atlantic sturgeon in the Delaware River: contemporary population status and identification of spawning areas. Completion Report: Award NA05NMF4051093, Dover, Delaware.
- Simpson, P.C. 2008. Movements and Habitat Use of Delaware River Atlantic Sturgeon. Masters Thesis; Delaware State Univ.; Dover, DE; 141 pp.
- Stier, D.J. and J.H. Crance. 1985. Habitat suitability index models and instream flow suitability curves: American Shad. U.S. Fish & Wild. Serv. Biol. Rep. 82(10.88). 47 pp.
- Wang, J.C.S. and R.J. Kernehan. 1979. Fishes of the Delaware Estuaries: a guide to the early life histories. E.A Communications; Towson, MD; 410 pp.
- Wright, J.F. and R. Porges. 1971. Water quality planning and management experiences of the Delaware River Basin Commission. Proceedings of the 5<sup>th</sup> International Water Pollution Research Conference; Pergamon Press Ltd; pgs. I-3/1 to I-3/17.

Figure 1. USGS Dissolved Oxygen Data for the sensor at the Ben Franklin Bridge (station 01467200) showing daily average, daily minimum, and daily maximum values from April 1, 1967, thru November 1, 1967 (data available at [waterdata.usgs.gov](http://waterdata.usgs.gov))






Figure 2. USGS Dissolved Oxygen Data for the sensor at the Ben Franklin Bridge (station 01467200) the distribution of 24-hour daily average values during the month of July from 1965 thru 2014 (data available at [waterdata.usgs.gov](http://waterdata.usgs.gov)).




Figure 3. Map of the Delaware Estuary and DRBC Water Quality Zones



Figure 4. Map of the Delaware Estuary with DRBC Water Quality Zones and PSEG survey “zones” or reaches for 2002-2004 Ichthyoplankton surveys




Figure 5. American Shad data summary from PSEG Ichthyoplankton surveys of 2002-2004 showing data distribution for all 8 surveys in each zone for each year as box-and-whisker plots (DRBC water quality zones are separated by dashed lines; PSEG survey reaches are indicated by blue bars at the top of each graph)




Figure 6. American Shad Young-of-Year data summary from New Jersey Seine Surveys for DRBC Water Quality Zones 3, 4, and 5. Full data distributions across all surveys within a Zone for a given year are presented as box-and-whisker plots.




Figure 7. Striped Bass data summary from PSEG Ichthyoplankton surveys of 2002-2004 showing data distribution for all 8 surveys in each zone for each year as box-and-whisker plots (DRBC water quality zones are separated by dashed lines; PSEG survey reaches are indicated by blue bars at the top of each graph)




Figure 8. Striped Bass Young-of-Year data summary from New Jersey Seine Surveys for DRBC Water Quality Zones 3, 4, and 5. Full data distributions across all surveys within a Zone for a given year are presented as box-and-whisker plots.




Figure 9. White Perch data summary from PSEG Ichthyoplankton surveys of 2002-2004 showing data distribution for all 8 surveys in each zone for each year as box-and-whisker plots (DRBC water quality zones are separated by dashed lines; PSEG survey reaches are indicated by blue bars at the top of each graph)






Figure 10. White Perch data summary from New Jersey Seine Surveys for DRBC Water Quality Zones 3, 4, and 5. Full data distributions across all surveys within a Zone for a given year are presented as box-and-whisker plots. (note: catches are for multiple year classes; these data do not solely represent young-of-year fish)




Figure 11. Bay Anchovy data summary from PSEG Ichthyoplankton surveys of 2002-2004 showing data distribution for all 8 surveys in each zone for each year as box-and-whisker plots (DRBC water quality zones are separated by dashed lines; PSEG survey reaches are indicated by blue bars at the top of each graph; higher salinity reaches of PSEG surveys with peak abundance for Bay Anchovy are not shown)




Figure 12. Bay Anchovy data summary from New Jersey Seine Surveys for DRBC Water Quality Zones 4 and 5, with data from the two sub-zones of Zone 5 presented separately. Full data distributions across all surveys within a Zone for a given year are presented as box-and-whisker plots. (note: catches are for both juvenile and mature fish across multiple year classes; these data do not solely represent immature fish)




Figure 13. Atlantic Silverside data summary from PSEG Ichthyoplankton surveys of 2002-2004 showing data distribution for all 8 surveys in each zone for each year as box-and-whisker plots (DRBC water quality zones are separated by dashed lines; PSEG survey reaches are indicated by blue bars at the top of each graph; higher salinity reaches of PSEG surveys with peak abundance for Bay Anchovy are not shown)




Figure 14. Atlantic Silverside data summary from New Jersey Seine Surveys for DRBC Water Quality Zones 4 and 5, with data from the two sub-zones of Zone 5 presented separately. Full data distributions across all surveys within a Zone for a given year are presented as box-and-whisker plots. (note: catches are for both juvenile and mature fish across multiple year classes; these data do not solely represent immature fish)




Figure 15. Alewife data summary from PSEG Ichthyoplankton surveys of 2002-2004 showing data distribution for all 8 surveys in each zone for each year as box-and-whisker plots (DRBC water quality zones are separated by dashed lines; PSEG survey reaches are indicated by blue bars at the top of each graph)




Figure 16. Blueback Herring data summary from PSEG Ichthyoplankton surveys of 2002-2004 showing data distribution for all 8 surveys in each zone for each year as box-and-whisker plots (DRBC water quality zones are separated by dashed lines; PSEG survey reaches are indicated by blue bars at the top of each graph)




Figure 17. “*Alosa* spp.” (indistinguishable Alewife & Blueback Herring eggs & larvae) data summary from PSEG Ichthyoplankton surveys of 2002-2004 showing data distribution for all 8 surveys in each zone for each year as box-and-whisker plots (DRBC water quality zones are separated by dashed lines; PSEG survey reaches are indicated by blue bars at the top of each graph)






Figure 18. Alewife data summary from New Jersey Seine Surveys for DRBC Water Quality Zones 3, 4, and 5. Full data distributions across all surveys within a Zone for a given year are presented as box-and-whisker plots. (note: primarily young-of-year fish, with some age-1 and adult fish at times collected)




Figure 19. Blueback Herring data summary from New Jersey Seine Surveys for DRBC Water Quality Zones 3, 4, and 5. Full data distributions across all surveys within a Zone for a given year are presented as box-and-whisker plots. (note: primarily young-of-year fish, with some age-1 and adult fish at times collected)




Figure 20. Atlantic Menhaden data summary from PSEG Ichthyoplankton surveys of 2002-2004 showing data distribution for all 8 surveys in each zone for each year as box-and-whisker plots (DRBC water quality zones are separated by dashed lines; PSEG survey reaches are indicated by blue bars at the top of each graph)





Figure 21 Atlantic Menhaden data summary from New Jersey Seine Surveys for DRBC Water Quality Zones 4 and 5, with data from the two sub-zones of Zone 5 presented separately. Full data distributions across all surveys within a Zone for a given year are presented as box-and-whisker plots. (note: primarily young-of-year fish, with some age-1 and adult fish at times collected)



# Appendix A. 1967 Water Quality Regulations & DRBC Resolution Approval

A RESOLUTION to amend the Comprehensive Plan with respect to water quality standards for the Delaware River Estuary and Bay.

WHEREAS, the Commission undertook a basinwide water pollution abatement and control program as authorized by its resolution of May 6, 1966 (No. 66-7), and as a first step in the formulation of that program, the Commission conducted a major conference on water pollution problems of the basin in Philadelphia on July 27, 1966; and

WHEREAS, the Commission has received and considered the preliminary report of the Delaware River Estuary Comprehensive Study conducted and prepared by the Federal Water Pollution Control Administration of the United States Department of the Interior, and the various findings and conclusions thereof; and

WHEREAS, a public hearing was duly held by the Commission on the 26th day of January, 1967, at Trenton, New Jersey, upon proposed water quality standards, as authorized or required by the Federal Water Pollution Control Act, as amended, and Articles 5 and 13 of the Delaware River Basin Compact, following the publication of notice pursuant to law; and

WHEREAS, the Commission has duly considered the results of its staff investigations, the facts and arguments presented at the public hearings, and all of the data and communications received by the Commission, and has consulted with interested parties in accordance with the Compact, and in consideration thereof the Commission has made certain determinations with respect to water quality standards for the Delaware River Estuary and Bay, and the tributaries thereof; and

WHEREAS, upon such investigations, public hearings and otherwise, the Commission finds and determines that the effectuation of national policy as set forth in the Water Pollution Control Act (Public Law 84-660) as last amended by the Clean Water Restoration Act of 1966 (P.L. 89-753), and of the purposes and provisions of the Delaware River Basin Compact and the Comprehensive Plan adopted pursuant thereto, requires that the Commission assume jurisdiction to control future pollution and abate existing pollution in the waters of the Delaware River Basin, by amendment of the Comprehensive Plan and by the adoption and promulgation of rules, regulations and standards pursuant to Article 5 of the Delaware River Basin Compact; now therefore

BE IT RESOLVED by the Delaware River Basin Commission:


1. The Comprehensive Plan is hereby amended by renumbering the existing Section X to Section XII and by the addition thereto of a new section, to be designated Section X Water Quality Standards, to read as follows:


Section X

Water Quality Standards

(The text of the standards is attached)

2. This resolution shall take effect immediately. All resolutions and parts of resolutions, and all parts of the Comprehensive Plan, heretofore adopted which are inconsistent with any provision of this resolution are, to the extent of such inconsistency, hereby repealed. This resolution will not apply to existing effluents pending the adoption by the Commission of rules and regulations to implement the Comprehensive Plan as amended, pursuant to Article 5 of the Compact.

  
Chairman

  
Secretary

ADOPTED: March 2, 1967

Amended by 67-7

QUALITY STANDARDS FOR THE INTERSTATE WATERS OF THE  
DELAWARE RIVER BASIN

ARTICLE I

ESTUARY GENERALLY

Water uses

Uses paramount. Water uses shall be paramount in determining stream quality objectives which, in turn, shall be the basis for determining effluent quality requirements.

Uses to be protected. The quality of waters shall be maintained in a safe and satisfactory condition for the following uses: (a) agricultural, industrial, and public water supplies after reasonable treatment, except where natural salinity precludes such uses; (b) wildlife, fish and other aquatic life; (c) recreation; (d) navigation; (e) controlled and regulated waste assimilation in reasonable compatibility with other uses; and (f) such other uses as may be provided by the Comprehensive Plan.

Stream quality objectives

Limits. The waters shall not contain substances attributable to municipal, industrial, or other discharges in concentrations or amounts sufficient to preclude the specified water uses to be protected. Within this requirement, the waters shall be substantially free from unsightly or malodorous nuisances due to floating solids, sludge deposits, debris, oil, scum, substances, in concentrations or combinations which are toxic or harmful to human, animal, plant, or aquatic life, or that produce color, taste, or odor.

Effluent quality requirements

Disinfection. Wastes (exclusive of stormwater bypass) containing human excreta shall be effectively disinfected before being discharged into surface bodies of water.

Public safety. Effluents shall not create a menace to public health or safety at the point of discharge.



Limits. Discharges shall not contain more than negligible amounts of debris, oil, scum or other floating materials, suspended matter which will settle to form sludge, toxic substances or substances that produce color, taste and odors.

#### Other considerations

Combined sewers. Any new facility or project combining sanitary or industrial waste with stormwater drainage, which would have a substantial effect on the quality of waters of the basin, shall not be permitted, whether or not any such project or facility discharges into an existing combined system.

Access and reports. The Commission, or its duly authorized representatives, shall have access, at reasonable hours, to observe and inspect waste treatment facilities and to collect samples for analyses. Upon written request, operation reports shall be submitted to the Commission.

Zones. The Delaware River and Bay may be divided into zones which will facilitate the management of surface and underground water quality. The required percentage reduction of biochemical oxygen demand of all wastes will be uniform within zones.

#### Definitions

Biochemical oxygen demand. Biochemical oxygen demand as determined under standard laboratory procedures for 5-days at 20° C.

Carbonaceous oxygen demand. That part of the ultimate oxygen demand associated with biochemical oxidation of carbonaceous, as distinct from nitrogenous, material.

Effective disinfection. The destruction of pathogenic organisms in such manner and under such controls as shall be prescribed by Commission regulations.

River mile. The distance, in statute miles, of a location or item measured from "mile zero" essentially along the centerline of the navigation channel. Mile zero is located at

intersection of a line between the Cape May Light and the tip of Cape Henlopen.

Application

This Article shall apply to the Delaware River Estuary and Bay, including the tidal portions of the tributaries thereof.

## ARTICLE II

### ZONE 2

#### Description

Zone 2 is that part of the Delaware River extending from the head of tidewater at Trenton, New Jersey, R.M. (River Mile) 133.4 (Trenton-Morrisville Toll Bridge) to R.M. 108.4 below the mouth of Pennypack Creek, including the tidal portions of the tributaries thereof.

#### Water uses

Uses to be protected. The quality of Zone 2 waters shall be maintained in a safe and satisfactory condition for the following uses: (a) agricultural, industrial, and public water supplies after reasonable treatment; (b) wildlife, maintenance and propagation of resident fish and other aquatic life, passage of anadromous fish; (c) recreation; and (d) navigation.

#### Stream quality objectives

Dissolved oxygen, daily average concentration shall not be less than 5.0 mg/l (milligrams per liter) except during the periods from April 1 to June 15 and September 16 to December 31 when the dissolved oxygen shall not average less than 6.5 mg/l; chlorides, maximum 15-day mean 50 mg/l; pH; between 6.5 and 8.5; total alkalinity, between 20 and 100 mg/l; hardness, maximum monthly mean 95 mg/l; temperature, shall not exceed 5° F above the average daily temperature gradient displayed during the 1961 - 1966 period, or a maximum of 86° F, whichever is less; phenols, maximum 0.005 mg/l; threshold odor number not to exceed 24 @ 60° C; syndets (synthetic detergents) measured as M.B.A.S. (methylene blue active substances), maximum monthly mean 0.5 mg/l; turbidity, maximum monthly mean 40 units, maximum 150 units; radioactivity, alpha emitters, maximum 3 pc/l (picocuries per liter), beta emitters, maximum 1,000 pc/l.

#### Effluent quality requirements

All discharges shall meet the effluent quality requirements of Article 1. The biochemical oxygen demand shall be reduced at all times by at least 85 percent. The carbonaceous oxygen demand from all outfalls in the Zone (exclusive of stormwater bypass) shall not exceed that assigned by Commission regulations.

## ZONE 3

### Description

Zone 3 is that part of the Delaware River extending from R. M. 108.4 to R. M. 95.0, below the mouth of Big Timber Creek, including the tidal portions of the tributaries thereof.

### Water uses

Uses to be protected. The quality of Zone 3 waters shall be maintained in a safe and satisfactory condition for the following uses: (a) agricultural, industrial, and public water supplies after reasonable treatment; (b) wildlife, maintenance of resident fish and other aquatic life, passage of anadromous fish; (c) recreation; and (d) navigation.

### Stream quality objectives

Dissolved oxygen, daily average concentration shall not be less than 3.5 mg/l, except during the periods from April 1 to June 15 and September 16 to December 31 when the dissolved oxygen shall not average less than 6.5 mg/l; chlorides, maximum 200 mg/l; pH, between 6.5 and 8.5; total alkalinity, between 20 and 120 mg/l; hardness, maximum monthly mean 150 mg/l; temperature, shall not exceed 5° F above the average daily temperature gradient displayed during the 1961 - 1966 period, or a maximum of 86° F, whichever is less; phenols, maximum 0.005 mg/l; threshold odor number not to exceed 24 @ 60° C; syndets (M.B.A.S.), maximum monthly mean 1.0 mg/l; turbidity, maximum monthly mean 40 units; maximum 150 units; radioactivity, alpha emitters, maximum 3 pc/l, beta emitters, maximum 1,000 pc/l.

### Effluent quality requirements

All discharges shall meet the effluent quality requirements of Article I. The biochemical oxygen demand shall be reduced at all times by at least 85 percent. The carbonaceous oxygen demand from all outfalls in the Zone (exclusive of stormwater bypass) shall not exceed that assigned by Commission regulations.

## ZONE 4

### Description

Zone 4 is that part of the Delaware River extending from R. M. 95.0 to R. M. 78.8, the Pennsylvania-Delaware boundary line, including the tidal portions of the tributaries thereof.

### Water uses

Uses to be protected. The quality of Zone 4 waters shall be maintained in a safe and satisfactory condition for the following uses: (a) industrial water supply after reasonable treatment; (b) wildlife, maintenance of resident fish and other aquatic life, passage of anadromous fish; (c) recreation; and (d) navigation.

### Stream quality objectives

Dissolved oxygen, daily average concentration shall not be less than 3.5 mg/l, except during the periods from April 1 to June 15 and September 16 to December 31 when the dissolved oxygen shall not average less than 6.5 mg/l; chlorides, maximum 250 mg/l at R. M. 92.47; pH, between 6.5 and 8.5; total alkalinity, between 20 and 120 mg/l; temperature, shall not exceed 5° F above the average daily temperature gradient displayed during the 1961 - 1966 period, or a maximum of 86° F, whichever is less; phenols, maximum 0.02 mg/l; threshold odor number not to exceed 24 @ 60° C; syndets (M.B.A.S.), maximum monthly mean 1.0 mg/l; turbidity, maximum monthly mean 40 units, maximum 150 units; radioactivity, alpha emitters, maximum 3 pc/l, beta emitters, maximum 1,000 pc/l.

### Effluent quality requirements

All discharges shall meet the effluent quality requirements of Article 1. The biochemical oxygen demand shall be reduced at all times by at least 85 percent. The carbonaceous oxygen demand from all outfalls in the Zone (exclusive of stormwater bypass) shall not exceed that assigned by Commission regulations.

## ZONE 5

### Description

Zone 5 is that part of the Delaware River extending from R. M. 78.8 to R. M. 48.2, Liston Point, including the tidal portions of the tributaries thereof.

### Water uses

Uses to be protected. The quality of Zone 5 waters shall be maintained in a safe and satisfactory condition for the following uses: (a) industrial water supply after reasonable treatment; (b) wildlife, maintenance of resident fish and other aquatic life, propagation of resident fish from R. M. 70.0 to R. M. 48.2, passage of anadromous fish; (c) recreation; and (d) navigation.

### Stream quality objectives

Dissolved oxygen, daily average concentration shall not be less than 3.5 mg/l at R. M. 78.8, 4.5 mg/l at R. M. 70.0 and 6.0 mg/l at R. M. 59.5, except during periods from April 1 to June 15 and September 16 to December 31 when the dissolved oxygen shall not average less than 6.5 mg/l in the entire zone; pH, between 6.5 and 8.5; total alkalinity, between 20 and 120 mg/l; temperature shall not exceed 5° F above the average daily temperature gradient displayed during the 1961 - 1966 period, or a maximum of 86° F, whichever is less; phenols, maximum 0.02 mg/l; threshold odor number not to exceed 24 @ 60° C; syndets (M.B.A.S.), maximum monthly mean 1.0 mg/l; turbidity, maximum monthly mean 40 units, maximum 150 units; radioactivity, alpha emitters, maximum 3 pc/l, beta emitters, maximum 1,000 pc/l.

### Effluent quality requirements

All discharges shall meet the effluent quality requirements of Article 1. The biochemical oxygen demand shall be reduced at all times by at least 85 percent. The carbonaceous oxygen demand from all outfalls in the Zone (exclusive of stormwater bypass) shall not exceed that assigned by Commission regulations.

## ZONE 6

### Description

Zone 6 is that part of Delaware Bay extending from R. M. 48.2 to R. M. 0.00, the Atlantic Ocean, including the tidal portions of the tributaries thereof.

### Water uses

Uses to be protected. The quality of Zone 6 waters shall be maintained in a safe and satisfactory condition for the following uses: (a) industrial water supply after reasonable treatment; (b) wildlife, maintenance and propagation of resident fish, shellfish, and other aquatic life, and passage of anadromous fish; (c) recreation and (d) navigation.

### Stream quality objectives

Dissolved oxygen, daily average concentration shall not be less than 6.0 mg/l; pH, between 6.5 and 8.5; total alkalinity, between 20 and 120 mg/l; temperature, shall not exceed 5° F above the average daily temperature gradient displayed during the 1961 - 1966 period, or a maximum of 86° F, whichever is less; phenols, maximum 0.02 mg/l; threshold odor number not to exceed 24 @ 60° C; syndets (M.B.A.S.), maximum monthly mean 1.0 mg/l; turbidity, maximum monthly mean 40 units, maximum 150 units; radioactivity, alpha emitters, maximum 3 pc/l, beta emitters, maximum 1,000 pc/l; coliform, MPN (most probable number) not to exceed U. S. Public Health Service's shellfish standards in designated shellfish areas.

### Effluent quality requirements

All discharges shall meet the effluent quality requirements of Article 1. The biochemical oxygen demand shall be reduced at all times by at least 85 percent.

Appendix B. Summary of Dissolved Oxygen  
Data from 2000 to 2014 from USGS Sensors  
at the Ben Franklin Bridge (01467200)  
and Chester, PA (01477050)

## Motivation for Dissolved Oxygen Data Summary

USEPA guidance on existing uses indicates that the description of an existing use should include both a description of the use itself as well as documentation of the water quality associated with, and protective of, that existing use (USEPA 2008<sup>1</sup>). The water quality in the Delaware Estuary consists of many complex and interacting components, from pH and dissolved oxygen, to turbidity and suspended sediments, to metals and toxic compounds, etc. It is beyond the scope of this Existing Use Evaluation to quantify the spatial and temporal variation in all water quality parameters for Zones 3, 4, and upper Zone 5. However, the historic hypoxia in the Delaware Estuary served as both the motivation for early estuary restoration efforts as well as regulatory actions such as DRBC's wasteload allocation of CBOD (aka, FSOD or First Stage Oxygen Demand). Moreover, the restoration in the fishery resources described in this Existing Use Evaluation could not have been attained without significant improvements in dissolved oxygen concentrations for the estuary. Because of the key link between this one facet of water quality and the restoration of the fishery resources, this appendix summarizes dissolved oxygen conditions for the same data window used in the fishery evaluation: 2000 to 2014. The cautions presented in the Conclusion section of this report are worth reiterating; the presentation of dissolved oxygen data does not presume that dissolved oxygen alone is responsible for the population variations, and either the recovery or the lack of recovery for any estuarine fish species.

### Dissolved Oxygen Data

The continuous water quality sensors maintained by the USGS and in part funded by the DRBC at the Ben Franklin Bridge and at Chester, PA, provide the most comprehensive documentation on the timing and magnitude of the dissolved oxygen conditions in Zones 3, 4 and the upper 8.8 miles of Zone 5. These sensors were established in the early 1960s and have been maintained every year for over 50 years, including the period 2000 to 2014 that serves as the basis for the fishery data evaluation of this report. The Ben Franklin Bridge sensor sits at the end of a pier on the Pennsylvania side of the river at RM 100 near the center of Zone 3. The Chester, PA, sensor is located at the Kimberly-Clark manufacturing facility at RM 83 near the lower limit of Zone 4. Each sensor records a suite of *in situ* water quality parameters at set intervals each day (currently every 15 minutes; see stations 01467200 and 01477050 at [waterdata.usgs.gov](http://waterdata.usgs.gov)) and the sensors are deployed in all months except winter (typically April through November).

As the graphs and tables included in this appendix demonstrate, the dissolved oxygen conditions in Zones 3 and 4 remain highly variable, with important changes seen among weeks, among months, and among years. To adequately capture the nature and details of these patterns of variation, full data distributions are presented across daily, monthly, and yearly time steps in either graphical or tabular form. Because DRBC's current dissolved oxygen criteria for these estuary zones rely on a 24-hour mean value, the focus for much of this appendix is on the patterns in the daily average dissolved oxygen concentration. But because of diel and tidal variations

---

<sup>1</sup> EPA 2008: September 5, 2008 letter from USEPA HQ (D. Keehner, Director, Standards and Health Protection Division) to Oklahoma Water Resources Board (D. Smithee) in response to Oklahoma Water Resources Board (D. Smithee, Chief, Water Quality Programs Division) letter and questions to USEPA-HQ (D. Keehner). {available at [water.epa.gov/scitech/swguidance/standards/upload/Smithee-existing-uses-2008-09-23.pdf](http://water.epa.gov/scitech/swguidance/standards/upload/Smithee-existing-uses-2008-09-23.pdf)}



within each day, and because aquatic organisms can be highly sensitive to minimum dissolved oxygen levels, data distributions are also presented for the daily minimum dissolved oxygen concentrations.

Although the data summaries themselves provide the best insight into the nature of dissolved oxygen variations from 2000 to 2014, a couple general statements can be made:

- Dissolved oxygen typically remains above the 3.5 mg/L criterion for most years, but both daily minima and 24-hour daily averages at times fall below 3.5 mg/L (more often at Ben Franklin Bridge than Chester);
- Dissolved oxygen typically is somewhat lower at the Ben Franklin Bridge site than the Chester site, but the reverse can be true;
- Patterns in high or low dissolved oxygen may be related to a number of contributing factors (e.g., freshwater inflow, water temperature, individual storm events) but the relative importance of these factors is not immediately evident when comparing among years

The following yearly summaries also provide narrative descriptions of key components of the variation within and among years at each of the two stations:

#### Conditions during 2000

- Ben Franklin daily average dissolved oxygen remained above 4.0 mg/L for most days during the summer, with 10% of days during September 2000 falling between 3.6 and 4.0 mg/L. Minimum dissolved oxygen at Ben Franklin remained above 3.5 mg/L for most of the summer, with a short period below 3.5 mg/L in September.
- Chester daily average dissolved oxygen remained above 4.3 mg/L during 2000, with higher concentrations earlier in the summer. Minimum dissolved oxygen fell below 4 mg/L on only one day during 2000, and remained between 4 mg/L and 5 mg/L for all of August and over half of September.

#### Conditions during 2001

- Ben Franklin daily average dissolved oxygen remained above 3.5 mg/L for the entire summer, but there were two periods (June, August) when mean dissolved oxygen fell below 4 mg/L. Minimum dissolved oxygen fell below 3.5 mg/L during those same two periods. Overall, however, dissolved oxygen remained above 4 mg/L (both minimum and daily average) for most of the summer.
- Chester daily average dissolved oxygen fell to 3.5 mg/L and remained between 3.5 mg/L and 4.0 mg/L for a period in June 2001, and fell to 3.7 mg/L during August 2001; between these low periods, daily average dissolved oxygen remained above 5 mg/L for most of the time. Minimum dissolved oxygen likewise was lowest during windows of reduced dissolved oxygen in June (3.1 mg/L) and August (3.2 mg/L).

#### Conditions during 2002

- Ben Franklin daily average dissolved oxygen fell below 3.5 mg/L on one day in August 2002. In addition, multiple periods of either mean or minimum dissolved oxygen below 4 mg/L were noted. Dissolved oxygen remained in the 4 mg/L to 5 mg/L range for a protracted period, extending through September 2002.
- Chester daily average dissolved oxygen remained above 5 mg/L for most days in 2002, with periods between 4 mg/L and 5 mg/L during June, July, and August. Minimum dissolved oxygen fell below 4 mg/L on one during 2002, and remained between 4 mg/L and 5 mg/L for much of the summer.

#### Conditions during 2003

- Ben Franklin dissolved oxygen concentrations remained relatively high throughout 2003, with the median of the daily averages remaining above 5 mg/L and the minimum dissolved oxygen never falling below 4 mg/L.
- Chester dissolved oxygen remained high through early July 2003, and then fell to more modest levels for the remainder of the summer. Daily average dissolved oxygen remained above 6 mg/L in June 2003, with median values of 5.2 mg/L in July and 4.6 mg/L in August. Daily minimum dissolved oxygen also remained above 6 mg/L in June 2003, but was lower in July, August, and early September.

#### Conditions during 2004

- Ben Franklin daily average dissolved oxygen remained above 3.5 mg/L and only ranged between 3.5 mg/L and 4 mg/L for a few days in late June and early July [*note: missing data in late July 2004 could be a period of reduced dissolved oxygen*]. Minimum dissolved oxygen fell as low as 3.2 mg/L in late June, but typically remained above 4 mg/L, and also remained above 5 mg/L for over half the days in June and August.
- Chester daily average dissolved oxygen fell below 4 mg/L during two periods in July and August 2004, but otherwise remained above 4 mg/L. Minimum dissolved oxygen also fell below 4 mg/L only during these same windows in July and August, and remained between 4 mg/L and 5 mg/L for most of June, July, and August.

#### Conditions during 2005

- Ben Franklin dissolved oxygen was moderately depressed in 2005. Daily average dissolved oxygen fell below 3.5 mg/L in July and remained between 3.5 mg/L and 4 mg/L for most days in both July and August. Minimum dissolved oxygen fell below 3.5 mg/L in June, July, August, and September, falling as low as 3.1 mg/L in July 2005.
- Chester daily average dissolved oxygen fell below 4 mg/L for four days in July 2005, and remained between 4 mg/L and 5 mg/L for most of the summer. Daily minimum dissolved oxygen fell as low as 3.4 mg/L on two days during July 2005 and had short periods in June and July below 4 mg/L; from June into September, daily minima typically ranged between 4 and 5 mg/L.

#### Conditions during 2006

- Ben Franklin dissolved oxygen remained moderately high during 2006 *[note: historic flooding occurred in the middle of summer 2006 on the non-tidal Delaware River; see [pubs.usgs.gov/of/2008/1203/pdf/ofr2008-1203.pdf](https://pubs.usgs.gov/of/2008/1203/pdf/ofr2008-1203.pdf)]*. Periods of low dissolved oxygen occurred in early June 2006 and in August 2006, but daily average dissolved oxygen remained above 4 mg/L for most days. Minimum dissolved oxygen fell below 3.5 mg/L briefly in August and remained between 3.5 mg/L and 4 mg/L on most days during August, but was above 4 mg/L for most days through the rest of summer.
- Chester daily average dissolved oxygen remained above 4 mg/L during 2006, with most days during summer 2006 above 5 mg/L. Daily minima dissolved oxygen typically ranged between 4 mg/L and 5 mg/L for June, July, and August 2006.

#### Conditions during 2007

- Ben Franklin dissolved oxygen was depressed in June 2007, although dissolved oxygen remained moderate for the rest of the summer *[note: too much missing data from early September precludes an accurate summary of September 2007 data]*. Average daily dissolved oxygen fell below 3.5 mg/L on five separate days in June 2007, and minimum dissolved oxygen fell to 2.5 mg/L. Minimum dissolved oxygen fell below 3.5 mg/L again at the end of August, at which point the USGS censored the data presumably because of sensor problems.
- Chester average daily dissolved oxygen remained above 5 mg/L for all of 2007 except a period during June when average dissolved oxygen fell between 4.7 mg/L and 5 mg/L. Minimum dissolved oxygen remained above 4.5 mg/L except for that same period in June 2007 when minimum dissolved oxygen fell between 4.2 mg/L and 4.5 mg/L.

#### Conditions during 2008

- Ben Franklin daily average dissolved oxygen remained above 3.5 mg/L during 2008, although periods of dissolved oxygen below 4 mg/L were seen in both June and July. Daily minimum dissolved oxygen fell as low as 2.2 mg/L in June and 2.7 mg/L in July, among the lowest dissolved oxygen values recorded during the 2000-2014 data window. Overall, dissolved oxygen remained between 4 mg/L and 5 mg/L for a protracted period, extending toward the end of September 2008.
- Chester daily average dissolved oxygen fluctuated around 5 mg/L during the summer of 2008, with the lowest daily average of 4.2 mg/L in July. Minimum dissolved oxygen readings typically were above 4 mg/L, with one day in June, two days in July, and two days in August with daily minima between 3.7 mg/L and 3.9 mg/L.

#### Conditions during 2009

- Ben Franklin dissolved oxygen concentrations remained relatively high for most of 2009, with only a single period in late July where dissolved oxygen fell substantially. Daily average dissolved oxygen remained above 6 mg/L for nearly all of June and through half of July, which was followed by a short duration where average dissolved oxygen fell to between 4 and 5 mg/L. Minimum dissolved oxygen likewise remained high for most of the summer, with minimum dissolved oxygen below 4 mg/L only during that short period in late July, when the lowest value of 3.3 mg/L was recorded.

- Chester dissolved oxygen remained notably high during 2009. Daily average dissolved oxygen exceeded 6.5 mg/L during most days this summer, with more typical values around 5 mg/L during only one period from late July into early August. Daily minima were also relatively high, often approaching or exceeding 6 mg/L, and again ranging in the more typical 4 mg/L to 5 mg/L range only during late July and early August.

#### Conditions during 2010

- Ben Franklin dissolved oxygen in 2010 is difficult to characterize because of equipment problems at this station. Real-time measurements in late June and early July indicated marked depressions in dissolved oxygen, but QA/QC checks showed the sensor was biased low in its readings during this period. Although spot measurements included dissolved oxygen readings as low as 3.3 mg/L, insufficient data exist during the period when the Ben Franklin sensor was inoperable to adequately and accurately characterize dissolved oxygen for 2010, although some depressed dissolved oxygen readings were recorded.
- Chester daily average dissolved oxygen exceeded 5 mg/L on most days during 2010 [*note: sensor failure in late June and early July occurred during a period when spot measurements showed depressed dissolved oxygen*]. Minimum dissolved oxygen remained above 4 mg/L on all days with approved data, and exceeded 5 mg/L on most days during the summer of 2010.

#### Conditions during 2011

- Ben Franklin dissolved oxygen remained relatively high through 2011, with a small depression in June and a more substantial drop in dissolved oxygen in late July and early August. Daily average dissolved oxygen remained above 5 mg/L for most of the year, and fell below 4 mg/L only on 2 days in August. Daily minimum dissolved oxygen fell to 3.4 mg/L on two days in August and remained between 4 and 5 mg/L through early August; minimum dissolved oxygen values were typically above 5 mg/L, however, during the remainder of 2011.
- Chester daily average dissolved oxygen remained above 5 mg/L for nearly every day during 2011, with just 5 days across June, July, and August with recorded average dissolved oxygen below 5 mg/L. Minimum dissolved oxygen remained above 4.2 mg/L during the summer, and stayed above 5 mg/L on most days during 2011.

#### Conditions during 2012

- Ben Franklin dissolved oxygen remained high in June 2012, but then fell to moderate to low values for the remainder of the summer. Daily average dissolved oxygen fell to 3.7 mg/L in July, and remained between 4 mg/L and 5 mg/L for all of August. Daily minimum dissolved oxygen fell to 3.1 mg/L in July, and remained between 3.5 and 4 mg/L for over half of August.
- Chester dissolved oxygen remained high through June 2012, and then fell to more typical values for July, August, and September. The median of the June daily averages was 6.7 mg/L, with medians in July, August, and September ranging from 4.9 to 5.5 mg/L. Minimum dissolved oxygen fell below 4 mg/L on 10 days during 2012, with the lowest value of 3.4 mg/L in August 2012.

#### Conditions during 2013


- Ben Franklin daily average dissolved oxygen remained relatively high throughout 2013, with only brief periods of daily average dissolved oxygen below 5 mg/L in August and September. Daily minimum dissolved oxygen remained above 4 mg/L for nearly all of the summer, falling below 4 mg/L on 2 days in September, with a minimum of 3.7 mg/L recorded that month.
- Chester daily average dissolved oxygen remained above 5 mg/L during the summer of 2013, with median or typical values approaching or exceeding 6 mg/L. Minimum dissolved oxygen remained above 5 mg/L for most days during 2013, falling to 4.9 mg/L on one day in July and as low as 4.7 mg/L on five days in August 2013.

#### Conditions during 2014

- Ben Franklin dissolved oxygen exhibited the highest sustained levels during 2014 compared to other years between 2000 and 2014, with daily average dissolved oxygen remaining above 5 mg/L for the only year during this period. Daily minimum dissolved oxygen also remained relatively high, with most minima at or above 6 mg/L through June and July, and the lowest value of 4.3 mg/L recorded in August.
- Official approved records for minimum dissolved oxygen during 2014 show a peculiar pattern, with dissolved oxygen concentrations as low as 3 mg/L despite the moderate to high daily average dissolved oxygen values. Official approved daily average dissolved oxygen remained near or above 5 mg/L. *[note: Closer inspection of the 15-minute interval data from USGS shows an unusual pattern of highly erratic dissolved oxygen readings during 2014. The dissolved oxygen data for 2014 are therefore so highly unusual that definitive statements on their distribution are tentative, as best.]*

BEN FRANKLIN BRIDGE (RM 100)  
DATA DISTRIBUTIONS


Distribution of USGS Dissolved Oxygen Data from Ben Franklin Bridge for the Year 2000



Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2000	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2000	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2000	Mar	0	<i>insufficient data to estimate percentiles and statistics</i>							
2000	Apr	30	9.2	9.3	9.5	9.7	9.7	9.8	10.0	10.1
2000	May	31	6.2	6.3	6.8	8.2	8.0	8.9	9.7	9.8
2000	June	25	5.9	6.6	6.9	7.9	7.6	8.1	8.3	8.8
2000	July	25	4.4	4.4	4.5	5.1	5.2	5.6	6.0	6.4
2000	Aug	29	4.3	4.4	4.4	4.6	4.7	5.0	5.1	5.2
2000	Sept	30	3.6	4.0	4.4	4.8	4.7	5.1	5.2	5.8
2000	Oct	31	4.8	4.9	5.6	5.8	5.9	6.4	6.6	7.4
2000	Nov	29	7.2	7.4	7.6	7.9	8.2	9.2	9.2	9.7
2000	Dec	3	<i>insufficient data to estimate percentiles and statistics</i>							

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2000	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2000	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2000	Mar	0	<i>insufficient data to estimate percentiles and statistics</i>							
2000	Apr	30	9.0	9.1	9.3	9.4	9.4	9.7	9.8	9.9
2000	May	31	5.7	6.0	6.4	7.9	7.7	8.6	9.4	9.7
2000	June	25	5.7	6.3	6.5	7.6	7.4	7.9	8.2	8.5
2000	July	25	3.7	3.9	4.3	4.8	4.8	5.2	5.7	6.1
2000	Aug	29	3.9	4.0	4.1	4.3	4.3	4.5	4.7	4.9
2000	Sept	30	3.2	3.5	4.1	4.3	4.3	4.7	4.8	5.4
2000	Oct	31	4.6	4.7	5.2	5.5	5.5	5.8	6.0	7.1
2000	Nov	29	6.9	7.1	7.3	7.6	7.9	8.9	9.1	9.3
2000	Dec	3	<i>insufficient data to estimate percentiles and statistics</i>							

Distribution of USGS Dissolved Oxygen Data from Ben Franklin Bridge for the Year **2001**




Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2001	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2001	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2001	Mar	4	<i>insufficient data to estimate percentiles and statistics</i>							
2001	Apr	30	9.3	9.5	9.7	10.1	10.2	10.9	11.2	11.2
2001	May	31	4.4	4.8	5.1	7.4	7.0	8.8	9.1	9.2
2001	June	30	3.6	3.7	4.0	5.8	6.0	8.1	8.4	8.7
2001	July	31	4.3	4.8	5.0	5.1	5.1	5.2	5.4	5.5
2001	Aug	31	3.5	3.8	4.5	4.7	4.8	5.3	5.5	5.7
2001	Sept	30	4.6	4.8	4.9	5.1	5.1	5.2	5.3	6.0
2001	Oct	31	5.7	6.0	6.1	6.3	6.4	6.7	7.1	7.3
2001	Nov	26	6.9	6.9	7.0	7.1	7.1	7.2	7.3	7.5
2001	Dec	3	<i>insufficient data to estimate percentiles and statistics</i>							

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2001	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2001	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2001	Mar	4	<i>insufficient data to estimate percentiles and statistics</i>							
2001	Apr	30	9.3	9.3	9.6	9.9	10.1	10.7	11.0	11.1
2001	May	31	3.8	4.3	4.7	6.6	6.5	8.5	8.9	9.0
2001	June	30	3.0	3.1	3.5	5.0	5.5	7.7	7.9	8.4
2001	July	31	3.8	4.5	4.6	4.8	4.7	4.9	5.1	5.2
2001	Aug	31	3.1	3.4	4.1	4.4	4.4	4.9	5.1	5.3
2001	Sept	30	4.3	4.6	4.7	4.8	4.8	4.9	4.9	5.7
2001	Oct	31	5.5	5.7	5.9	6.0	6.1	6.4	6.7	7.0
2001	Nov	26	6.2	6.7	6.7	6.8	6.8	6.9	7.0	7.2
2001	Dec	3	<i>insufficient data to estimate percentiles and statistics</i>							




Distribution of USGS Dissolved Oxygen Data from Ben Franklin Bridge for the Year 2002



Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2002	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2002	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2002	Mar	5	<i>insufficient data to estimate percentiles and statistics</i>							
2002	Apr	30	5.9	6.0	6.7	8.9	8.4	9.7	10.1	10.3
2002	May	28	7.0	7.4	7.7	8.0	7.9	8.1	8.3	8.6
2002	June	30	4.7	5.3	5.7	6.0	5.9	6.2	6.8	7.2
2002	July	31	3.9	4.0	4.2	4.4	4.5	4.7	5.1	5.3
2002	Aug	31	3.4	3.7	3.8	4.3	4.1	4.4	4.5	4.5
2002	Sept	30	4.3	4.3	4.6	4.7	4.7	4.8	4.9	5.0
2002	Oct	31	4.5	4.8	5.2	7.1	7.0	9.0	9.1	9.2
2002	Nov	30	9.4	9.6	9.6	9.9	10.1	10.7	10.9	11.1
2002	Dec	1	<i>insufficient data to estimate percentiles and statistics</i>							

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2002	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2002	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2002	Mar	5	<i>insufficient data to estimate percentiles and statistics</i>							
2002	Apr	30	5.5	5.9	6.4	8.7	8.2	9.6	10.0	10.2
2002	May	28	6.6	6.9	7.3	7.7	7.6	7.9	8.1	8.4
2002	June	30	4.4	5.0	5.4	5.7	5.6	5.8	6.5	7.0
2002	July	31	3.7	3.7	3.9	4.1	4.1	4.3	4.5	4.9
2002	Aug	31	3.2	3.3	3.6	4.0	3.8	4.1	4.2	4.3
2002	Sept	30	3.9	4.0	4.3	4.5	4.4	4.5	4.6	4.8
2002	Oct	31	4.2	4.4	4.8	6.5	6.7	8.8	8.9	9.0
2002	Nov	30	9.1	9.4	9.4	9.7	9.9	10.5	10.8	10.8
2002	Dec	1	<i>insufficient data to estimate percentiles and statistics</i>							


Distribution of USGS Dissolved Oxygen Data from Ben Franklin Bridge for the Year **2003**



Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2003	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2003	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2003	Mar	3	<i>insufficient data to estimate percentiles and statistics</i>							
2003	Apr	25	9.6	9.9	10.3	10.5	10.6	11.0	11.5	11.8
2003	May	24	7.0	7.4	7.7	7.9	8.1	8.4	9.4	10.0
2003	June	30	6.9	7.0	7.2	7.5	7.6	7.9	8.3	8.5
2003	July	26	4.7	4.8	4.8	5.1	5.3	5.5	6.4	7.2
2003	Aug	31	4.4	4.6	5.1	5.4	5.4	5.7	6.1	6.4
2003	Sept	30	4.4	4.8	5.8	6.5	6.2	6.8	6.9	7.0
2003	Oct	31	6.7	7.9	8.3	8.5	8.4	8.7	8.8	9.2
2003	Nov	30	8.9	9.1	9.3	9.8	9.8	10.3	10.3	10.8
2003	Dec	1	<i>insufficient data to estimate percentiles and statistics</i>							

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2003	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2003	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2003	Mar	3	<i>insufficient data to estimate percentiles and statistics</i>							
2003	Apr	25	9.4	9.7	10.1	10.4	10.4	10.6	11.3	11.7
2003	May	24	6.4	7.1	7.2	7.4	7.7	7.7	9.3	9.8
2003	June	30	6.4	6.7	6.9	7.1	7.3	7.8	8.0	8.3
2003	July	26	4.3	4.5	4.6	4.7	5.0	5.1	6.0	6.9
2003	Aug	31	4.0	4.2	4.6	5.0	5.0	5.3	5.6	5.9
2003	Sept	30	4.1	4.5	5.2	6.0	5.8	6.5	6.6	6.8
2003	Oct	31	5.1	7.7	8.1	8.3	8.2	8.5	8.6	8.9
2003	Nov	30	8.7	8.9	9.1	9.4	9.6	10.0	10.1	10.7
2003	Dec	1	<i>insufficient data to estimate percentiles and statistics</i>							


Distribution of USGS Dissolved Oxygen Data from Ben Franklin Bridge for the Year **2004**



Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2004	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2004	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2004	Mar	2	<i>insufficient data to estimate percentiles and statistics</i>							
2004	Apr	30	8.0	8.4	9.1	9.8	9.6	10.2	10.4	10.7
2004	May	31	5.3	5.8	6.2	7.3	7.1	8.1	8.3	8.4
2004	June	30	3.6	4.0	4.5	5.4	5.2	6.0	6.3	6.4
2004	July	22	3.9	4.0	4.3	4.6	4.7	5.0	5.1	6.4
2004	Aug	31	4.5	4.8	5.1	5.6	5.5	5.9	6.0	6.5
2004	Sept	28	4.9	5.3	5.6	6.0	6.2	6.9	7.5	7.7
2004	Oct	31	6.5	6.9	7.5	7.9	8.0	8.9	9.0	9.1
2004	Nov	27	8.5	8.8	8.9	9.0	9.1	9.2	9.6	9.9
2004	Dec	0	<i>insufficient data to estimate percentiles and statistics</i>							

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2004	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2004	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2004	Mar	2	<i>insufficient data to estimate percentiles and statistics</i>							
2004	Apr	30	7.8	8.0	8.8	9.5	9.3	10.1	10.1	10.6
2004	May	31	5.0	5.4	5.8	7.0	6.8	7.8	8.1	8.3
2004	June	30	3.2	3.6	4.1	5.0	4.8	5.7	5.7	5.9
2004	July	22	3.6	3.7	3.9	4.1	4.2	4.5	4.7	5.7
2004	Aug	31	4.1	4.4	4.6	5.3	5.1	5.6	5.7	5.9
2004	Sept	28	4.6	4.9	5.2	5.6	5.9	6.7	7.4	7.6
2004	Oct	31	6.3	6.7	7.3	7.7	7.7	8.5	8.7	8.8
2004	Nov	27	7.8	8.0	8.5	8.8	8.6	8.9	9.0	9.3
2004	Dec	0	<i>insufficient data to estimate percentiles and statistics</i>							


Distribution of USGS Dissolved Oxygen Data from Ben Franklin Bridge for the Year **2005**



Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2005	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2005	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2005	Mar	2	<i>insufficient data to estimate percentiles and statistics</i>							
2005	Apr	26	9.3	9.4	9.7	10.4	10.3	10.7	11.5	11.8
2005	May	31	5.5	5.9	6.6	8.6	8.2	9.8	9.8	10.0
2005	June	27	3.7	3.9	4.1	5.0	4.9	5.7	5.9	6.0
2005	July	30	3.4	3.5	3.6	3.8	3.8	4.0	4.1	4.4
2005	Aug	27	3.6	3.6	3.7	3.9	3.9	4.0	4.1	4.3
2005	Sept	30	3.6	3.7	4.0	4.5	4.3	4.6	4.7	4.7
2005	Oct	25	4.4	4.5	4.7	8.1	7.5	8.5	10.1	10.5
2005	Nov	28	8.7	8.9	9.0	9.5	9.7	10.4	10.9	11.0
2005	Dec	0	<i>insufficient data to estimate percentiles and statistics</i>							

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2005	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2005	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2005	Mar	2	<i>insufficient data to estimate percentiles and statistics</i>							
2005	Apr	26	9.2	9.3	9.5	10.3	10.2	10.5	11.3	11.7
2005	May	31	5.2	5.7	6.5	8.3	7.9	9.6	9.7	9.8
2005	June	27	3.4	3.7	3.8	4.3	4.5	5.4	5.5	5.7
2005	July	30	3.1	3.2	3.3	3.4	3.5	3.7	3.8	4.0
2005	Aug	27	3.3	3.4	3.5	3.6	3.6	3.7	3.9	4.0
2005	Sept	30	3.4	3.5	3.7	4.2	4.1	4.4	4.5	4.6
2005	Oct	25	4.2	4.3	4.5	8.0	7.2	8.2	9.9	10.3
2005	Nov	28	8.5	8.7	8.9	9.3	9.5	10.2	10.6	10.8
2005	Dec	0	<i>insufficient data to estimate percentiles and statistics</i>							


Distribution of USGS Dissolved Oxygen Data from Ben Franklin Bridge for the Year **2006**



Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2006	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2006	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2006	Mar	1	<i>insufficient data to estimate percentiles and statistics</i>							
2006	Apr	30	6.9	7.4	8.2	8.6	9.3	11.1	11.6	11.7
2006	May	27	6.2	6.5	6.9	7.3	7.2	7.7	7.7	7.8
2006	June	30	4.0	4.4	4.8	5.1	5.3	5.8	6.1	6.8
2006	July	31	4.4	4.6	5.0	5.9	5.7	6.4	6.6	7.0
2006	Aug	25	3.7	4.0	4.3	4.4	4.3	4.5	4.5	4.6
2006	Sept	30	4.7	5.0	5.2	6.1	5.9	6.4	6.8	7.0
2006	Oct	31	6.2	6.4	6.6	6.8	7.2	7.8	8.5	9.8
2006	Nov	30	8.4	8.5	8.7	9.7	9.5	9.9	10.6	10.6
2006	Dec	0	<i>insufficient data to estimate percentiles and statistics</i>							

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2006	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2006	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2006	Mar	1	<i>insufficient data to estimate percentiles and statistics</i>							
2006	Apr	30	6.4	6.9	7.8	8.4	9.0	10.8	11.4	11.5
2006	May	27	5.9	6.1	6.5	6.9	6.9	7.5	7.5	7.6
2006	June	30	3.7	4.2	4.5	4.8	4.9	5.1	5.7	6.6
2006	July	31	4.2	4.4	4.7	5.6	5.4	6.0	6.2	6.5
2006	Aug	25	3.2	3.4	3.8	3.9	3.8	4.0	4.1	4.2
2006	Sept	30	4.1	4.7	5.0	5.6	5.5	6.1	6.4	6.6
2006	Oct	31	5.5	5.9	6.1	6.4	6.8	7.5	8.3	9.4
2006	Nov	30	7.9	8.1	8.3	9.3	9.1	9.6	10.2	10.4
2006	Dec	0	<i>insufficient data to estimate percentiles and statistics</i>							

Distribution of USGS Dissolved Oxygen Data from Ben Franklin Bridge for the Year 2007




Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2007	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2007	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2007	Mar	1	<i>insufficient data to estimate percentiles and statistics</i>							
2007	Apr	30	9.1	9.4	10.6	11.0	10.7	11.1	11.3	11.5
2007	May	31	4.8	5.0	5.4	7.3	7.0	8.3	8.9	9.0
2007	June	30	3.2	3.4	3.7	4.0	4.3	5.0	5.3	5.4
2007	July	31	4.6	4.7	4.8	5.2	5.1	5.3	5.4	5.5
2007	Aug	26	4.1	4.4	4.6	4.6	4.6	4.7	4.8	4.8
2007	Sept <sup>a</sup>	13	<i>insufficient data to estimate percentiles and statistics</i>							
2007	Oct	31	5.6	5.7	5.9	6.4	6.5	6.9	7.8	8.4
2007	Nov	30	8.5	9.1	9.3	9.6	10.4	11.9	12.0	12.6
2007	Dec	18	12.6	12.7	12.9	13.3	13.2	13.4	13.6	13.7

<sup>a</sup> – sensor failure early in September 2007 could lead to bias in representing data distribution

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2007	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2007	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2007	Mar	1	<i>insufficient data to estimate percentiles and statistics</i>							
2007	Apr	30	8.7	9.0	10.3	10.7	10.4	10.8	11.0	11.2
2007	May	31	3.5	3.8	4.5	6.4	6.1	7.1	8.5	8.7
2007	June	30	2.5	2.8	2.9	3.3	3.5	3.9	4.7	4.8
2007	July	31	3.9	4.1	4.2	4.5	4.5	4.8	4.9	5.2
2007	Aug	26	3.0	3.7	4.0	4.2	4.1	4.3	4.3	4.4
2007	Sept <sup>a</sup>	13	<i>insufficient data to estimate percentiles and statistics</i>							
2007	Oct	31	5.3	5.4	5.5	6.1	6.2	6.4	7.2	8.1
2007	Nov	30	8.0	8.6	8.7	9.1	9.9	11.7	11.8	12.4
2007	Dec	18	12.2	12.5	12.7	12.9	12.9	13.2	13.3	13.4


Distribution of USGS Dissolved Oxygen Data from Ben Franklin Bridge for the Year **2008**



Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2008	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2008	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2008	Mar	0	<i>insufficient data to estimate percentiles and statistics</i>							
2008	Apr	30	7.7	8.8	9.8	10.3	10.4	11.4	11.8	12.0
2008	May	31	6.8	7.0	7.1	7.3	7.3	7.4	7.5	7.7
2008	June	30	3.7	3.8	4.0	5.3	5.6	7.2	7.5	7.9
2008	July	28	3.6	3.7	3.8	4.1	4.1	4.3	4.4	4.8
2008	Aug	31	4.2	4.3	4.5	4.7	4.6	4.8	4.9	5.0
2008	Sept	29	4.0	4.2	4.3	4.6	4.7	4.8	5.4	5.5
2008	Oct	31	5.4	5.7	5.9	6.0	6.4	6.9	7.3	9.1
2008	Nov	30	8.8	9.0	9.2	9.5	9.5	9.7	10.2	10.7
2008	Dec	17	10.7	11.3	11.5	11.6	11.6	11.7	12.1	12.2

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2008	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2008	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2008	Mar	0	<i>insufficient data to estimate percentiles and statistics</i>							
2008	Apr	30	6.5	7.9	9.1	9.6	9.8	11.0	11.4	11.6
2008	May	31	5.3	5.8	6.1	6.3	6.2	6.5	6.6	6.9
2008	June	30	2.2	2.6	3.0	4.1	4.3	5.8	6.1	6.7
2008	July	28	2.7	3.1	3.3	3.5	3.5	3.8	3.9	4.0
2008	Aug	31	3.5	3.8	4.1	4.2	4.2	4.4	4.5	4.7
2008	Sept	29	3.5	3.9	4.0	4.3	4.4	4.5	5.2	5.3
2008	Oct	31	4.9	5.4	5.6	5.7	6.1	6.5	6.9	8.9
2008	Nov	30	8.5	8.7	8.9	9.2	9.2	9.5	9.9	10.4
2008	Dec	17	10.4	10.9	11.1	11.3	11.3	11.4	11.8	12.1

Distribution of USGS Dissolved Oxygen Data from Ben Franklin Bridge for the Year **2009**




Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2009	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2009	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2009	Mar	0	<i>insufficient data to estimate percentiles and statistics</i>							
2009	Apr	29	7.8	8.4	9.0	9.3	9.2	9.5	10.0	10.4
2009	May	31	5.8	6.1	6.2	6.8	6.8	7.3	7.6	7.7
2009	June	30	5.6	5.7	6.0	6.5	6.5	7.0	7.3	7.4
2009	July	31	4.1	4.2	4.4	7.0	6.1	7.2	7.4	7.5
2009	Aug	31	5.0	5.2	5.4	5.5	5.7	6.1	6.3	6.5
2009	Sept	30	5.6	5.8	6.3	6.6	6.7	7.2	7.4	7.5
2009	Oct	31	6.8	6.9	7.3	7.5	7.8	8.5	8.8	9.3
2009	Nov	30	8.7	8.8	9.1	9.5	9.4	9.7	9.7	9.9
2009	Dec	0	<i>insufficient data to estimate percentiles and statistics</i>							

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2009	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2009	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2009	Mar	0	<i>insufficient data to estimate percentiles and statistics</i>							
2009	Apr	29	6.5	7.4	8.5	8.8	8.7	9.0	9.6	10.1
2009	May	31	4.8	5.1	5.3	6.0	5.9	6.2	6.6	7.1
2009	June	30	4.5	4.8	5.0	5.4	5.6	6.3	6.6	7.0
2009	July	31	3.3	3.5	3.6	6.2	5.4	6.5	6.8	7.1
2009	Aug	31	4.1	4.3	4.4	4.8	4.9	5.3	5.7	5.8
2009	Sept	30	4.8	5.2	5.8	6.0	6.1	6.7	7.0	7.2
2009	Oct	31	6.4	6.6	6.8	7.1	7.4	8.1	8.3	8.6
2009	Nov	30	8.2	8.5	8.7	8.9	8.9	9.1	9.3	9.5
2009	Dec	0	<i>insufficient data to estimate percentiles and statistics</i>							



Distribution of USGS Dissolved Oxygen Data from Ben Franklin Bridge for the Year **2010**




Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2010	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2010	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2010	Mar	0	<i>insufficient data to estimate percentiles and statistics</i>							
2010	Apr	30	8.6	8.7	8.9	9.3	9.6	9.9	11.3	11.7
2010	May	24	7.8	7.9	8.2	8.5	8.5	8.7	8.9	10.1
2010	June <sup>a</sup>	8	<i>insufficient data to estimate percentiles and statistics</i>							
2010	July <sup>a</sup>	0	<i>insufficient data to estimate percentiles and statistics</i>							
2010	Aug	25	4.4	4.6	4.7	4.8	4.8	5.0	5.1	5.2
2010	Sept	30	5.0	5.1	5.2	5.3	5.3	5.4	5.8	5.9
2010	Oct	31	5.8	7.0	7.9	8.0	7.9	8.2	8.3	8.6
2010	Nov	30	8.1	8.5	9.0	9.5	9.5	10.1	10.3	10.5
2010	Dec	0	<i>insufficient data to estimate percentiles and statistics</i>							

<sup>a</sup> – although sensor failed, spot measurements indicated D.O. as low as 3.3 mg/L during July 2010

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2010	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2010	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2010	Mar	0	<i>insufficient data to estimate percentiles and statistics</i>							
2010	Apr	30	8.0	8.1	8.3	8.7	9.0	9.4	10.9	11.5
2010	May	24	6.6	6.9	7.0	7.4	7.5	7.8	8.1	9.5
2010	June <sup>a</sup>	8	<i>insufficient data to estimate percentiles and statistics</i>							
2010	July <sup>a</sup>	0	<i>insufficient data to estimate percentiles and statistics</i>							
2010	Aug	25	4.1	4.3	4.4	4.5	4.5	4.7	4.7	4.9
2010	Sept	30	4.6	4.8	4.8	5.0	5.0	5.1	5.4	5.5
2010	Oct	31	5.2	6.8	7.4	7.6	7.5	7.9	7.9	8.2
2010	Nov	30	7.7	8.0	8.5	8.9	9.0	9.8	9.9	10.0
2010	Dec	0	<i>insufficient data to estimate percentiles and statistics</i>							


Distribution of USGS Dissolved Oxygen Data from Ben Franklin Bridge for the Year **2011**



Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2011	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2011	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2011	Mar	3	<i>insufficient data to estimate percentiles and statistics</i>							
2011	Apr	30	9.5	10.6	10.9	11.3	11.4	11.8	12.5	12.7
2011	May	31	8.1	8.2	8.5	8.7	9.0	9.6	9.7	9.9
2011	June	30	5.7	5.9	6.2	7.2	7.1	7.9	8.1	8.2
2011	July	31	4.2	4.6	5.7	6.6	6.2	6.9	7.0	7.5
2011	Aug	31	3.8	4.0	4.6	5.7	5.4	6.0	6.4	7.2
2011	Sept	30	7.3	7.4	7.6	7.8	7.8	8.1	8.2	8.3
2011	Oct	31	7.9	8.7	8.9	9.0	9.0	9.2	9.4	9.7
2011	Nov	30	9.9	10.4	10.6	10.9	10.8	11.0	11.0	11.1
2011	Dec	12	10.9	10.9	11.1	11.2	11.2	11.3	11.6	11.8

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2011	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2011	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2011	Mar	3	<i>insufficient data to estimate percentiles and statistics</i>							
2011	Apr	30	9.3	10.0	10.6	11.1	11.1	11.5	12.4	12.6
2011	May	31	7.7	7.9	8.1	8.5	8.7	9.4	9.5	9.6
2011	June	30	4.8	5.0	5.3	6.7	6.4	7.4	7.6	7.8
2011	July	31	3.7	4.0	5.1	6.1	5.7	6.6	6.6	7.1
2011	Aug	31	3.4	3.7	3.9	5.1	4.8	5.5	5.8	6.9
2011	Sept	30	7.0	7.2	7.3	7.5	7.5	7.7	7.9	8.0
2011	Oct	31	7.4	8.2	8.7	8.8	8.7	9.0	9.2	9.4
2011	Nov	30	9.6	10.1	10.3	10.6	10.5	10.7	10.8	10.8
2011	Dec	12	10.7	10.7	10.9	10.9	11.0	11.1	11.3	11.6


Distribution of USGS Dissolved Oxygen Data from Ben Franklin Bridge for the Year **2012**



Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2012	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2012	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2012	Mar	5	<i>insufficient data to estimate percentiles and statistics</i>							
2012	Apr	30	7.5	7.7	8.6	9.8	9.4	10.0	10.1	10.2
2012	May	31	6.7	7.2	7.2	7.5	7.6	8.0	8.1	8.2
2012	June	28	5.9	6.4	6.5	7.2	7.1	7.5	8.0	8.1
2012	July	31	3.7	3.8	3.9	4.4	4.4	4.9	5.2	5.5
2012	Aug	31	4.0	4.1	4.1	4.4	4.4	4.6	4.7	4.8
2012	Sept	30	4.2	4.3	4.6	5.2	5.4	6.4	6.8	7.0
2012	Oct	31	6.2	6.3	6.5	7.2	7.3	8.2	8.4	8.8
2012	Nov	30	8.6	9.1	10.0	10.3	10.1	10.4	10.5	10.6
2012	Dec	2	<i>insufficient data to estimate percentiles and statistics</i>							

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2012	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2012	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2012	Mar	5	<i>insufficient data to estimate percentiles and statistics</i>							
2012	Apr	30	7.3	7.4	8.1	9.5	9.0	9.6	9.8	9.8
2012	May	31	6.3	6.7	6.9	7.1	7.0	7.3	7.3	7.6
2012	June	28	4.9	5.7	5.8	6.3	6.2	6.6	6.9	7.2
2012	July	31	3.1	3.3	3.3	3.8	3.8	4.2	4.5	4.7
2012	Aug	31	3.5	3.6	3.8	3.9	3.9	4.1	4.1	4.2
2012	Sept	30	3.5	3.7	4.1	4.7	4.9	6.0	6.4	6.6
2012	Oct	31	5.6	5.8	6.0	6.7	6.9	8.0	8.2	8.6
2012	Nov	30	8.4	8.7	9.5	10.0	9.7	10.0	10.1	10.3
2012	Dec	2	<i>insufficient data to estimate percentiles and statistics</i>							


Distribution of USGS Dissolved Oxygen Data from Ben Franklin Bridge for the Year **2013**



Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2013	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2013	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2013	Mar	5	<i>insufficient data to estimate percentiles and statistics</i>							
2013	Apr	30	7.6	7.9	8.7	9.2	9.7	11.0	11.5	11.7
2013	May	31	6.2	6.4	6.7	7.7	7.9	9.0	9.9	10.1
2013	June	30	5.6	6.1	6.4	7.0	7.1	7.6	8.3	9.1
2013	July	31	5.0	5.1	5.3	5.8	5.9	6.3	6.8	7.2
2013	Aug	31	4.9	5.0	5.2	5.5	5.7	6.1	6.6	6.8
2013	Sept	30	4.6	4.9	4.9	5.3	5.2	5.5	5.6	5.7
2013	Oct	31	5.3	5.6	5.7	6.5	6.4	6.8	7.3	7.5
2013	Nov	30	7.3	7.7	7.9	9.0	8.8	9.1	9.8	10.7
2013	Dec	5	<i>insufficient data to estimate percentiles and statistics</i>							

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2013	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2013	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2013	Mar	5	<i>insufficient data to estimate percentiles and statistics</i>							
2013	Apr	30	7.1	7.5	8.6	9.0	9.3	10.8	11.1	11.4
2013	May	31	5.8	6.0	6.3	7.2	7.4	8.2	9.4	9.7
2013	June	30	4.9	5.6	5.8	6.6	6.5	7.3	7.6	8.3
2013	July	31	4.2	4.4	4.8	5.3	5.3	5.8	6.2	6.5
2013	Aug	31	4.1	4.2	4.5	4.7	4.9	5.4	5.8	6.1
2013	Sept	30	3.7	4.2	4.3	4.6	4.6	5.0	5.1	5.1
2013	Oct	31	4.0	4.7	5.0	5.8	5.6	6.1	6.5	6.9
2013	Nov	30	6.5	7.1	7.5	8.5	8.3	8.7	9.4	10.1
2013	Dec	5	<i>insufficient data to estimate percentiles and statistics</i>							

Distribution of USGS Dissolved Oxygen Data from Ben Franklin Bridge for the Year **2014**




Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2014	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2014	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2014	Mar	5	<i>insufficient data to estimate percentiles and statistics</i>							
2014	Apr	30	9.4	9.6	9.9	10.4	10.7	11.3	12.0	12.3
2014	May	31	6.9	7.4	7.8	8.3	8.3	9.0	9.2	9.7
2014	June	30	6.1	6.4	6.9	7.4	7.4	8.2	8.4	8.7
2014	July	31	5.7	6.1	6.6	7.2	7.0	7.5	7.6	7.9
2014	Aug	31	5.1	5.1	5.2	5.2	5.4	5.5	5.6	6.1
2014	Sept	30	5.1	5.2	5.2	5.3	5.5	5.8	6.1	6.2
2014	Oct	31	5.8	5.9	6.2	6.5	6.7	7.3	7.5	7.6
2014	Nov	30	7.6	8.1	8.3	8.7	9.2	10.5	10.6	10.9
2014	Dec	3	<i>insufficient data to estimate percentiles and statistics</i>							

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2014	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2014	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2014	Mar	5	<i>insufficient data to estimate percentiles and statistics</i>							
2014	Apr	30	9.1	9.2	9.6	10.0	10.3	10.9	11.7	11.9
2014	May	31	6.4	7.0	7.3	7.8	7.9	8.7	9.0	9.4
2014	June	30	5.3	5.9	6.2	6.8	6.8	7.6	7.9	8.2
2014	July	31	4.5	5.1	5.7	6.1	5.9	6.3	6.6	7.0
2014	Aug	31	4.3	4.4	4.4	4.6	4.6	4.8	5.0	5.2
2014	Sept	30	4.4	4.5	4.7	4.9	5.0	5.2	5.6	5.9
2014	Oct	31	5.3	5.5	5.8	6.0	6.2	6.9	7.1	7.2
2014	Nov	30	7.3	7.8	7.9	8.3	8.8	10.0	10.2	10.6
2014	Dec	3	<i>insufficient data to estimate percentiles and statistics</i>							

CHESTER, PA (RM 83)  
DATA DISTRIBUTIONS


Distribution of USGS Dissolved Oxygen Data from Chester, PA, for the Year **2000**



Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2000	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2000	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2000	Mar	0	<i>insufficient data to estimate percentiles and statistics</i>							
2000	Apr	12	8.7	8.8	9.1	9.1	9.1	9.2	9.5	9.6
2000	May	29	5.4	6.1	6.7	7.8	7.6	8.4	9.2	9.3
2000	June	30	5.0	5.3	5.6	6.6	6.7	8.0	8.4	8.6
2000	July	31	4.4	4.8	5.2	5.4	5.5	5.8	6.2	6.6
2000	Aug	31	4.4	4.5	4.6	4.9	4.8	5.1	5.2	5.3
2000	Sept	30	4.3	4.5	4.6	5.0	5.0	5.3	5.6	5.8
2000	Oct	31	5.1	5.2	5.4	5.6	5.7	5.9	6.2	7.2
2000	Nov	5	<i>insufficient data to estimate percentiles and statistics</i>							
2000	Dec	0	<i>insufficient data to estimate percentiles and statistics</i>							

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2000	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2000	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2000	Mar	0	<i>insufficient data to estimate percentiles and statistics</i>							
2000	Apr	12	8.5	8.6	8.9	8.9	8.9	9.0	9.2	9.5
2000	May	29	5.0	5.5	6.3	7.4	7.3	8.3	9.0	9.1
2000	June	30	4.2	4.8	5.1	6.1	6.3	7.5	7.9	8.1
2000	July	31	4.1	4.3	4.9	5.1	5.1	5.4	5.8	6.2
2000	Aug	31	4.0	4.1	4.3	4.5	4.5	4.8	4.9	5.0
2000	Sept	30	3.7	4.1	4.3	4.5	4.6	5.0	5.2	5.5
2000	Oct	31	4.8	4.9	5.1	5.2	5.4	5.6	5.7	7.0
2000	Nov	5	<i>insufficient data to estimate percentiles and statistics</i>							
2000	Dec	0	<i>insufficient data to estimate percentiles and statistics</i>							

Distribution of USGS Dissolved Oxygen Data from Chester, PA, for the Year **2001**




Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2001	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2001	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2001	Mar	4	<i>insufficient data to estimate percentiles and statistics</i>							
2001	Apr	26	9.2	9.2	9.4	9.9	9.8	10.3	10.5	10.6
2001	May	30	5.2	5.8	6.2	6.8	7.3	9.0	9.2	9.5
2001	June	28	3.5	3.6	3.7	4.9	5.4	7.2	7.9	8.5
2001	July	31	4.5	5.1	5.2	5.5	5.8	6.4	6.9	7.1
2001	Aug	29	3.7	3.9	4.3	4.9	4.7	5.2	5.4	5.6
2001	Sept	25	4.9	5.1	5.3	5.4	5.4	5.5	5.5	6.0
2001	Oct	31	6.1	6.2	6.8	6.9	6.9	7.2	7.6	7.7
2001	Nov	30	7.4	7.5	7.6	8.1	8.1	8.4	8.5	8.6
2001	Dec	2	<i>insufficient data to estimate percentiles and statistics</i>							

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2001	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2001	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2001	Mar	4	<i>insufficient data to estimate percentiles and statistics</i>							
2001	Apr	26	8.8	8.9	9.1	9.6	9.6	10.0	10.3	10.5
2001	May	30	4.9	5.3	5.6	6.5	6.8	8.3	8.8	8.9
2001	June	28	3.1	3.2	3.4	4.5	5.0	6.7	7.0	7.7
2001	July	31	3.8	4.6	4.8	5.2	5.3	5.9	6.3	6.9
2001	Aug	29	3.2	3.5	3.8	4.4	4.2	4.7	4.7	5.0
2001	Sept	25	4.5	4.6	4.8	4.9	4.9	5.0	5.1	5.5
2001	Oct	31	5.5	5.8	6.2	6.6	6.5	6.7	7.0	7.3
2001	Nov	30	7.0	7.1	7.2	7.9	7.7	8.1	8.2	8.3
2001	Dec	2	<i>insufficient data to estimate percentiles and statistics</i>							




Distribution of USGS Dissolved Oxygen Data from Chester, PA, for the Year **2002**



Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2002	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2002	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2002	Mar	5	<i>insufficient data to estimate percentiles and statistics</i>							
2002	Apr	30	6.2	6.3	7.0	9.0	8.5	9.7	9.9	10.1
2002	May	10	<i>insufficient data to estimate percentiles and statistics</i>							
2002	June	30	4.4	4.5	4.7	5.5	5.7	6.6	7.4	7.4
2002	July	31	4.5	4.6	5.0	5.2	5.1	5.3	5.5	5.5
2002	Aug	28	4.7	4.9	5.1	5.3	5.3	5.4	5.6	5.8
2002	Sept	30	5.8	5.9	6.0	6.1	6.1	6.3	6.4	6.6
2002	Oct	31	5.7	5.9	6.1	6.5	6.9	7.9	8.2	8.3
2002	Nov	30	8.4	8.7	8.7	9.0	9.3	9.9	10.2	10.6
2002	Dec	0	<i>insufficient data to estimate percentiles and statistics</i>							

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2002	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2002	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2002	Mar	5	<i>insufficient data to estimate percentiles and statistics</i>							
2002	Apr	30	6.0	6.2	6.6	8.7	8.3	9.6	9.8	9.9
2002	May	10	<i>insufficient data to estimate percentiles and statistics</i>							
2002	June	30	4.2	4.2	4.4	5.4	5.4	6.3	7.1	7.3
2002	July	31	3.9	4.2	4.5	4.7	4.6	4.8	5.0	5.2
2002	Aug	28	4.0	4.3	4.6	4.8	4.8	4.9	5.1	5.4
2002	Sept	30	5.2	5.4	5.5	5.7	5.7	5.9	6.0	6.3
2002	Oct	31	5.3	5.4	5.6	6.1	6.5	7.5	8.0	8.2
2002	Nov	30	8.3	8.4	8.6	8.9	9.1	9.6	9.9	10.3
2002	Dec	0	<i>insufficient data to estimate percentiles and statistics</i>							


Distribution of USGS Dissolved Oxygen Data from Chester, PA, for the Year **2003**



Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2003	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2003	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2003	Mar	5	<i>insufficient data to estimate percentiles and statistics</i>							
2003	Apr	30	9.0	9.4	9.8	10.3	10.2	10.8	10.9	11.0
2003	May	25	5.5	6.2	6.6	6.9	7.0	7.3	7.9	8.7
2003	June	30	6.1	6.4	6.7	6.9	6.9	7.2	7.4	7.5
2003	July	31	4.5	4.6	4.8	5.2	5.6	6.7	7.2	7.3
2003	Aug	31	4.2	4.3	4.5	4.6	4.6	4.9	5.1	5.2
2003	Sept	30	3.8	4.4	4.8	5.9	5.6	6.2	6.4	7.0
2003	Oct	31	6.6	7.3	7.4	7.7	7.7	7.9	8.2	9.1
2003	Nov	30	8.2	8.3	8.5	9.1	9.0	9.3	9.4	10.3
2003	Dec	1	<i>insufficient data to estimate percentiles and statistics</i>							

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2003	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2003	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2003	Mar	5	<i>insufficient data to estimate percentiles and statistics</i>							
2003	Apr	30	8.8	9.2	9.6	10.2	10.1	10.7	10.8	10.9
2003	May	25	5.2	6.1	6.2	6.5	6.6	6.9	7.4	7.9
2003	June	30	6.0	6.3	6.5	6.7	6.7	7.1	7.2	7.4
2003	July	31	4.3	4.4	4.5	4.9	5.3	6.4	6.8	7.0
2003	Aug	31	3.9	4.1	4.2	4.4	4.4	4.7	4.9	5.0
2003	Sept	30	3.6	3.9	4.4	5.8	5.3	6.0	6.3	6.6
2003	Oct	31	6.5	7.0	7.2	7.4	7.5	7.6	8.0	8.8
2003	Nov	30	8.1	8.2	8.4	8.9	8.8	9.2	9.3	10.1
2003	Dec	1	<i>insufficient data to estimate percentiles and statistics</i>							


Distribution of USGS Dissolved Oxygen Data from Chester, PA, for the Year **2004**



Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2004	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2004	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2004	Mar	2	<i>insufficient data to estimate percentiles and statistics</i>							
2004	Apr	30	7.8	7.9	8.3	10.3	9.7	10.7	11.0	11.4
2004	May	30	4.9	5.1	5.6	7.1	6.5	7.4	7.7	7.8
2004	June	30	4.6	4.8	4.9	5.1	5.1	5.4	5.5	5.9
2004	July	30	3.9	3.9	4.3	4.6	4.6	4.9	5.1	5.2
2004	Aug	29	3.8	4.0	4.3	5.0	4.9	5.4	5.5	5.7
2004	Sept	28	4.7	4.9	5.1	5.4	5.8	6.8	7.2	7.3
2004	Oct	17	6.7	6.7	6.7	6.9	7.0	7.1	7.4	7.6
2004	Nov	30	7.0	7.2	7.7	8.0	8.0	8.4	8.6	9.4
2004	Dec	9	<i>insufficient data to estimate percentiles and statistics</i>							

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2004	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2004	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2004	Mar	2	<i>insufficient data to estimate percentiles and statistics</i>							
2004	Apr	30	7.6	7.8	8.1	10.2	9.5	10.5	10.9	11.2
2004	May	30	4.6	4.7	5.3	6.8	6.2	7.2	7.4	7.6
2004	June	30	4.2	4.4	4.6	4.8	4.9	5.2	5.2	5.8
2004	July	30	3.7	3.8	4.0	4.3	4.3	4.6	4.8	5.0
2004	Aug	29	3.6	3.8	4.2	4.7	4.7	5.2	5.3	5.5
2004	Sept	28	4.5	4.6	4.9	5.2	5.6	6.7	7.0	7.2
2004	Oct	17	6.4	6.5	6.6	6.8	6.8	7.0	7.2	7.5
2004	Nov	30	6.8	7.1	7.6	7.8	7.8	8.2	8.3	9.2
2004	Dec	9	<i>insufficient data to estimate percentiles and statistics</i>							


Distribution of USGS Dissolved Oxygen Data from Chester, PA, for the Year **2005**



Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2005	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2005	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2005	Mar	2	<i>insufficient data to estimate percentiles and statistics</i>							
2005	Apr	30	7.5	7.9	8.3	9.2	9.1	10.0	10.2	10.4
2005	May	31	5.6	5.7	6.1	6.9	6.7	7.3	7.5	7.6
2005	June	21	4.8	4.9	5.0	5.1	5.4	5.9	6.1	6.2
2005	July	31	3.7	3.9	4.2	4.4	4.4	4.5	4.6	5.1
2005	Aug	26	4.5	4.6	4.7	4.9	4.8	4.9	5.2	5.2
2005	Sept	22	4.9	5.0	5.1	5.5	5.5	5.8	6.2	6.4
2005	Oct	22	6.0	6.5	7.0	7.3	7.3	7.6	8.2	8.5
2005	Nov	26	8.3	8.4	8.6	8.7	8.7	8.7	9.1	9.3
2005	Dec	12	9.3	9.6	10.0	10.2	10.1	10.3	10.4	10.5

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2005	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2005	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2005	Mar	2	<i>insufficient data to estimate percentiles and statistics</i>							
2005	Apr	30	7.4	7.7	8.2	9.0	8.9	9.7	10.0	10.3
2005	May	31	5.3	5.5	6.0	6.7	6.5	7.2	7.3	7.4
2005	June	21	3.9	4.2	4.5	4.9	4.9	5.3	5.7	5.8
2005	July	31	3.4	3.5	3.9	4.1	4.0	4.2	4.3	4.6
2005	Aug	26	4.1	4.1	4.2	4.4	4.4	4.5	4.8	4.8
2005	Sept	22	4.2	4.4	4.6	5.1	5.1	5.5	5.7	6.0
2005	Oct	22	5.8	6.1	6.6	7.2	7.1	7.5	7.9	8.3
2005	Nov	26	8.2	8.3	8.4	8.5	8.6	8.6	8.9	9.2
2005	Dec	12	9.2	9.4	9.9	10.0	9.9	10.1	10.3	10.4


Distribution of USGS Dissolved Oxygen Data from Chester, PA, for the Year **2006**



Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2006	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2006	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2006	Mar	1	<i>insufficient data to estimate percentiles and statistics</i>							
2006	Apr	27	6.6	6.8	7.4	7.9	8.1	9.2	9.5	9.9
2006	May	31	5.8	6.1	6.3	6.6	6.7	7.1	7.4	7.6
2006	June	30	4.6	4.7	4.8	5.1	5.3	5.5	6.3	7.0
2006	July	31	4.6	4.7	4.8	4.9	5.4	6.0	6.8	7.1
2006	Aug	31	4.6	4.8	5.0	5.2	5.1	5.4	5.5	5.7
2006	Sept	30	5.2	5.4	5.5	5.8	5.8	6.0	6.3	6.7
2006	Oct	22	6.0	6.2	6.5	7.3	7.6	8.8	9.5	9.8
2006	Nov	27	8.8	9.1	9.5	10.0	10.0	10.5	11.2	11.2
2006	Dec	0	<i>insufficient data to estimate percentiles and statistics</i>							

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2006	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2006	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2006	Mar	1	<i>insufficient data to estimate percentiles and statistics</i>							
2006	Apr	27	6.3	6.6	7.0	7.6	7.9	9.0	9.3	9.7
2006	May	31	5.6	5.7	6.0	6.3	6.4	6.8	7.2	7.3
2006	June	30	4.3	4.3	4.5	4.8	5.0	5.3	6.0	6.7
2006	July	31	4.3	4.4	4.6	4.8	5.3	5.9	6.6	7.0
2006	Aug	31	4.2	4.3	4.6	4.8	4.8	5.0	5.2	5.5
2006	Sept	30	5.0	5.1	5.3	5.6	5.6	5.9	6.2	6.4
2006	Oct	22	5.8	6.0	6.3	7.1	7.3	8.3	8.9	9.4
2006	Nov	27	8.6	8.9	9.2	9.6	9.7	10.3	10.9	11.1
2006	Dec	0	<i>insufficient data to estimate percentiles and statistics</i>							


Distribution of USGS Dissolved Oxygen Data from Chester, PA, for the Year **2007**



Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2007	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2007	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2007	Mar	1	<i>insufficient data to estimate percentiles and statistics</i>							
2007	Apr	26	11.1	11.5	11.7	12.4	12.2	12.5	13.0	13.2
2007	May	31	6.5	6.7	6.8	7.9	7.7	8.5	8.8	9.4
2007	June	26	4.7	4.9	4.9	5.2	5.3	5.7	5.9	6.5
2007	July	29	5.4	5.6	5.8	5.9	5.9	6.0	6.2	6.5
2007	Aug	22	5.1	5.2	5.3	5.4	5.4	5.5	5.6	5.7
2007	Sept	30	5.2	5.5	5.6	5.9	6.1	6.6	6.7	6.9
2007	Oct	19	6.6	6.9	6.9	7.1	7.0	7.1	7.2	7.2
2007	Nov	21	7.3	7.8	8.3	8.6	8.7	9.5	10.0	10.4
2007	Dec	18	10.6	10.8	11.2	11.3	11.3	11.4	11.9	12.3

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2007	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2007	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2007	Mar	1	<i>insufficient data to estimate percentiles and statistics</i>							
2007	Apr	26	10.8	11.3	11.5	12.1	12.0	12.4	12.8	13.0
2007	May	31	5.8	5.9	6.4	7.7	7.4	8.4	8.7	9.2
2007	June	26	4.2	4.3	4.5	4.8	4.8	5.1	5.4	5.8
2007	July	29	4.9	5.1	5.3	5.5	5.5	5.6	5.9	6.2
2007	Aug	22	4.6	4.8	4.9	4.9	5.0	5.0	5.2	5.5
2007	Sept	30	4.9	5.2	5.3	5.6	5.7	6.0	6.2	6.5
2007	Oct	19	6.2	6.4	6.6	6.7	6.7	6.8	6.9	7.2
2007	Nov	21	7.2	7.6	8.1	8.4	8.5	9.1	9.8	10.2
2007	Dec	18	10.4	10.6	11.1	11.2	11.2	11.3	11.6	12.2


Distribution of USGS Dissolved Oxygen Data from Chester, PA, for the Year **2008**



Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2008	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2008	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2008	Mar	0	<i>insufficient data to estimate percentiles and statistics</i>							
2008	Apr	25	8.4	8.9	9.6	10.0	9.8	10.3	10.4	10.6
2008	May	31	6.6	6.7	6.8	7.4	7.4	8.0	8.2	8.9
2008	June	30	4.6	4.7	4.8	5.2	6.2	7.8	8.7	9.4
2008	July	31	4.2	4.5	5.0	5.2	5.1	5.5	5.5	5.7
2008	Aug	31	4.5	5.0	5.2	5.4	5.3	5.5	5.5	5.7
2008	Sept	30	4.9	5.0	5.2	5.4	5.7	6.2	6.8	7.1
2008	Oct	31	6.2	6.4	6.5	6.8	7.2	8.1	8.3	8.5
2008	Nov	30	8.2	8.4	8.4	8.6	8.8	9.2	9.6	9.7
2008	Dec	17	9.8	9.9	9.9	10.6	10.6	11.3	11.5	11.8

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2008	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2008	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2008	Mar	0	<i>insufficient data to estimate percentiles and statistics</i>							
2008	Apr	25	8.1	8.7	9.3	9.9	9.7	10.2	10.3	10.5
2008	May	31	6.1	6.3	6.4	7.0	7.0	7.6	7.8	8.2
2008	June	30	3.9	4.2	4.3	4.8	5.7	7.3	7.7	8.7
2008	July	31	3.7	4.2	4.4	4.7	4.6	4.9	5.1	5.3
2008	Aug	31	3.9	4.2	4.8	4.9	4.8	5.0	5.1	5.3
2008	Sept	30	4.1	4.6	4.7	5.0	5.2	5.6	6.3	6.8
2008	Oct	31	5.6	5.8	6.1	6.4	6.8	7.7	8.0	8.3
2008	Nov	30	8.0	8.1	8.2	8.4	8.6	8.9	9.5	9.6
2008	Dec	17	9.6	9.7	9.8	10.3	10.4	11.0	11.3	11.6

Distribution of USGS Dissolved Oxygen Data from Chester, PA, for the Year **2009**




Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2009	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2009	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2009	Mar	0	<i>insufficient data to estimate percentiles and statistics</i>							
2009	Apr	29	8.3	8.5	8.8	9.1	9.2	9.6	9.9	10.2
2009	May	31	6.3	6.9	7.0	7.2	7.5	7.9	8.6	9.2
2009	June	30	5.8	6.0	6.1	6.4	6.7	7.2	7.7	9.0
2009	July	31	4.9	5.1	5.4	7.4	7.3	9.2	9.6	9.8
2009	Aug	28	5.0	5.2	5.4	6.5	6.3	7.0	7.4	7.7
2009	Sept	30	5.9	6.2	6.4	6.6	6.7	6.9	7.6	8.0
2009	Oct	31	6.2	6.5	6.7	7.3	7.1	7.6	7.7	7.9
2009	Nov	30	7.6	7.7	7.7	8.2	8.1	8.4	8.6	8.7
2009	Dec	10	<i>insufficient data to estimate percentiles and statistics</i>							

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2009	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2009	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2009	Mar	0	<i>insufficient data to estimate percentiles and statistics</i>							
2009	Apr	29	8.0	8.3	8.6	8.9	8.9	9.4	9.6	10.1
2009	May	31	5.9	6.5	6.6	6.8	7.1	7.5	8.3	8.6
2009	June	30	5.3	5.6	5.7	6.1	6.3	6.8	7.4	8.4
2009	July	31	4.6	4.9	5.0	6.7	6.7	8.2	8.8	9.0
2009	Aug	28	4.4	4.8	4.9	5.8	5.7	6.3	6.7	6.9
2009	Sept	30	5.6	5.8	6.1	6.2	6.3	6.5	7.0	7.3
2009	Oct	31	5.6	6.1	6.2	6.9	6.7	7.2	7.4	7.7
2009	Nov	30	7.3	7.4	7.5	7.9	7.9	8.1	8.3	8.5
2009	Dec	10	<i>insufficient data to estimate percentiles and statistics</i>							



Distribution of USGS Dissolved Oxygen Data from Chester, PA, for the Year **2010**




Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2010	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2010	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2010	Mar	0	<i>insufficient data to estimate percentiles and statistics</i>							
2010	Apr	30	8.7	8.8	9.3	10.0	9.9	10.6	10.9	11.2
2010	May	31	6.4	6.7	6.8	8.1	8.2	9.7	10.1	10.3
2010	June	24	5.3	5.4	5.5	5.9	5.9	6.0	6.6	6.7
2010	July <sup>a</sup>	18	<i>insufficient data to estimate percentiles and statistics</i>							
2010	Aug	31	5.2	5.3	5.5	5.6	5.7	6.0	6.1	6.3
2010	Sept	30	5.6	5.7	6.3	6.5	6.4	6.6	6.7	6.9
2010	Oct	24	5.7	6.7	6.9	7.1	7.0	7.3	7.3	7.7
2010	Nov	30	7.7	7.8	8.3	8.9	8.7	9.0	9.4	9.7
2010	Dec	1	<i>insufficient data to estimate percentiles and statistics</i>							

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2010	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2010	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2010	Mar	0	<i>insufficient data to estimate percentiles and statistics</i>							
2010	Apr	30	8.3	8.5	8.7	9.5	9.5	10.1	10.6	10.8
2010	May	31	6.0	6.2	6.4	7.6	7.7	9.2	9.5	9.8
2010	June	24	4.9	5.0	5.1	5.3	5.4	5.6	6.2	6.4
2010	July <sup>a</sup>	18	<i>insufficient data to estimate percentiles and statistics</i>							
2010	Aug	31	4.6	4.8	5.0	5.2	5.2	5.6	5.7	6.0
2010	Sept	30	5.3	5.5	5.7	6.1	6.0	6.2	6.2	6.6
2010	Oct	24	4.0	5.8	6.4	6.7	6.5	7.0	7.2	7.6
2010	Nov	30	7.4	7.5	8.0	8.6	8.4	8.8	9.1	9.5
2010	Dec	1	<i>insufficient data to estimate percentiles and statistics</i>							

<sup>a</sup> – missing data during early July 2010 precludes accurate estimation; known period of depressed D.O.


Distribution of USGS Dissolved Oxygen Data from Chester, PA, for the Year **2011**



Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2011	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2011	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2011	Mar	0	<i>insufficient data to estimate percentiles and statistics</i>							
2011	Apr	26	9.0	9.5	9.7	9.9	10.1	10.5	11.1	11.4
2011	May	27	7.5	7.7	7.7	8.5	8.4	8.8	9.2	9.4
2011	June	30	4.9	5.1	5.3	6.2	6.6	7.6	9.2	9.3
2011	July	31	4.8	5.0	5.2	5.7	5.7	6.3	6.4	6.8
2011	Aug	30	4.9	5.1	5.2	5.5	5.5	5.7	6.1	6.4
2011	Sept	30	6.2	6.3	7.0	7.3	7.2	7.6	7.7	8.0
2011	Oct	31	7.3	7.8	8.2	8.3	8.3	8.5	8.6	8.9
2011	Nov	30	8.9	9.2	9.6	9.7	9.7	9.9	10.0	10.2
2011	Dec	1	<i>insufficient data to estimate percentiles and statistics</i>							

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2011	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2011	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2011	Mar	0	<i>insufficient data to estimate percentiles and statistics</i>							
2011	Apr	26	8.7	9.4	9.5	9.7	9.9	10.4	10.9	11.2
2011	May	27	7.3	7.3	7.5	8.2	8.1	8.6	8.9	9.1
2011	June	30	4.4	4.7	4.9	5.6	6.0	7.0	8.2	8.8
2011	July	31	4.2	4.6	4.9	5.5	5.4	5.9	6.1	6.2
2011	Aug	30	4.5	4.7	4.8	5.2	5.1	5.3	5.8	6.0
2011	Sept	30	6.0	6.1	6.8	7.1	7.0	7.3	7.6	7.8
2011	Oct	31	7.1	7.6	7.9	8.1	8.1	8.3	8.4	8.7
2011	Nov	30	8.7	9.0	9.3	9.5	9.5	9.7	9.8	10.0
2011	Dec	1	<i>insufficient data to estimate percentiles and statistics</i>							


Distribution of USGS Dissolved Oxygen Data from Chester, PA, for the Year **2012**



Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2012	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2012	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2012	Mar	3	<i>insufficient data to estimate percentiles and statistics</i>							
2012	Apr	30	6.7	6.8	7.1	8.8	8.3	9.1	9.4	9.9
2012	May	21	5.4	5.7	5.9	6.3	6.5	6.9	7.7	7.8
2012	June	30	5.6	5.7	6.2	6.7	6.8	7.7	7.8	8.0
2012	July	31	4.2	4.5	4.7	4.9	5.0	5.3	5.5	5.8
2012	Aug	31	4.2	4.6	4.8	4.9	4.9	5.2	5.4	5.5
2012	Sept	28	4.7	5.0	5.4	5.5	5.5	5.7	6.2	6.2
2012	Oct	31	5.3	5.7	6.0	6.8	6.7	7.2	7.4	8.5
2012	Nov	30	8.4	9.1	9.4	9.6	9.6	9.8	10.2	10.2
2012	Dec	2	<i>insufficient data to estimate percentiles and statistics</i>							

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2012	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2012	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2012	Mar	3	<i>insufficient data to estimate percentiles and statistics</i>							
2012	Apr	30	6.5	6.5	6.7	8.7	8.0	8.9	9.2	9.7
2012	May	21	5.1	5.3	5.6	6.1	6.1	6.6	6.9	7.2
2012	June	30	5.4	5.5	5.8	6.3	6.3	7.0	7.2	7.5
2012	July	31	3.7	3.9	4.2	4.4	4.5	4.8	5.1	5.5
2012	Aug	31	3.4	3.9	4.3	4.4	4.3	4.6	4.6	5.1
2012	Sept	28	4.1	4.5	4.7	5.0	5.0	5.4	5.6	6.0
2012	Oct	31	4.9	5.2	5.5	6.6	6.3	6.7	7.1	8.3
2012	Nov	30	8.1	8.7	9.1	9.4	9.3	9.5	9.9	10.2
2012	Dec	2	<i>insufficient data to estimate percentiles and statistics</i>							


Distribution of USGS Dissolved Oxygen Data from Chester, PA, for the Year **2013**



Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2013	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2013	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2013	Mar	3	<i>insufficient data to estimate percentiles and statistics</i>							
2013	Apr	30	9.0	9.2	9.5	9.7	10.0	11.0	11.2	11.3
2013	May	31	6.3	6.4	7.2	7.4	7.5	7.9	8.6	9.1
2013	June	30	5.2	5.6	5.9	6.3	6.2	6.5	6.8	7.5
2013	July	31	5.3	5.5	5.8	6.2	6.4	7.0	7.8	8.1
2013	Aug	26	5.0	5.1	5.5	5.7	5.7	6.0	6.1	6.2
2013	Sept	26	5.6	5.9	6.0	6.5	6.4	6.7	6.8	7.0
2013	Oct	31	6.1	6.3	6.8	7.5	7.4	7.9	8.6	8.6
2013	Nov	30	8.1	8.3	8.5	9.5	9.3	9.6	10.3	10.5
2013	Dec	2	<i>insufficient data to estimate percentiles and statistics</i>							

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2013	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2013	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2013	Mar	3	<i>insufficient data to estimate percentiles and statistics</i>							
2013	Apr	30	8.8	8.8	8.9	9.2	9.7	10.9	11.0	11.1
2013	May	31	5.8	6.1	6.7	7.0	7.2	7.6	8.2	8.7
2013	June	30	5.0	5.2	5.6	6.0	5.9	6.1	6.4	7.1
2013	July	31	4.9	5.1	5.4	5.6	5.9	6.5	7.3	7.4
2013	Aug	26	4.7	4.8	5.1	5.4	5.3	5.7	5.7	6.0
2013	Sept	26	5.3	5.5	5.6	5.9	5.9	6.3	6.4	6.5
2013	Oct	31	5.4	5.6	6.1	7.0	6.9	7.6	8.1	8.4
2013	Nov	30	7.7	7.9	8.2	9.2	9.0	9.3	10.2	10.3
2013	Dec	2	<i>insufficient data to estimate percentiles and statistics</i>							

Distribution of USGS Dissolved Oxygen Data from Chester, PA, for the Year **2014\*\***



Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Means**							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2014	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2014	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2014	Mar	3	<i>insufficient data to estimate percentiles and statistics</i>							
2014	Apr	30	9.2	9.4	9.5	10.0	10.1	10.7	11.2	11.4
2014	May	31	6.9	7.1	7.3	7.9	8.0	8.6	8.9	9.4
2014	June	30	5.9	6.2	7.0	7.4	7.5	8.1	9.2	10.0
2014	July	31	5.0	5.3	5.5	5.8	5.8	6.2	6.5	6.8
2014	Aug	31	5.0	5.1	5.3	5.5	5.4	5.6	5.7	6.0
2014	Sept	30	4.9	4.9	5.5	5.9	6.0	6.5	6.9	7.3
2014	Oct	31	6.6	6.8	7.0	7.2	7.2	7.5	7.7	7.8
2014	Nov	29	7.9	8.3	8.5	8.7	9.2	10.2	10.3	10.4
2014	Dec	31	10.1	10.3	10.9	11.3	11.1	11.5	11.6	11.7

\*\* - erratic 15-min interval dissolved oxygen data raises questions about these statistical summaries

Year	Month	n	Dissolved Oxygen: Distribution of 24-hour Minimums**							
			min	10 <sup>th</sup>	25 <sup>th</sup>	median	mean	75 <sup>th</sup>	90 <sup>th</sup>	max
2014	Jan	0	<i>insufficient data to estimate percentiles and statistics</i>							
2014	Feb	0	<i>insufficient data to estimate percentiles and statistics</i>							
2014	Mar	3	<i>insufficient data to estimate percentiles and statistics</i>							
2014	Apr	30	7.2	8.2	8.9	9.1	9.1	9.5	10.0	10.6
2014	May	31	5.7	5.9	6.5	7.1	7.2	7.9	8.6	8.8
2014	June	30	4.5	4.8	5.9	6.4	6.5	7.1	8.0	8.9
2014	July	31	3.5	3.8	3.9	4.4	4.4	4.9	5.1	5.4
2014	Aug	31	3.0	3.2	3.7	4.1	4.1	4.5	4.8	5.0
2014	Sept	30	3.2	3.6	4.5	5.4	5.2	5.8	6.2	6.7
2014	Oct	31	5.5	6.0	6.2	6.7	6.6	6.8	7.1	7.2
2014	Nov	29	7.1	7.7	7.9	8.2	8.6	9.7	9.8	10.0
2014	Dec	31	9.0	9.8	10.3	10.8	10.6	11.1	11.3	11.3

## Appendix C. Table 1 and Selected Text from Nadeau et al. 1979

Nadeau, R., R. Hemmett, J. Miller, R.W. Miller, D.M. Clark, D. Jacangelo, L. Skinner, R.W. Marshall, T.A. Strekal, C. Emery, S.D. Selzer, D.P. Pollison, R.C. Albert, R.C. Kausch, and A.A. Anderson. 1979. Dissolved Oxygen Requirements of a “Fishable” Delaware River Estuary. Report to the Delaware River Basin Commission by the Ad Hoc Task Force to Evaluate Dissolved Oxygen Requirements of Indigenous Estuary Fish. {available online at [http://www.state.nj.us/drbc/library/documents/del-estuary\\_DOrequirements\\_1979report.pdf](http://www.state.nj.us/drbc/library/documents/del-estuary_DOrequirements_1979report.pdf)}

Nadeau et al. 1979 introduce Table 1 on pages 1 & 2 of their report:

*“Based on the knowledge of individual Task Force members and data concerning past and present Delaware Estuary Fisheries, the Task Force developed a list of thirteen migratory fish, thirteen resident fish, and one endangered fish species (Table 1). This tabulation represents a target Estuary Fish population in terms of sensitive game, forage, commercial and desired fish species rather than a total anticipated population. Total estuary fish populations are assumed to increase proportionately to the target species.*

*“For each target fish, Table 1 presents existing and potential activities (spawning, nursery, passage) by Estuary zone. The Task Force feels that current limitations on existing activities and population sizes, particularly in Zone 4, do not meet the intent of the National “fishable” goal. Restoration of fish populations to include the listed potential activities, however, will result in a minimal level of fisheries required to satisfy the national goal.”*

TABLE I

Selected Fish of Interest by Zone and Activity

Aquatic Species	DELAWARE RIVER ZONE*											
	2			3			4			5		
	S	N	P	S	N	P	S	N	P	S	N	P
<u>Migratory or Marine/Estuarine:</u>												
Striped bass	o	o	x	o	o	x	o	o	x	x	x	x
Alewife	x	x	x	x	x		o	x		x	x	
Blueback herring	x	x	x	x	x		o	x		x	x	
American shad	o	x	x	o	o	x	o	o	x			x
Mummichog			x**			x	x	x	x	x	x	x
Bay anchovy											x	x
Menhaden							x	x			x	x
Bluefish											x	x
Weakfish											x	x
Spot											x	x
Atlantic croaker											x	x
Atlantic silversides											x	x
Blue crab			x**			x**			x**		x	x
<u>Resident (freshwater/brackish water):</u>												
Pumpkinseed/bluegill	x	x	x	x	x	x	o	o	x	x	x	x
Largemouth bass	x	x	x	o	o	x	o	o	o	o	o	o
Walleye	x <sup>+</sup>	x	x									
Channel catfish	x	x	x	x	x	x	o	o	x	x	x	x
Black crappie	x	x	x	o	o	o	o	o	o			x
Golden shiner	x	x	x	x	x	x	o	o	o			x
White perch	x	x	x	x	x	x	o	o	x	x	x	x
Silvery minnow	x	x	x	x	x	x	o	o	x	x	x	x
Carp	x	x	x	x	x	x	x	x	x	x	x	x
Goldfish	x	x	x	x	x	x	x	x	x	x	x	x
Brown bullhead	x	x	x	x	x	x	x	x	x	x	x	x
Yellow perch												x
<u>Endangered Species</u>												
Short nose sturgeon	x	x	x	o	x		o	x		x	x	

\*Key

S = Spawning

N = Nursery

P = Passage

o = potential

x = existing

\*\*Limited or occasional presence (limited for reasons other than water quality)

+ Limited spawning in area (limited for reasons other than water quality)



## Appendix D. PSEG Ichthyoplankton Data from 2002, 2003, and 2004 Biological Monitoring Program Annual Reports

### **Report Citations:**

- PSEG (Public Service Enterprise Group). 2002. Biological Monitoring Program Annual Report-2002. 754 pp.
- PSEG (Public Service Enterprise Group). 2003. Biological Monitoring Program Annual Report-2003. 740 pp.
- PSEG (Public Service Enterprise Group). 2004. Biological Monitoring Program Annual Report-2004. 762 pp.

# 2002 DATA

Table 4-33 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 1) for each life stage of each target  
 species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	1,118.14	0.00	1,035.82	5,681.23	153.39	157.84	0.00	0.00	0.00
yolk sac larvae	84.45	0.00	13.04	506.70	9.54	1.52	0.00	0.00	0.00
post-yolk sac larvae	80.17	7.44	11.08	142.98	326.23	1.38	2.73	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	12.12	0.00	15.82	60.33	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.05	0.00	0.00	0.33	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	1.28	0.00	0.00	5.19	1.47	1.38	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.14	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.41
undetermined larvae	0.26	0.00	0.00	0.00	1.65	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	33,247.03	0.00	306.40	48,213.95	43,354.13	107,106.65	19,064.96	1,073.10	489.09
yolk sac larvae	0.05	0.00	0.00	0.29	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	151.96	0.28	11.67	92.68	13.69	297.82	413.67	38.63	508.23
juvenile	2.46	0.00	0.00	0.00	0.00	0.00	0.00	0.00	24.04
adult	0.25	0.00	0.71	0.29	0.00	0.60	0.01	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.05	0.00	0.00	0.00	0.30	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-33 (page 2 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 1) for each life stage of each target  
 species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	7.23	0.00	0.00	0.00	45.43	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	2.97	0.00	0.00	0.00	17.76	0.89	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.04	0.00	0.25	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	5,765.58	0.00	1.00	15,502.52	7,990.41	7,768.09	1,718.81	28,835.66	0.00
yolk sac larvae	6.03	0.00	0.00	21.12	0.00	16.77	0.00	0.00	0.00
post-yolk sac larvae	20.41	0.00	10.24	33.82	26.45	35.77	44.02	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>40,501</b>	<b>8</b>	<b>1,406</b>	<b>70,261</b>	<b>51,940</b>	<b>115,389</b>	<b>21,244</b>	<b>29,947</b>	<b>1,023</b>
<b>Gammarus spp.</b>	281	133	387	166	542	455	13	39	116
<b>Neomysis americana</b>	210,668	9,565	212,993	171,560	687,653	53,729	228,877	367,598	33,827

Note: Event 1 occurred April 9-18.  
 Event 2 occurred April 23-May 2.  
 Event 3 occurred May 6-21.  
 Event 4 occurred May 22-30.  
 Event 5 occurred June 7-18.  
 Event 6 occurred June 24-29.  
 Event 7 occurred July 8-17.  
 Event 8 occurred July 23-30.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-34 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 2) for each life stage of each target  
 species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	1,352.49	0.00	248.59	8,317.16	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	91.22	0.00	0.36	575.51	1.87	0.00	0.00	0.00	0.00
post-yolk sac larvae	31.49	11.77	2.86	142.08	42.34	0.50	0.00	0.00	0.00
juvenile	0.13	0.00	0.00	0.00	0.00	1.06	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	15.20	0.00	13.10	83.15	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	1.30	0.00	0.00	7.09	0.53	0.00	0.00	0.00	0.00
post-yolk sac larvae	22.37	0.00	3.30	45.97	80.61	6.80	3.85	0.00	0.00
juvenile	0.17	0.00	0.00	0.00	0.00	1.36	0.00	0.00	0.00
adult	0.16	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.53
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	14,432.95	0.00	77.04	49,019.94	32,529.57	6,948.29	7,518.83	276.54	162.93
yolk sac larvae	0.07	0.00	0.00	0.00	0.47	0.00	0.00	0.00	0.00
post-yolk sac larvae	405.18	0.00	7.99	184.16	7.73	1,509.84	1,250.51	374.90	558.38
juvenile	30.05	0.00	0.00	0.00	0.00	0.00	0.00	5.85	282.53
adult	5.49	4.76	11.20	12.59	3.35	1.06	1.93	0.00	1.53
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-34 (page 2 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 2) for each life stage of each target species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.47	0.45	2.53	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	8,462.40	0.00	0.00	49,850.05	864.43	1,503.97	2,762.46	526.51	1.12
yolk sac larvae	3.19	0.00	0.00	17.78	0.00	3.11	0.00	0.00	0.00
post-yolk sac larvae	46.66	0.00	1.75	76.68	4.97	142.91	174.48	2.92	4.57
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.12	0.00	0.00	0.00	0.00	1.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>24,901</b>	<b>17</b>	<b>369</b>	<b>108,332</b>	<b>33,536</b>	<b>10,120</b>	<b>11,712</b>	<b>1,187</b>	<b>1,013</b>
<b>Gammarus spp.</b>	1,465	412	2,652	704	2,375	2,767	334	1,550	271
<b>Neomysis americana</b>	174,555	36,898	191,769	158,156	273,737	111,460	612,125	28,015	13,280

Note: Event 1 occurred April 9-18.  
 Event 2 occurred April 23-May 2.  
 Event 3 occurred May 6-21.  
 Event 4 occurred May 22-30.  
 Event 5 occurred June 7-18.  
 Event 6 occurred June 24-29.  
 Event 7 occurred July 8-17.  
 Event 8 occurred July 23-30.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-35 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 3) for each life stage of each target  
 species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	2.26	0.00	10.42	2.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	356.61	0.00	1,052.20	0.00	1,309.44	123.50	0.00	0.00	0.00
yolk sac larvae	8.22	0.00	20.93	0.00	0.72	23.87	0.00	0.00	0.00
post-yolk sac larvae	42.55	2.48	78.14	114.63	15.82	22.10	0.00	0.00	0.00
juvenile	28.91	0.00	1.07	157.94	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.26	0.00	0.00	1.43	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	34.46	0.00	9.03	173.56	2.79	4.40	4.28	0.00	0.00
juvenile	0.68	0.00	0.46	0.00	0.00	0.00	16.54	0.00	0.00
adult	1.63	0.00	1.27	0.89	0.00	0.00	0.00	9.73	0.00
undetermined larvae	2.62	0.00	0.00	14.42	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	20,154.21	0.00	156.07	1,180.70	19,645.13	97,644.28	0.00	114.34	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	716.43	0.39	10.72	4.19	42.91	2,888.13	3,183.33	636.79	0.00
juvenile	157.05	0.00	0.00	0.00	0.00	2.39	1,647.28	759.88	0.00
adult	24.36	2.49	69.63	55.40	0.00	0.99	12.83	4.13	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-35 (page 2 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 3) for each life stage of each target  
 species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.84	0.00	0.00	0.00	2.68	2.99	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	2,521.60	0.00	0.00	459.67	450.35	13,131.09	0.00	11.20	0.00
yolk sac larvae	0.25	0.00	0.00	0.00	0.68	0.96	0.00	0.00	0.00
post-yolk sac larvae	38.59	0.00	0.00	0.45	44.65	173.34	16.54	11.90	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>24,092</b>	<b>5</b>	<b>1,410</b>	<b>2,165</b>	<b>21,515</b>	<b>114,018</b>	<b>4,881</b>	<b>1,548</b>	<b>0</b>
<b>Gammarus spp.</b>	<b>3,718</b>	<b>205</b>	<b>477</b>	<b>13,742</b>	<b>1,000</b>	<b>524</b>	<b>6,139</b>	<b>5,251</b>	<b>0</b>
<b>Neomysis americana</b>	<b>485,405</b>	<b>81,323</b>	<b>202,453</b>	<b>1,960,906</b>	<b>623,378</b>	<b>33,164</b>	<b>6,175</b>	<b>23,743</b>	<b>0</b>

Note: Event 1 occurred April 9-18.  
 Event 2 occurred April 23-May 2.  
 Event 3 occurred May 6-21.  
 Event 4 occurred May 22-30.  
 Event 5 occurred June 7-18.  
 Event 6 occurred June 24-29.  
 Event 7 occurred July 8-17.  
 Event 8 occurred July 23-30.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.



Table 4-36 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 4) for each life stage of each target  
 species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.99	0.00	4.71	1.68	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	1.34	0.00	0.00	7.23	0.87	0.00	0.00	0.00	0.00
post-yolk sac larvae	34.31	30.37	7.15	151.88	0.00	6.76	0.00	0.00	0.00
juvenile	7.14	0.00	0.00	40.32	0.52	2.57	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	13.39	0.00	1.33	79.17	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	2.81	0.00	0.00	16.44	0.45	0.00	0.00	0.00	0.00
post-yolk sac larvae	39.72	0.00	6.29	231.83	0.95	0.00	0.00	0.00	0.00
juvenile	4.03	0.00	0.00	0.00	0.00	0.00	0.00	114.72	1.25
adult	0.28	0.00	0.00	0.00	0.00	0.00	0.00	4.08	0.84
undetermined larvae	11.36	0.00	0.00	68.16	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	32,976.39	0.37	2.80	35,789.19	159,837.60	21,238.81	712.40	131.31	1,052.64
yolk sac larvae	1.63	0.00	0.00	0.00	10.89	0.00	0.00	0.00	0.00
post-yolk sac larvae	659.28	0.47	2.59	11.00	4.65	1,320.84	5,213.47	2,443.21	1,349.71
juvenile	76.92	0.00	0.00	1.91	0.00	0.00	40.31	299.35	391.68
adult	11.78	27.13	6.23	32.49	0.55	0.00	20.15	0.00	0.93
undetermined larvae	5.41	0.00	1.42	0.00	0.00	0.00	0.00	0.00	31.18
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.07	0.00	0.00	0.41	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-36 (page 2 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 4) for each life stage of each target  
 species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	410.85	0.00	0.56	1,699.21	0.00	866.93	0.00	0.00	71.85
yolk sac larvae	0.72	0.00	0.00	0.84	3.58	0.00	0.00	0.00	0.00
post-yolk sac larvae	5.73	0.00	0.00	7.54	0.43	31.77	5.04	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>34,264</b>	<b>58</b>	<b>33</b>	<b>38,139</b>	<b>159,860</b>	<b>23,468</b>	<b>5,991</b>	<b>2,993</b>	<b>2,900</b>
<b>Gammarus spp.</b>	1,985	1,148	203	1,801	112	1,356	220	18,011	4,137
<b>Neomysis americana</b>	145,806	95,735	562,307	104,271	55,579	76,286	221,479	15,301	19,367

Note: Event 1 occurred April 9-18.  
 Event 2 occurred April 23-May 2.  
 Event 3 occurred May 6-21.  
 Event 4 occurred May 22-30.  
 Event 5 occurred June 7-18.  
 Event 6 occurred June 24-29.  
 Event 7 occurred July 8-17.  
 Event 8 occurred July 23-30.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-37 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 5) for each life stage of each target  
 species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	1.91	0.00	0.00	0.00	15.26	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	13.35	0.00	0.00	0.00	106.79	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	6.36	0.00	0.00	0.00	50.85	0.00	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.43	1.68	0.00	0.88	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.14	0.00	1.13	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	12.61	85.48	11.99	1.69	0.00	0.00	0.00	0.00	0.00
juvenile	3.75	0.00	0.60	13.98	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	3.12	0.00	2.61	11.17	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	18.96	0.00	34.99	52.30	0.69	7.55	3.83	0.00	0.00
juvenile	0.44	0.00	0.00	0.00	0.00	0.00	3.53	0.00	0.00
adult	0.08	0.61	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	924.51	0.91	0.00	743.33	2,995.26	706.36	1,532.57	674.34	0.00
yolk sac larvae	0.08	0.00	0.00	0.33	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	1,010.62	0.00	0.00	3.49	3,847.17	2,238.94	1,900.20	91.62	0.00
juvenile	22.14	0.00	0.00	0.00	3.27	24.38	55.53	93.98	0.00
adult	9.21	31.81	15.30	11.89	0.00	1.40	0.68	0.70	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-37 (page 2 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 5) for each life stage of each target  
 species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.90	0.00	0.00	3.60	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	6.99	0.00	0.00	0.00	55.94	0.00	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	0.61	0.00	0.00	2.46	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	5.01	0.00	0.00	19.35	0.00	1.40	0.00	0.00	0.00
post-yolk sac larvae	0.78	0.00	0.00	3.14	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	19.33	0.00	0.00	19.64	0.00	115.36	0.00	0.00	0.00
yolk sac larvae	0.07	0.00	0.00	0.28	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	7.41	0.00	0.00	0.00	7.34	46.54	4.72	0.66	0.00
juvenile	1.25	0.00	0.00	0.00	0.00	9.98	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.09	0.00	0.00	0.36	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	225.66	0.00	0.00	0.00	1,805.29	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>2,296</b>	<b>120</b>	<b>67</b>	<b>888</b>	<b>8,888</b>	<b>3,152</b>	<b>3,501</b>	<b>861</b>	<b>0</b>
<b>Gammarus spp.</b>	<b>4,178</b>	<b>895</b>	<b>1,275</b>	<b>5,521</b>	<b>1,610</b>	<b>14,004</b>	<b>2,759</b>	<b>1,837</b>	<b>0</b>
<b>Neomysis americana</b>	<b>136,561</b>	<b>258,373</b>	<b>25,478</b>	<b>106,353</b>	<b>400,286</b>	<b>134,830</b>	<b>21,502</b>	<b>39,312</b>	<b>0</b>

Note: Event 1 occurred April 9-18.  
 Event 2 occurred April 23-May 2.  
 Event 3 occurred May 6-21.  
 Event 4 occurred May 22-30.  
 Event 5 occurred June 7-18.  
 Event 6 occurred June 24-29.  
 Event 7 occurred July 8-17.  
 Event 8 occurred July 23-30.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-38 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 6) for each life stage of each target  
 species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.20	0.00	0.00	1.59	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	9.90	8.69	11.54	43.33	15.67	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	12.84	5.14	16.06	0.00	0.00	79.13	2.38	0.00	0.00
juvenile	4.59	0.00	2.86	9.58	24.26	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.57	0.00	0.00	0.00	0.00	3.52	1.03	0.00	0.00
yolk sac larvae	3.28	0.00	0.00	0.00	7.51	18.75	0.00	0.00	0.00
post-yolk sac larvae	6.91	0.00	0.00	0.00	2.39	29.76	15.95	5.97	1.21
juvenile	0.28	0.00	0.00	0.00	0.00	0.00	0.00	1.01	1.20
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.30	0.00	0.00	0.00	0.00	2.42	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	2,493.46	0.00	0.00	0.00	554.53	596.41	17,951.44	441.24	404.08
yolk sac larvae	45.33	0.00	0.00	0.00	0.00	0.00	362.61	0.00	0.00
post-yolk sac larvae	1,621.48	0.00	0.00	0.00	19.65	3,596.82	7,556.32	1,585.80	213.29
juvenile	34.12	0.00	5.51	0.00	0.00	2.40	135.56	52.79	76.73
adult	15.44	4.01	17.78	6.73	91.66	2.37	0.00	0.00	0.95
undetermined larvae	0.15	0.00	0.00	0.00	0.00	0.00	0.00	1.22	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-38 (page 2 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 6) for each life stage of each target  
 species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.98	0.00	0.00	7.80	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.30	0.00	0.00	2.36	0.00	0.00	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	50.05	0.00	0.00	400.38	0.00	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	12.05	0.00	0.00	96.40	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	99.61	1.15	0.00	766.75	28.94	0.00	0.00	0.00	0.00
post-yolk sac larvae	52.64	0.00	0.00	302.08	109.38	9.67	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	16.89	0.00	0.00	0.00	2.50	0.00	132.63	0.00	0.00
yolk sac larvae	3.70	0.00	0.00	0.00	10.88	0.00	18.70	0.00	0.00
post-yolk sac larvae	28.67	0.00	0.00	0.00	0.00	151.76	77.59	0.00	0.00
juvenile	3.26	0.00	1.03	0.00	0.00	17.94	7.13	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.59	0.00	0.00	2.36	2.39	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.12	0.00	0.00	0.96	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>4,518</b>	<b>19</b>	<b>55</b>	<b>1,640</b>	<b>870</b>	<b>4,511</b>	<b>26,261</b>	<b>2,088</b>	<b>697</b>
<b>Gammarus spp.</b>	935	44	245	2,790	830	965	2,342	202	64
<b>Neomysis americana</b>	23,067	3,015	32,163	9,991	39,508	20,938	26,765	14,719	37,438

Note: Event 1 occurred April 9-18.  
 Event 2 occurred April 23-May 2.  
 Event 3 occurred May 6-21.  
 Event 4 occurred May 22-30.  
 Event 5 occurred June 7-18.  
 Event 6 occurred June 24-29.  
 Event 7 occurred July 8-17.  
 Event 8 occurred July 23-30.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-39 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 7) for each life stage of each target species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	1.91	0.00	0.00	0.00	15.26	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	18.14	114.68	6.90	12.94	7.17	3.40	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	14.65	57.41	11.47	0.00	2.63	42.02	3.66	0.00	0.00
juvenile	4.15	3.55	0.00	11.46	10.11	2.45	3.76	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.13	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.07
yolk sac larvae	6.80	0.00	0.00	13.02	0.00	3.54	29.38	8.45	0.00
post-yolk sac larvae	13.32	0.00	1.49	4.70	2.34	32.30	21.74	43.35	3.56
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	6.65	0.00	0.00	0.00	24.47	0.00	0.00	27.51	1.19
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	166.49	0.00	3.90	0.00	0.00	227.49	510.47	370.01	220.07
juvenile	15.20	3.86	3.90	0.00	0.00	0.00	49.87	16.67	47.27
adult	18.63	41.52	34.00	25.85	42.84	0.00	3.66	0.00	1.19
undetermined larvae	26.19	0.00	0.00	0.00	0.00	0.00	0.00	209.48	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-39 (page 2 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 7) for each life stage of each target species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.14	0.00	0.00	1.14	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	3.35	0.00	0.00	2.30	6.52	17.16	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	5.51	0.00	0.00	7.11	25.98	11.03	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	86.37	0.00	0.00	661.76	2.26	26.97	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.14	0.00	0.00	1.14	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	22.81	5.92	5.01	134.24	36.22	1.08	0.00	0.00	0.00
yolk sac larvae	82.03	2.37	7.99	323.21	320.40	2.31	0.00	0.00	0.00
post-yolk sac larvae	317.59	0.00	0.00	236.32	1,809.49	490.31	4.62	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	37.75	0.00	0.00	0.00	302.03	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.14	0.00	0.00	0.00	1.13	0.00	0.00	0.00	0.00
yolk sac larvae	0.28	0.00	0.00	0.00	2.26	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.89	0.00	0.00	0.00	0.00	1.09	6.03	0.00	0.00
juvenile	2.93	2.89	0.00	0.00	0.00	0.00	20.56	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	1.64	1.18	0.00	5.74	6.17	0.00	0.00	0.00	0.00
post-yolk sac larvae	14.78	0.00	0.00	1.14	7.67	105.65	3.76	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>869</b>	<b>233</b>	<b>75</b>	<b>1,442</b>	<b>2,625</b>	<b>967</b>	<b>658</b>	<b>675</b>	<b>274</b>
<b>Gammarus spp.</b>	<b>5,257</b>	<b>2,736</b>	<b>221</b>	<b>7,106</b>	<b>15,039</b>	<b>5,612</b>	<b>9,608</b>	<b>1,162</b>	<b>572</b>
<b>Neomysis americana</b>	<b>18,395</b>	<b>8,808</b>	<b>26,684</b>	<b>2,249</b>	<b>67,371</b>	<b>1,671</b>	<b>13,674</b>	<b>17,079</b>	<b>9,625</b>

Note: Event 1 occurred April 9-18.  
 Event 2 occurred April 23-May 2.  
 Event 3 occurred May 6-21.  
 Event 4 occurred May 22-30.  
 Event 5 occurred June 7-18.  
 Event 6 occurred June 24-29.  
 Event 7 occurred July 8-17.  
 Event 8 occurred July 23-30.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.



Table 4-40 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 8) for each life stage of each target  
 species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	107.71	0.00	4.08	8.37	839.58	9.65	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	29.47	40.51	172.09	4.18	7.01	9.91	2.06	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	12.70	29.11	58.02	0.00	0.00	2.21	12.24	0.00	0.00
juvenile	1.94	0.00	0.00	6.28	4.67	2.48	2.06	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	4.18	2.28	0.00	2.58	0.00	0.00	2.06	10.45	16.88
post-yolk sac larvae	6.70	0.00	8.47	2.09	0.00	0.00	16.31	17.12	9.64
juvenile	0.82	0.00	6.58	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	43.51	0.00	17.55	0.00	0.00	0.00	65.12	119.28	146.16
juvenile	15.17	0.00	38.58	0.00	0.00	0.00	20.48	2.07	60.23
adult	3.01	4.56	7.56	11.93	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.29	2.28	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-40 (page 2 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 8) for each life stage of each target species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	18.61	0.00	0.00	9.99	131.97	6.91	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	112.32	0.00	0.00	0.00	652.81	245.74	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	207.54	0.00	0.00	0.00	615.88	1,044.43	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	2,111.79	34.46	227.75	16,437.24	194.90	0.00	0.00	0.00	0.00
yolk sac larvae	1,682.94	2.28	983.27	11,685.59	792.38	0.00	0.00	0.00	0.00
post-yolk sac larvae	208.66	0.00	7.86	495.62	935.02	193.72	37.05	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.30	0.00	0.00	0.00	0.00	0.00	0.00	0.00	2.41
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	34.23	32.18	61.42	129.06	48.98	2.21	0.00	0.00	0.00
post-yolk sac larvae	218.65	0.00	10.36	15.80	861.61	842.98	18.42	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.58	0.00	0.00	0.00	4.67	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>4,821</b>	<b>148</b>	<b>1,604</b>	<b>28,809</b>	<b>5,089</b>	<b>2,360</b>	<b>176</b>	<b>149</b>	<b>235</b>
<b>Gammarus spp.</b>	40,420	2,040	23,397	20,899	75,881	135,009	35,174	12,113	18,849
<b>Neomysis americana</b>	6,391	53	63	230	65	263	22	15,918	34,509

Note: Event 1 occurred April 9-18.  
 Event 2 occurred April 23-May 2.  
 Event 3 occurred May 6-21.  
 Event 4 occurred May 22-30.  
 Event 5 occurred June 7-18.  
 Event 6 occurred June 24-29.  
 Event 7 occurred July 8-17.  
 Event 8 occurred July 23-30.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-41 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 9) for each life stage of each target  
 species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.11	0.88	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	89.26	2.52	21.26	75.56	515.45	79.28	19.17	0.87	0.00
juvenile	0.22	0.00	0.00	0.00	0.00	0.00	0.88	0.87	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.72	0.00	5.75	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.45	0.00	0.00	0.00	0.00	0.00	0.00	3.56	0.00
juvenile	0.43	0.00	0.00	0.00	0.00	0.00	0.00	3.45	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.22	0.00	0.00	0.00	1.76	0.00	0.00	0.00	0.00
yolk sac larvae	15.61	0.00	0.00	0.00	0.00	124.92	0.00	0.00	0.00
post-yolk sac larvae	5.50	0.00	0.00	0.00	0.00	44.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.10	0.00	0.81	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	3.48	0.83	10.76	10.89	3.78	0.84	0.75	0.00	0.00
adult	0.11	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.85
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	15.39	9.13	63.32	1.83	0.00	35.78	12.09	0.95	0.00
juvenile	2.56	0.00	0.00	4.35	8.56	0.00	1.48	6.09	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.18	0.00	0.00	0.00	0.00	0.00	1.48	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	11.81	0.00	0.00	0.00	0.00	0.00	5.94	15.77	72.79
juvenile	6.46	0.00	0.00	0.00	0.00	0.00	5.16	0.00	46.56
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.11	0.00	0.00	0.00	0.00	0.00	0.00	0.88	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-41 (page 2 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 9) for each life stage of each target species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	64.93	0.85	4.08	324.58	134.45	59.26	7.65	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	291.44	0.00	0.81	362.67	1,380.64	587.37	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	164.20	0.00	0.00	439.18	287.32	579.28	1.75	6.03	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	2,724.59	9.91	290.49	21,200.22	294.33	1.75	0.00	0.00	0.00
yolk sac larvae	1,437.35	0.00	6,272.87	4,465.36	759.66	0.89	0.00	0.00	0.00
post-yolk sac larvae	233.00	0.00	18.95	699.48	968.38	146.53	61.56	3.41	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	46.32	0.00	0.00	0.00	370.59	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	5.54	29.92	0.81	7.97	5.59	0.00	0.00	0.00	0.00
yolk sac larvae	41.32	59.23	63.31	122.40	66.69	18.88	0.00	0.00	0.00
post-yolk sac larvae	412.92	0.00	155.68	255.37	491.29	2,092.69	247.35	60.02	0.92
juvenile	0.21	0.00	1.67	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>5,575</b>	<b>113</b>	<b>6,911</b>	<b>27,970</b>	<b>5,288</b>	<b>3,771</b>	<b>365</b>	<b>102</b>	<b>121</b>
<b>Gammarus spp.</b>	<b>31,177</b>	<b>344</b>	<b>9,596</b>	<b>42,369</b>	<b>73,441</b>	<b>72,400</b>	<b>26,912</b>	<b>11,301</b>	<b>13,050</b>
<b>Neomysis americana</b>	<b>108</b>	<b>0</b>	<b>11</b>	<b>0</b>	<b>2</b>	<b>3</b>	<b>1</b>	<b>0</b>	<b>846</b>

Note: Event 1 occurred April 9-18.  
 Event 2 occurred April 23-May 2.  
 Event 3 occurred May 6-21.  
 Event 4 occurred May 22-30.  
 Event 5 occurred June 7-18.  
 Event 6 occurred June 24-29.  
 Event 7 occurred July 8-17.  
 Event 8 occurred July 23-30.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-42 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 10) for each life stage of each target  
 species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	148.02	0.93	121.50	523.93	313.12	172.65	22.41	0.00	0.00
juvenile	0.12	0.00	0.00	0.00	0.00	0.00	0.90	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	3.71	4.53	16.13	5.39	2.90	0.00	0.00	0.00	0.00
yolk sac larvae	14.53	0.00	33.74	55.04	24.53	0.00	0.00	0.00	0.00
post-yolk sac larvae	1.02	0.00	0.00	0.00	0.00	0.00	7.96	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.13	0.00	0.00	0.99	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	12.15	0.00	0.00	0.00	26.04	66.77	0.00	1.92	0.00
post-yolk sac larvae	13.68	0.00	0.00	8.52	38.76	41.86	2.65	9.25	0.00
juvenile	0.12	0.00	0.00	0.00	0.00	0.00	0.00	0.96	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.12	0.00	0.00	0.90	0.00	0.00	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	1.42	0.00	2.79	3.24	1.96	0.85	0.00	1.91	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	3.04	0.72	16.51	0.96	3.75	0.00	1.80	0.00	0.00
juvenile	0.56	0.00	0.00	0.00	0.00	0.00	0.00	4.39	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	0.66	0.00	0.00	0.00	0.00	5.17	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	1.05	0.00	0.00	0.00	0.00	0.00	0.88	1.79	6.88
juvenile	0.34	0.00	0.00	0.00	0.00	0.00	0.00	0.88	2.24
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.58	0.00	0.00	0.00	0.00	1.65	0.00	2.86	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-42 (page 2 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 10) for each life stage of each target  
 species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	29.87	0.72	17.05	22.00	7.26	172.83	8.94	8.40	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	248.38	0.00	0.00	476.68	365.28	1,045.83	0.00	2.82	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	1.64	0.00	0.00	12.79	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	428.68	0.00	0.00	140.35	1,523.19	1,680.19	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	11.81	1.72	0.00	83.19	7.22	0.00	0.00	0.00	0.00
yolk sac larvae	5.21	0.00	31.57	5.39	1.94	0.00	1.76	0.00	0.00
post-yolk sac larvae	124.96	0.00	15.43	140.56	489.99	323.99	4.68	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	46.63	0.00	0.00	0.00	363.72	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.11	0.00	0.00	0.00	0.00	0.87	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	31.91	72.45	4.69	128.26	37.48	6.05	0.00	0.00	0.00
yolk sac larvae	67.08	32.14	280.52	76.60	97.37	33.75	2.84	0.00	0.00
post-yolk sac larvae	836.10	0.00	543.80	211.97	3,562.74	1,586.83	477.57	137.76	1.12
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.09	0.72	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>2,034</b>	<b>114</b>	<b>1,084</b>	<b>1,897</b>	<b>6,867</b>	<b>5,139</b>	<b>532</b>	<b>173</b>	<b>10</b>
<b>Gammarus spp.</b>	<b>37,561</b>	<b>279</b>	<b>28,261</b>	<b>20,251</b>	<b>105,818</b>	<b>42,128</b>	<b>33,840</b>	<b>48,501</b>	<b>17,369</b>
<b>Neomysis americana</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>3</b>	<b>0</b>	<b>3</b>	<b>0</b>

Note: Event 1 occurred April 9-18.  
 Event 2 occurred April 23-May 2.  
 Event 3 occurred May 6-21.  
 Event 4 occurred May 22-30.  
 Event 5 occurred June 7-18.  
 Event 6 occurred June 24-29.  
 Event 7 occurred July 8-17.  
 Event 8 occurred July 23-30.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-43 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 11) for each life stage of each target  
 species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	1.10	0.00	0.00	0.00	8.81	0.00	0.00	0.00	0.00
post-yolk sac larvae	164.16	1.85	1,019.88	63.55	14.21	189.62	17.50	6.69	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	2.64	0.00	0.00	0.00	21.14	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	53.05	5.90	270.24	144.06	2.60	1.61	0.00	0.00	0.00
yolk sac larvae	103.16	0.00	710.45	53.48	51.50	9.85	0.00	0.00	0.00
post-yolk sac larvae	13.32	0.00	0.00	0.00	83.07	0.00	23.52	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	3.84	0.00	0.00	0.00	30.72	0.00	0.00	0.00	0.00
yolk sac larvae	7.43	0.00	1.84	3.39	0.88	53.34	0.00	0.00	0.00
post-yolk sac larvae	7.72	0.00	1.86	25.03	5.16	28.77	0.00	0.96	0.00
juvenile	0.11	0.00	0.00	0.00	0.88	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.21	0.00	0.00	1.65	0.00	0.00	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.11	0.00	0.85	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.98	3.53	4.32	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.36	0.00	0.00	0.00	0.00	0.00	0.00	2.87	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.11	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.88
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-43 (page 2 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 11) for each life stage of each target  
 species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	2.53	0.00	0.00	0.00	20.26	0.00	0.00	0.00	0.00
post-yolk sac larvae	13.66	0.00	8.66	1.65	3.55	63.09	13.37	18.12	0.86
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	97.61	0.00	2.45	247.29	205.51	290.17	0.00	3.83	1.71
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	48.48	0.00	0.00	0.00	0.00	387.83	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	86.65	0.00	0.00	263.51	0.00	429.69	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	1.68	2.59	0.00	1.71	5.23	3.94	0.00	0.00	0.00
yolk sac larvae	0.53	0.00	0.00	0.00	1.74	1.64	0.85	0.00	0.00
post-yolk sac larvae	5.80	0.00	3.65	21.11	11.01	4.93	0.00	5.68	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	59.86	73.73	8.35	322.06	48.55	26.21	0.00	0.00	0.00
yolk sac larvae	121.15	27.43	602.28	16.71	252.13	63.84	6.79	0.00	0.00
post-yolk sac larvae	1,056.64	2.90	1,208.16	309.89	627.21	6,061.54	107.66	134.88	0.86
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	30.93	0.00	0.00	2.46	244.97	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>1,884</b>	<b>118</b>	<b>3,843</b>	<b>1,478</b>	<b>1,639</b>	<b>7,616</b>	<b>170</b>	<b>173</b>	<b>4</b>
<b>Gammarus spp.</b>	<b>17,688</b>	<b>226</b>	<b>6,505</b>	<b>27,186</b>	<b>17,182</b>	<b>21,073</b>	<b>9,048</b>	<b>35,832</b>	<b>24,455</b>
<b>Neomysis americana</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>1</b>	<b>4</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>

Note: Event 1 occurred April 9-18.  
 Event 2 occurred April 23-May 2.  
 Event 3 occurred May 6-21.  
 Event 4 occurred May 22-30.  
 Event 5 occurred June 7-18.  
 Event 6 occurred June 24-29.  
 Event 7 occurred July 8-17.  
 Event 8 occurred July 23-30.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.



Table 4-44 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 12) for each life stage of each target  
 species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	1.46	0.00	0.00	12.01	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	428.22	0.68	3,005.00	14.46	273.29	201.18	15.82	0.97	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	181.19	159.69	501.05	813.67	11.37	0.00	0.00	0.00	0.00
yolk sac larvae	256.94	3.22	1,788.97	28.11	278.24	8.39	0.00	0.00	0.00
post-yolk sac larvae	19.19	0.00	0.00	0.00	149.82	0.00	3.55	3.96	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	5.13	17.47	13.81	0.00	10.78	0.00	0.00	0.00	0.00
yolk sac larvae	7.96	0.00	20.88	17.28	27.11	0.00	0.00	0.00	0.00
post-yolk sac larvae	46.75	0.00	99.70	206.15	34.61	39.10	0.00	5.28	0.00
juvenile	0.11	0.00	0.00	0.00	0.00	0.00	0.89	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	2.59	0.00	0.00	21.26	0.00	0.00	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	4.30	0.00	0.00	0.00	0.00	35.28	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.83	0.00	0.00	0.00	0.00	0.00	0.89	0.74	4.35
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.11	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.73
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-44 (page 2 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 12) for each life stage of each target  
 species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	21.50	0.00	0.00	0.00	76.76	44.04	30.21	6.26	17.27
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	62.53	0.00	0.00	12.06	271.17	226.77	0.00	2.74	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	123.89	0.00	0.00	99.10	630.65	286.16	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	1.55	0.00	9.39	0.86	2.46	0.00	0.00	0.00	0.00
yolk sac larvae	1.07	0.00	6.13	0.00	0.00	0.00	2.66	0.00	0.00
post-yolk sac larvae	3.14	0.00	0.00	2.56	1.63	18.92	2.66	0.00	0.00
juvenile	0.11	0.00	0.00	0.00	0.00	0.00	0.89	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	208.49	465.31	64.08	1,087.85	92.38	0.00	0.00	0.00	0.00
yolk sac larvae	210.20	215.45	827.35	31.95	403.03	244.07	1.78	0.00	0.00
post-yolk sac larvae	1,009.17	0.00	2,619.90	174.91	2,539.05	2,017.48	734.86	186.31	2.21
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>2,596</b>	<b>862</b>	<b>8,956</b>	<b>2,522</b>	<b>4,802</b>	<b>3,121</b>	<b>794</b>	<b>206</b>	<b>25</b>
<b>Gammarus spp.</b>	12,881	63	3,610	19,525	10,469	1,609	8,185	37,121	20,869
<b>Neomysis americana</b>	7	0	0	0	0	53	0	3	0

Note: Event 1 occurred April 9-18.  
 Event 2 occurred April 23-May 2.  
 Event 3 occurred May 6-21.  
 Event 4 occurred May 22-30.  
 Event 5 occurred June 7-18.  
 Event 6 occurred June 24-29.  
 Event 7 occurred July 8-17.  
 Event 8 occurred July 23-30.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-45 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 13) for each life stage of each target  
 species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	128.71	0.95	700.15	1.61	86.00	220.23	11.26	8.92	9.06
juvenile	0.20	0.00	0.00	0.00	0.00	0.00	0.00	1.57	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	6.41	11.60	10.47	0.00	0.00	29.17	0.00	0.00	0.00
yolk sac larvae	61.63	5.21	417.04	38.94	0.00	31.00	0.84	0.00	0.00
post-yolk sac larvae	0.46	0.00	0.00	0.00	0.00	0.00	3.66	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	15.96	0.00	117.97	0.00	5.38	4.32	0.00	0.00	0.00
yolk sac larvae	30.43	0.00	201.12	16.71	14.71	10.91	0.00	0.00	0.00
post-yolk sac larvae	153.90	0.00	847.10	153.97	146.09	78.25	0.00	5.77	0.00
juvenile	0.20	0.00	0.00	0.00	0.00	0.00	0.00	1.62	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	3.46	0.00	0.00	0.00	0.00	27.67	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.10	0.00	0.00	0.00	0.00	0.00	0.00	0.79	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.10	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.81
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-45 (page 2 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 13) for each life stage of each target  
 species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	88.50	0.00	38.10	0.00	669.88	0.00	0.00	0.00	0.00
post-yolk sac larvae	43.03	0.00	41.08	0.00	145.55	100.18	32.95	5.48	31.01
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	207.96	0.00	19.13	0.00	202.70	1,429.00	5.75	7.10	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	32.72	0.00	0.00	0.00	0.00	261.75	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	0.33	0.00	0.00	0.80	1.82	0.00	0.00	0.00	0.00
yolk sac larvae	25.34	0.00	0.00	202.68	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.52	0.00	0.00	0.00	0.00	1.72	1.64	0.83	0.00
juvenile	0.11	0.00	0.00	0.00	0.00	0.00	0.00	0.84	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	32.05	8.70	2.73	2.63	14.94	227.44	0.00	0.00	0.00
yolk sac larvae	302.72	13.43	324.64	52.93	1,712.02	314.69	1.62	0.00	0.00
post-yolk sac larvae	251.45	0.00	939.96	7.57	199.01	662.85	122.21	78.50	1.51
juvenile	0.42	0.00	0.00	0.00	0.00	0.00	0.00	3.33	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>1,387</b>	<b>40</b>	<b>3,659</b>	<b>478</b>	<b>3,198</b>	<b>3,399</b>	<b>180</b>	<b>115</b>	<b>42</b>
<b>Gammarus spp.</b>	10,254	70	491	2,349	3,073	13,273	5,673	28,595	28,509
<b>Neomysis americana</b>	3	0	0	3	17	0	0	1	2

Note: Event 1 occurred April 9-18.  
 Event 2 occurred April 23-May 2.  
 Event 3 occurred May 6-21.  
 Event 4 occurred May 22-30.  
 Event 5 occurred June 7-18.  
 Event 6 occurred June 24-29.  
 Event 7 occurred July 8-17.  
 Event 8 occurred July 23-30.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-46 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 14) for each life stage of each target  
 species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	17.26	0.00	58.20	0.00	49.95	21.47	1.71	2.59	4.19
juvenile	0.11	0.00	0.00	0.00	0.00	0.00	0.89	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	45.50	15.63	348.34	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	57.77	0.00	174.07	22.27	182.85	82.95	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.73	0.00	1.79	0.80	3.28	0.00	0.00	0.00	0.00
post-yolk sac larvae	11.01	0.00	60.01	10.72	0.00	16.40	0.00	0.91	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.20	0.00	0.78	0.80	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.10	0.00	0.00	0.00	0.81	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-46 (page 2 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 14) for each life stage of each target  
 species caught during the Delaware Baywide Ichthyoplankton Effort, April-July 2002

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	13.37	0.00	105.28	0.00	1.70	0.00	0.00	0.00	0.00
post-yolk sac larvae	31.28	0.00	25.42	0.00	183.95	7.07	0.00	1.83	32.81
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	116.88	0.00	0.00	0.00	749.24	184.20	0.00	1.60	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.11	0.00	0.00	0.00	0.00	0.00	0.00	0.86	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.11	0.00	0.00	0.00	0.00	0.00	0.85	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	226.20	28.66	0.00	0.00	1,780.93	0.00	0.00	0.00	0.00
yolk sac larvae	260.00	13.41	77.55	260.63	1,394.46	333.95	0.00	0.00	0.00
post-yolk sac larvae	3.35	0.00	8.94	0.00	0.00	4.87	6.02	6.09	0.85
juvenile	0.11	0.00	0.00	0.00	0.00	0.00	0.89	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>784</b>	<b>58</b>	<b>860</b>	<b>295</b>	<b>4,347</b>	<b>651</b>	<b>10</b>	<b>14</b>	<b>38</b>
<b>Gammarus spp.</b>	<b>6,677</b>	<b>152</b>	<b>142</b>	<b>2,862</b>	<b>1,182</b>	<b>2,128</b>	<b>38,793</b>	<b>4,328</b>	<b>3,825</b>
<b>Neomysis americana</b>	<b>2</b>	<b>0</b>	<b>0</b>	<b>4</b>	<b>5</b>	<b>3</b>	<b>0</b>	<b>0</b>	<b>0</b>

Note: Event 1 occurred April 9-18.  
 Event 2 occurred April 23-May 2.  
 Event 3 occurred May 6-21.  
 Event 4 occurred May 22-30.  
 Event 5 occurred June 7-18.  
 Event 6 occurred June 24-29.  
 Event 7 occurred July 8-17.  
 Event 8 occurred July 23-30.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

# 2003 DATA

Table 4-33 (page 1 of 28)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 1) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic Croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	553.66	0.00	0.00	3,780.08	0.00	573.80	35.88	0.00	0.00
yolk sac larvae	1.19	0.00	0.00	5.26	4.51	0.00	0.00	0.00	0.00
post-yolk sac larvae	1.00	0.00	0.00	3.21	2.35	0.00	2.55	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.17	0.00	0.00	0.00	0.29	0.00	1.04	0.00	0.00
juvenile	0.04	0.00	0.00	0.00	0.00	0.00	0.29	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	27,948.25	0.00	0.00	1,132.80	30,692.63	72,242.18	26,748.43	63,878.49	29,087.50
yolk sac larvae	0.64	0.00	0.00	0.00	0.00	0.00	0.00	0.00	5.06
post-yolk sac larvae	695.64	0.00	0.00	0.00	0.00	0.28	26.72	2,908.75	2,579.65
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.68	0.32	0.00	0.00	0.30	1.98	1.37	1.47	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00



Table 4-33 (page 2 of 28)  
PSEG Estuary Enhancement Program  
Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 1) for each life stage  
of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	142.40	0.00	3.77	0.00	251.84	0.00	823.88	53.03	14.53
yolk sac larvae	0.07	0.00	0.00	0.00	0.00	0.00	0.00	0.59	0.00
post-yolk sac larvae	9.09	0.00	0.00	0.00	0.30	0.00	0.99	52.78	18.04
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>29,353</b>	<b>0</b>	<b>4</b>	<b>4,921</b>	<b>30,952</b>	<b>72,818</b>	<b>27,641</b>	<b>66,895</b>	<b>31,705</b>
<b>Gammarus spp.</b>	279	87	64	322	376	175	265	732	220
<b>Neomysis americana</b>	98,573	3,624	1,324	20,092	124,409	58,956	323,311	118,727	123,929

Note: Event 1 occurred April 1-14.  
Event 2 occurred April 15-24.  
Event 3 occurred May 6-15.  
Event 4 occurred May 19-27.  
Event 5 occurred June 2-10.  
Event 6 occurred June 17-25.  
Event 7 occurred July 1-15.  
Event 8 occurred July 15-25.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-34 (page 3 of 28)  
PSEG Estuary Enhancement Program  
Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 2) for each life stage  
of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic Croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	149.45	0.00	0.00	1,098.37	0.37	113.40	0.00	0.00	0.00
yolk sac larvae	0.89	0.00	0.00	6.25	0.86	0.00	0.00	0.00	0.00
post-yolk sac larvae	1.36	0.97	0.00	1.04	6.94	1.29	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	1.03	0.44	0.00	0.00	0.38	6.66	0.86	0.00	0.00
juvenile	0.10	0.00	0.00	0.00	0.00	0.00	0.85	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	23,517.46	0.00	0.00	0.00	9,375.70	35,432.24	124,206.37	11,344.44	9,352.28
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	1,363.06	0.00	0.00	0.00	0.00	0.89	134.46	941.48	9,979.11
juvenile	0.43	0.00	0.00	0.00	0.00	0.00	0.42	0.00	3.05
adult	5.72	0.51	0.00	2.20	22.56	8.87	0.92	8.45	0.38
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-34 (page 4 of 28)  
PSEG Estuary Enhancement Program  
Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 2) for each life stage  
of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	50.06	0.00	406.07	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	12.43	0.00	0.00	0.52	0.00	0.00	0.94	3.63	95.76
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>25,102</b>	<b>2</b>	<b>406</b>	<b>1,108</b>	<b>9,407</b>	<b>35,563</b>	<b>124,345</b>	<b>12,298</b>	<b>19,431</b>
<b>Gammarus spp.</b>	2,237	186	187	86	13,991	225	397	699	821
<b>Neomysis americana</b>	176,371	80,958	7,175	47,709	218,177	127,289	242,297	386,249	296,468

Note: Event 1 occurred April 1-14.  
Event 2 occurred April 15-24.  
Event 3 occurred May 6-15.  
Event 4 occurred May 19-27.  
Event 5 occurred June 2-10.  
Event 6 occurred June 17-25.  
Event 7 occurred July 1-15.  
Event 8 occurred July 15-25.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-35 (page 5 of 28)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 3) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.05	0.00	0.00	0.42	0.05	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.05	0.00	0.00	0.40	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic Croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	59.61	0.00	0.00	467.97	5.79	3.13	0.00	0.00	0.00
yolk sac larvae	1.02	0.00	0.00	3.38	4.34	0.00	0.47	0.00	0.00
post-yolk sac larvae	1.40	0.00	0.45	5.48	5.30	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.56	0.00	0.00	2.29	2.15	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	8,613.25	0.00	0.00	4,834.16	4,425.96	36,525.37	18,734.65	1,672.38	2,713.47
yolk sac larvae	0.05	0.00	0.00	0.42	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	1,219.15	0.00	0.00	0.48	0.00	0.43	3.31	4,635.29	5,113.70
juvenile	41.23	0.00	0.00	0.00	0.00	0.00	0.00	266.28	63.58
adult	3.60	0.00	0.00	23.40	4.19	0.81	0.43	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-35 (page 6 of 28)  
PSEG Estuary Enhancement Program  
Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 3) for each life stage  
of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.16	0.43	0.00	0.87	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	16.65	0.00	5.13	0.00	0.00	0.00	121.79	0.00	6.30
yolk sac larvae	0.39	0.00	0.00	0.00	0.00	0.00	0.00	3.13	0.00
post-yolk sac larvae	28.96	0.00	0.00	0.00	0.00	0.00	0.00	191.13	40.53
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>9,986</b>	<b>0</b>	<b>6</b>	<b>5,339</b>	<b>4,448</b>	<b>36,530</b>	<b>18,861</b>	<b>6,768</b>	<b>7,938</b>
<b>Gammarus spp.</b>	5,903	268	393	121	404	65	23,395	2,851	17,981
<b>Neomysis americana</b>	154,467	78,277	111,753	101,046	106,151	26,991	173,338	353,283	284,894

Note: Event 1 occurred April 1-14.  
Event 2 occurred April 15-24.  
Event 3 occurred May 6-15.  
Event 4 occurred May 19-27.  
Event 5 occurred June 2-10.  
Event 6 occurred June 17-25.  
Event 7 occurred July 1-15.  
Event 8 occurred July 15-25.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-36 (page 7 of 28)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 4) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic Croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	48.10	0.00	0.00	320.42	64.38	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	2.09	0.38	0.00	0.00	15.11	0.89	0.34	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.91	0.00	0.00	0.00	2.06	1.21	0.69	3.33	0.00
juvenile	0.08	0.00	0.00	0.00	0.00	0.00	0.67	0.00	0.00
adult	0.55	0.40	0.00	0.00	0.00	0.00	0.34	0.43	3.25
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	7,589.86	0.00	0.00	32.45	7,486.21	21,788.54	7,199.77	21,870.90	2,341.02
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	952.14	0.00	0.00	0.00	0.00	0.00	157.84	3,687.95	3,771.33
juvenile	27.60	0.00	0.00	0.00	0.00	0.00	0.00	0.00	220.83
adult	7.08	0.00	0.44	6.95	35.29	2.59	1.81	5.63	3.90
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.05	0.00	0.00	0.00	0.00	0.39	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-36 (page 8 of 28)  
PSEG Estuary Enhancement Program  
Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 4) for each life stage  
of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	16.90	0.00	5.21	0.42	0.00	0.00	0.00	129.54	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	3.96	0.00	0.00	0.00	0.00	0.00	5.17	18.63	7.90
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>8,649</b>	<b>1</b>	<b>6</b>	<b>360</b>	<b>7,603</b>	<b>21,794</b>	<b>7,367</b>	<b>25,716</b>	<b>6,348</b>
<b>Gammarus spp.</b>	406	125	117	155	765	148	793	1,018	129
<b>Neomysis americana</b>	97,138	23,908	15,443	251,430	79,626	63,736	91,817	194,767	56,381

Note: Event 1 occurred April 1-14.  
Event 2 occurred April 15-24.  
Event 3 occurred May 6-15.  
Event 4 occurred May 19-27.  
Event 5 occurred June 2-10.  
Event 6 occurred June 17-25.  
Event 7 occurred July 1-15.  
Event 8 occurred July 15-25.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-37 (page 9 of 28)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 5) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic Croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.08	0.00	0.00	0.67	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.63	0.00	0.00	1.03	2.34	0.53	1.16	0.00	0.00
juvenile	0.40	3.19	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.07	0.60	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	3.79	0.00	0.00	0.00	8.26	0.66	3.60	11.10	6.69
juvenile	0.35	0.54	0.00	0.00	0.00	0.00	2.26	0.00	0.00
adult	0.92	1.65	0.00	0.00	0.00	0.00	0.53	0.56	4.66
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	203.95	0.00	0.00	0.00	26.41	2.00	100.11	1,212.24	290.84
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	734.43	0.00	0.00	0.00	0.00	0.00	264.56	3,561.41	2,049.45
juvenile	44.56	0.00	0.00	0.00	0.00	0.00	0.00	0.00	356.46
adult	23.46	1.62	0.52	22.42	120.39	11.45	7.28	22.76	1.22
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00



Table 4-37 (page 10 of 28)  
PSEG Estuary Enhancement Program  
Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 5) for each life stage  
of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.75	0.00	0.00	0.00	0.00	0.00	6.00	0.00	0.00
juvenile	1.53	0.00	0.00	0.00	0.00	0.00	0.00	12.27	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.26	0.00	0.00	0.00	0.00	2.10	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	55.29	0.00	0.00	0.00	0.00	0.66	11.71	334.40	95.56
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>1,070</b>	<b>8</b>	<b>1</b>	<b>24</b>	<b>157</b>	<b>17</b>	<b>397</b>	<b>5,155</b>	<b>2,805</b>
<b>Gammarus spp.</b>	1,028	47	60	14	3,364	324	3,220	536	659
<b>Neomysis americana</b>	168,755	162,507	700,082	234,705	44,737	56,186	110,717	25,946	15,164

Note: Event 1 occurred April 1-14.  
Event 2 occurred April 15-24.  
Event 3 occurred May 6-15.  
Event 4 occurred May 19-27.  
Event 5 occurred June 2-10.  
Event 6 occurred June 17-25.  
Event 7 occurred July 1-15.  
Event 8 occurred July 15-25.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-38 (page 11 of 28)  
PSEG Estuary Enhancement Program  
Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 6) for each life stage  
of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.15	0.00	0.00	0.00	0.00	1.23	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic Croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.24	0.00	0.00	1.93	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.14	0.00	0.00	0.00	1.16	0.00	0.00	0.00	0.00
post-yolk sac larvae	1.03	0.00	1.23	0.00	7.01	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	3.77	0.00	0.00	0.00	3.15	3.69	0.00	14.67	8.63
juvenile	0.14	0.00	0.00	0.00	0.00	0.00	1.11	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	123.18	0.00	0.00	0.00	6.99	1.23	0.00	0.00	977.20
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	460.83	0.00	0.00	0.00	0.00	0.00	0.00	941.39	2,745.26
juvenile	0.31	0.00	0.00	0.00	0.00	0.00	0.00	0.00	2.45
adult	6.98	0.00	0.00	20.91	22.60	7.30	0.00	2.79	2.28
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.12	0.00	0.00	0.00	0.00	0.00	0.94	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-38 (page 12 of 28)  
PSEG Estuary Enhancement Program  
Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 6) for each life stage  
of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	0.58	0.00	0.00	1.14	3.47	0.00	0.00	0.00	0.00
yolk sac larvae	1.24	0.00	0.00	2.94	7.01	0.00	0.00	0.00	0.00
post-yolk sac larvae	9.82	0.00	0.00	0.00	26.80	3.67	32.56	15.57	0.00
juvenile	1.05	0.00	0.00	0.00	0.00	0.00	1.11	7.33	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	17.85	0.00	0.00	0.00	0.00	0.00	0.00	61.74	81.07
juvenile	2.46	0.00	0.00	0.00	0.00	0.00	0.00	3.77	15.91
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>630</b>	<b>0</b>	<b>1</b>	<b>27</b>	<b>78</b>	<b>17</b>	<b>36</b>	<b>1,047</b>	<b>3,833</b>
<b>Gammarus spp.</b>	19,495	14,681	1,474	1,025	1,650	3,065	122,670	4,967	6,424
<b>Neomysis americana</b>	436,387	7,472	130,343	3,053,597	56,741	41,804	77,677	86,081	37,383

Note: Event 1 occurred April 1-14.  
Event 2 occurred April 15-24.  
Event 3 occurred May 6-15.  
Event 4 occurred May 19-27.  
Event 5 occurred June 2-10.  
Event 6 occurred June 17-25.  
Event 7 occurred July 1-15.  
Event 8 occurred July 15-25.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-39 (page 13 of 28)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 7) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.58	0.00	0.00	0.00	4.62	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic Croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.13	0.00	0.00	0.00	1.05	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.13	0.00	0.00	0.00	0.00	0.00	1.02	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.13	0.00	0.00	0.00	0.00	0.00	1.07	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	0.12	0.00	0.00	0.00	0.00	0.98	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	113.13	0.00	0.00	0.00	0.00	0.00	0.00	221.16	683.86
juvenile	0.65	0.00	0.00	0.00	0.00	0.98	0.00	0.00	4.23
adult	1.23	0.00	0.00	2.16	5.62	0.98	0.00	0.00	1.07
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.12	0.00	0.00	0.00	0.00	0.00	0.98	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-39 (page 14 of 28)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 7) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.28	0.00	0.00	0.00	0.00	2.21	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.48	0.00	0.00	3.82	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	11.30	0.00	0.00	0.00	0.00	17.67	0.00	72.74	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	65.03	0.00	0.00	517.90	0.00	2.33	0.00	0.00	0.00
yolk sac larvae	156.75	0.00	0.00	1,216.26	36.85	0.00	0.93	0.00	0.00
post-yolk sac larvae	110.79	0.00	0.00	0.00	404.35	163.13	69.19	247.33	2.31
juvenile	1.95	0.00	0.00	0.00	0.00	0.00	1.07	7.56	6.94
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	11.77	0.00	0.00	0.00	0.00	0.00	0.00	0.00	94.15
juvenile	2.21	0.00	0.00	0.00	0.00	0.00	0.00	0.00	17.65
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	2.77	0.00	0.00	22.14	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	9.24	0.00	0.00	17.57	2.38	44.71	4.01	5.28	0.00
juvenile	0.38	0.00	0.00	0.00	0.00	0.00	1.96	1.09	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>489</b>	<b>0</b>	<b>0</b>	<b>1,780</b>	<b>455</b>	<b>233</b>	<b>80</b>	<b>555</b>	<b>810</b>
<b>Gammarus spp.</b>	92,262	4,943	1,400	33,023	15,424	123,673	441,296	70,765	47,570
<b>Neomysis americana</b>	46,393	23	622	152,879	8,860	27,273	241	49,527	131,715

Note: Event 1 occurred April 1-14.  
 Event 2 occurred April 15-24.  
 Event 3 occurred May 6-15.  
 Event 4 occurred May 19-27.  
 Event 5 occurred June 2-10.  
 Event 6 occurred June 17-25.  
 Event 7 occurred July 1-15.  
 Event 8 occurred July 15-25.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-40 (page 15 of 28)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 8) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	99.45	0.00	0.00	67.55	561.83	85.58	37.57	43.10	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic Croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.52	0.00	0.00	0.00	0.00	0.00	0.00	0.00	4.18
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	0.61	0.00	0.00	0.00	0.00	0.00	0.00	4.86	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	5,050.46	0.00	0.00	0.00	0.00	0.00	0.00	39,824.23	579.47
juvenile	29.93	0.00	0.00	0.00	0.00	0.00	0.00	0.00	239.47
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-40 (page 16 of 28)  
PSEG Estuary Enhancement Program  
Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 8) for each life stage  
of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	16.92	0.00	0.00	0.00	57.93	9.51	37.57	25.86	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	1.08	0.00	0.00	0.00	0.00	0.00	0.00	8.62	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	109.12	0.00	0.00	858.56	9.65	4.75	0.00	0.00	0.00
yolk sac larvae	286.34	0.00	0.00	1,949.17	196.84	9.51	135.24	0.00	0.00
post-yolk sac larvae	66,035.43	0.00	0.00	0.00	8,522.94	1,779.38	393.41	517,579.36	8.35
juvenile	3,117.02	0.00	0.00	0.00	0.00	0.00	0.00	24,892.30	43.89
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	5.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	40.10
juvenile	4.58	0.00	0.00	0.00	0.00	0.00	0.00	0.00	36.63
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	1.65	0.00	8.54	4.67	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	11.94	0.00	0.00	95.52	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	10,859.30	0.00	0.00	180.86	1,847.08	2,813.33	2,137.51	79,874.76	20.88
juvenile	6,845.10	0.00	0.00	0.00	0.00	0.00	0.00	54,758.32	2.44
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>92,474</b>	<b>0</b>	<b>9</b>	<b>3,156</b>	<b>11,196</b>	<b>4,702</b>	<b>2,741</b>	<b>717,011</b>	<b>975</b>
<b>Gammarus spp.</b>	340,119	379	8,804	21,183	1,032,951	666,243	534,997	329,003	127,394
<b>Neomysis americana</b>	18,206	0	152	660	0	14	18	131	144,671

Note: Event 1 occurred April 1-14.  
Event 2 occurred April 15-24.  
Event 3 occurred May 6-15.  
Event 4 occurred May 19-27.  
Event 5 occurred June 2-10.  
Event 6 occurred June 17-25.  
Event 7 occurred July 1-15.  
Event 8 occurred July 15-25.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-41 (page 17 of 28)  
PSEG Estuary Enhancement Program  
Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 9) for each life stage  
of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	138.98	0.00	1.22	35.22	141.46	512.20	0.00	394.19	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	3.82	0.00	30.41	2.48	0.00	3.00	0.00	0.00	0.00
yolk sac larvae	3.10	0.00	8.39	16.61	0.00	0.87	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	1.69	0.00	0.00	0.00	0.00	0.00	0.00	0.00	13.22
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	2.20	0.00	0.00	0.98	0.00	9.54	1.62	5.05	0.00
juvenile	0.97	0.00	0.00	0.00	0.00	0.00	0.81	2.58	4.14
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic Croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.22	0.00	0.00	0.00	0.00	0.87	0.00	0.83	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	0.41	0.00	0.00	0.00	0.00	0.86	2.37	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	28.49	0.00	0.00	0.00	0.00	0.00	0.00	0.00	222.22
juvenile	6.37	0.00	0.00	0.00	0.00	0.00	0.00	0.00	49.71
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.54	0.00	0.00	0.00	0.00	0.00	0.00	0.00	4.17
juvenile	5.00	0.00	0.00	0.00	0.00	0.00	0.86	0.00	38.13
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00



Table 4-41 (page 18 of 28)  
PSEG Estuary Enhancement Program  
Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 9) for each life stage  
of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.12	0.00	1.22	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	19.09	0.00	0.00	5.47	62.65	30.38	0.00	50.39	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	9.25	0.00	0.00	0.00	0.00	0.00	0.86	71.28	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	203.68	0.00	15.49	17.64	1,546.21	10.75	1.73	0.00	0.00
yolk sac larvae	35.81	0.00	0.00	39.40	47.52	188.37	3.17	0.83	0.00
post-yolk sac larvae	156.43	0.00	0.00	0.00	629.54	400.41	1.59	181.05	7.54
juvenile	8.88	0.00	0.00	0.00	0.00	0.00	0.81	2.51	65.96
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	66.07	0.00	489.00	121.63	0.99	1.50	0.00	0.00	0.00
yolk sac larvae	12.19	0.00	1.22	49.96	1.93	42.23	0.00	0.00	0.00
post-yolk sac larvae	305.69	0.00	0.00	153.79	502.50	526.59	2.47	1,141.23	57.79
juvenile	12.36	0.00	0.00	0.00	0.00	0.00	0.00	18.09	78.35
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>1,021</b>	<b>0</b>	<b>547</b>	<b>443</b>	<b>2,933</b>	<b>1,728</b>	<b>16</b>	<b>1,868</b>	<b>541</b>
<b>Gammarus spp.</b>	125,130	51	170	1,296	84,806	323,908	31,622	124,799	409,394
<b>Neomysis americana</b>	211	0	8	12	0	1	1,621	1	6

Note: Event 1 occurred April 1-14.  
Event 2 occurred April 15-24.  
Event 3 occurred May 6-15.  
Event 4 occurred May 19-27.  
Event 5 occurred June 2-10.  
Event 6 occurred June 17-25.  
Event 7 occurred July 1-15.  
Event 8 occurred July 15-25.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-42 (page 19 of 28)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 10) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	266.61	0.00	0.85	226.13	1,172.29	279.81	391.76	60.43	1.63
juvenile	0.10	0.00	0.00	0.00	0.00	0.00	0.00	0.83	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	6.66	8.13	9.48	19.28	12.32	4.04	0.00	0.00	0.00
yolk sac larvae	56.03	0.00	3.51	432.14	9.12	1.78	0.00	1.68	0.00
post-yolk sac larvae	0.20	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.61
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.11	0.00	0.00	0.85	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.86	0.00	0.00	1.84	0.00	5.04	0.00	0.00	0.00
post-yolk sac larvae	5.24	0.00	0.00	5.55	6.77	21.18	3.48	4.96	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic Croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.11	0.00	0.00	0.85	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	1.33	0.00	0.00	0.00	0.00	0.00	0.00	0.00	10.67
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.41	0.00	0.00	0.00	0.00	0.00	0.00	0.84	2.41
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-42 (page 20 of 28)  
PSEG Estuary Enhancement Program  
Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 10) for each life stage  
of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	33.57	0.00	0.00	60.38	100.03	17.44	57.21	21.65	6.50
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.98	0.00	0.00	7.86	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	61.80	0.00	0.00	7.36	472.22	14.81	0.00	0.00	0.00
yolk sac larvae	30.51	0.00	0.00	67.15	112.17	3.16	59.18	2.45	0.00
post-yolk sac larvae	169.76	0.00	0.00	0.00	1,253.39	9.49	87.03	8.20	0.00
juvenile	0.10	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.81
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.11	0.00	0.00	0.00	0.01	0.89	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	39.25	2.44	78.70	182.98	46.67	3.23	0.00	0.00	0.00
yolk sac larvae	27.02	0.00	0.00	104.84	48.50	27.49	15.66	19.62	0.00
post-yolk sac larvae	435.81	0.00	0.00	375.76	497.04	174.95	2,245.69	166.14	26.89
juvenile	0.21	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.71
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>1,137</b>	<b>11</b>	<b>93</b>	<b>1,493</b>	<b>3,731</b>	<b>563</b>	<b>2,860</b>	<b>287</b>	<b>52</b>
<b>Gammarus spp.</b>	47,626	17	49	1,437	29,494	55,213	54,786	16,458	223,553
<b>Neomysis americana</b>	3	0	1	6	4	6	1	0	3

Note: Event 1 occurred April 1-14.  
Event 2 occurred April 15-24.  
Event 3 occurred May 6-15.  
Event 4 occurred May 19-27.  
Event 5 occurred June 2-10.  
Event 6 occurred June 17-25.  
Event 7 occurred July 1-15.  
Event 8 occurred July 15-25.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-43 (page 21 of 28)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 11) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	489.53	0.00	2.53	1,190.32	652.02	1,773.00	195.43	98.70	4.24
juvenile	0.84	0.00	0.00	0.00	0.00	0.00	0.00	6.72	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	38.23	0.00	67.47	220.53	14.46	3.41	0.00	0.00	0.00
yolk sac larvae	451.06	0.86	5.69	3,564.95	17.62	19.38	0.00	0.00	0.00
post-yolk sac larvae	2.11	0.00	0.00	0.00	0.00	0.00	0.00	0.00	16.85
juvenile	0.11	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.86
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.46	0.00	0.86	0.94	1.89	0.00	0.00	0.00	0.00
yolk sac larvae	24.79	0.00	0.00	85.65	10.35	102.29	0.00	0.00	0.00
post-yolk sac larvae	53.40	0.00	0.00	47.96	12.62	302.40	34.09	29.21	0.90
juvenile	1.67	0.00	0.00	0.00	0.00	0.00	0.82	6.72	5.83
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic Croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	2.19	0.00	0.00	17.52	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.12	0.00	0.00	0.00	0.99	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	0.23	0.00	0.00	0.00	0.00	1.87	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.30	0.00	0.00	0.00	0.00	0.00	0.00	0.00	2.43
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	5.86	0.00	0.00	0.00	0.00	0.00	0.00	0.81	46.08
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-43 (page 22 of 28)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 11) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	93.78	0.00	0.00	490.12	78.59	67.45	19.02	16.59	14.92
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	37.01	0.00	0.00	5.72	212.39	75.17	2.76	0.00	0.00
yolk sac larvae	11.85	0.00	0.00	57.36	32.43	0.00	3.29	1.68	0.00
post-yolk sac larvae	54.83	0.00	0.00	4.23	410.40	17.30	0.00	5.15	1.57
juvenile	0.32	0.00	0.00	0.00	0.00	0.00	0.00	0.00	2.59
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	35.91	0.00	14.15	199.47	63.83	6.96	2.87	0.00	0.00
yolk sac larvae	107.42	0.00	0.00	212.88	27.22	549.69	53.67	15.94	0.00
post-yolk sac larvae	481.82	0.00	0.00	737.81	415.33	874.48	1,073.22	686.82	66.89
juvenile	0.32	0.00	0.00	0.00	0.00	0.00	0.00	0.00	2.58
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>1,894</b>	<b>1</b>	<b>91</b>	<b>6,835</b>	<b>1,950</b>	<b>3,793</b>	<b>1,385</b>	<b>868</b>	<b>166</b>
<b>Gammarus spp.</b>	<b>36,059</b>	<b>17</b>	<b>57</b>	<b>284</b>	<b>7,707</b>	<b>51,249</b>	<b>45,484</b>	<b>6,034</b>	<b>177,643</b>
<b>Neomysis americana</b>	<b>4</b>	<b>0</b>	<b>0</b>	<b>11</b>	<b>3</b>	<b>4</b>	<b>1</b>	<b>0</b>	<b>10</b>

Note: Event 1 occurred April 1-14.  
 Event 2 occurred April 15-24.  
 Event 3 occurred May 6-15.  
 Event 4 occurred May 19-27.  
 Event 5 occurred June 2-10.  
 Event 6 occurred June 17-25.  
 Event 7 occurred July 1-15.  
 Event 8 occurred July 15-25.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-44 (page 23 of 28)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 12) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	551.20	0.00	0.00	1,175.42	2,924.89	269.47	13.55	20.81	5.43
juvenile	0.34	0.00	0.00	0.00	0.00	0.00	0.00	2.72	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	132.57	0.00	72.85	714.61	269.34	3.75	0.00	0.00	0.00
yolk sac larvae	105.92	0.00	0.80	422.71	102.72	312.89	8.25	0.00	0.00
post-yolk sac larvae	2.56	0.00	0.00	0.00	0.00	0.00	0.00	0.00	20.48
juvenile	0.33	0.00	0.00	0.00	0.00	0.00	0.00	0.00	2.63
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	57.00	0.00	2.60	393.43	32.18	23.65	4.13	0.00	0.00
yolk sac larvae	35.36	0.00	0.00	148.89	102.27	29.06	2.64	0.00	0.00
post-yolk sac larvae	121.75	0.00	0.00	326.52	287.40	163.26	88.77	36.49	71.52
juvenile	1.77	0.00	0.00	0.00	1.77	0.00	0.00	3.42	10.77
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic Croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.11	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.90
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.11	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.89
juvenile	2.13	0.00	0.00	0.00	2.13	0.00	0.00	0.00	17.02
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-44 (page 24 of 28)  
PSEG Estuary Enhancement Program  
Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 12) for each life stage  
of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	42.58	0.00	0.80	127.93	179.19	11.71	17.05	0.00	9.06
juvenile	0.10	0.00	0.00	0.00	0.00	0.00	0.00	0.83	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	9.88	0.00	2.60	0.81	20.66	8.61	46.32	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	1.79	0.00	0.00	0.00	5.86	3.12	0.83	0.92	3.58
juvenile	0.11	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.88
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	37.91	0.00	13.86	102.56	166.41	16.85	3.58	0.00	0.00
yolk sac larvae	70.26	0.00	0.00	170.82	284.08	46.69	47.72	12.81	0.00
post-yolk sac larvae	303.47	0.00	0.00	89.32	570.07	243.91	3.65	756.08	764.76
juvenile	1.11	0.00	0.00	0.00	0.00	0.00	0.00	0.00	8.85
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>1,478</b>	<b>0</b>	<b>94</b>	<b>3,673</b>	<b>4,945</b>	<b>1,133</b>	<b>236</b>	<b>834</b>	<b>917</b>
<b>Gammarus spp.</b>	<b>38,349</b>	<b>11</b>	<b>27</b>	<b>197</b>	<b>1,958</b>	<b>18,549</b>	<b>11,924</b>	<b>9,393</b>	<b>264,730</b>
<b>Neomysis americana</b>	<b>7</b>	<b>0</b>	<b>1</b>	<b>50</b>	<b>1</b>	<b>3</b>	<b>0</b>	<b>1</b>	<b>0</b>

Note: Event 1 occurred April 1-14.  
Event 2 occurred April 15-24.  
Event 3 occurred May 6-15.  
Event 4 occurred May 19-27.  
Event 5 occurred June 2-10.  
Event 6 occurred June 17-25.  
Event 7 occurred July 1-15.  
Event 8 occurred July 15-25.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-45 (page 25 of 28)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 13) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	473.65	0.00	0.00	899.97	2,795.45	6.06	35.08	26.57	26.08
juvenile	0.21	0.00	0.00	0.00	0.00	0.00	0.00	1.67	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	11.84	3.28	0.00	8.49	8.57	38.49	35.91	0.00	0.00
yolk sac larvae	316.59	0.00	1.85	1,490.09	828.03	178.40	34.36	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	30.01	0.00	24.17	83.14	39.78	29.65	63.34	0.00	0.00
yolk sac larvae	65.40	0.00	0.00	355.28	136.02	6.66	24.41	0.81	0.00
post-yolk sac larvae	144.85	0.00	0.00	356.29	550.95	5.03	144.27	71.59	30.71
juvenile	1.67	0.00	0.00	0.00	0.00	0.00	0.00	4.38	8.99
adult	0.11	0.00	0.00	0.00	0.00	0.00	0.00	0.86	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic Croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.23	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.82
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.21	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.72
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00



Table 4-45 (page 26 of 28)  
PSEG Estuary Enhancement Program  
Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 13) for each life stage  
of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.21	0.00	0.00	1.71	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	17.91	0.00	0.00	4.36	88.18	2.05	10.56	7.47	32.10
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.64	0.00	0.00	0.00	0.00	0.00	0.00	5.16	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	1.39	0.00	0.00	0.00	0.00	8.22	2.89	0.00	0.00
yolk sac larvae	0.11	0.00	0.87	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.24	0.00	0.00	0.00	0.00	0.00	0.98	0.00	0.97
juvenile	0.11	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.85
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	19.70	0.00	0.00	26.84	48.13	77.04	5.57	0.00	0.00
yolk sac larvae	52.69	0.00	0.00	149.69	133.34	3.00	85.97	49.53	0.00
post-yolk sac larvae	37.71	0.00	0.00	0.00	20.40	0.00	1.97	40.75	238.59
juvenile	0.22	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.53
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>1,176</b>	<b>3</b>	<b>27</b>	<b>3,376</b>	<b>4,649</b>	<b>355</b>	<b>445</b>	<b>209</b>	<b>343</b>
<b>Gammarus spp.</b>	<b>8,056</b>	<b>18</b>	<b>23</b>	<b>273</b>	<b>2,691</b>	<b>1,055</b>	<b>3,548</b>	<b>6,166</b>	<b>50,676</b>
<b>Neomysis americana</b>	<b>357</b>	<b>3</b>	<b>0</b>	<b>33</b>	<b>0</b>	<b>9</b>	<b>10</b>	<b>1</b>	<b>2,804</b>

Note: Event 1 occurred April 1-14.  
Event 2 occurred April 15-24.  
Event 3 occurred May 6-15.  
Event 4 occurred May 19-27.  
Event 5 occurred June 2-10.  
Event 6 occurred June 17-25.  
Event 7 occurred July 1-15.  
Event 8 occurred July 15-25.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-46 (page 27 of 28)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 14) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	130.83	0.00	0.00	16.93	855.79	1.88	6.09	1.89	19.07
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	22.79	0.00	27.63	100.71	0.98	24.04	3.48	25.48	0.00
yolk sac larvae	326.41	0.00	0.00	368.68	1,598.65	637.25	6.71	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	1.93	0.00	0.00	3.48	2.45	7.53	1.95	0.00	0.00
yolk sac larvae	1.26	0.00	0.00	1.98	7.15	0.00	0.00	0.94	0.00
post-yolk sac larvae	34.74	0.00	0.00	0.00	231.82	1.02	0.87	9.29	34.95
juvenile	2.58	0.00	0.00	0.00	0.00	0.00	0.00	0.90	19.70
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic Croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.34	0.00	0.00	0.00	0.00	0.00	0.00	0.00	2.74
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.78	0.00	0.00	0.00	0.00	0.00	0.00	0.00	6.25
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-46 (page 28 of 28)  
PSEG Estuary Enhancement Program  
Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 14) for each life stage  
of each target species caught during the Ichthyoplankton Effort, April-July 2003

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	2.40	0.00	0.00	0.87	17.37	0.00	0.00	0.94	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	0.49	0.00	0.00	0.00	0.00	0.00	3.90	0.00	0.00
yolk sac larvae	0.12	0.00	0.00	0.00	0.00	0.00	0.00	0.97	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.12	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.92
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	63.03	0.00	0.00	162.80	205.40	37.71	98.32	0.00	0.00
yolk sac larvae	59.72	0.00	0.00	9.43	131.84	7.19	139.88	189.45	0.00
post-yolk sac larvae	4.43	0.00	0.00	0.87	30.08	1.79	0.00	0.90	1.79
juvenile	0.33	0.00	0.00	0.00	0.00	0.00	0.00	0.00	2.67
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>652</b>	<b>0</b>	<b>28</b>	<b>666</b>	<b>3,082</b>	<b>718</b>	<b>261</b>	<b>231</b>	<b>88</b>
<b>Gammarus spp.</b>	4,603	27	41	281	4,445	1,071	894	4,531	25,530
<b>Neomysis americana</b>	12	0	1	87	1	3	0	0	3

Note: Event 1 occurred April 1-14.  
Event 2 occurred April 15-24.  
Event 3 occurred May 6-15.  
Event 4 occurred May 19-27.  
Event 5 occurred June 2-10.  
Event 6 occurred June 17-25.  
Event 7 occurred July 1-15.  
Event 8 occurred July 15-25.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

# 2004 DATA

Table 4-33 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 1) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	230.75	0.00	8.56	1,148.30	418.79	270.35	0.00	0.00	0.00
yolk sac larvae	0.03	0.00	0.00	0.00	0.28	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.91	0.00	0.00	0.00	3.83	3.42	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.76	0.00	0.00	0.00	6.05	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.17	0.00	0.00	0.00	0.28	0.82	0.00	0.30	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	66,811.36	0.00	0.00	2.96	121,447.44	367,156.82	11,071.54	32,429.71	2,382.44
yolk sac larvae	0.11	0.00	0.00	0.00	0.00	0.84	0.00	0.00	0.00
post-yolk sac larvae	5,048.84	0.00	0.00	0.00	1.39	152.58	6,916.87	3,801.20	29,518.71
juvenile	2.68	0.00	0.00	0.00	0.00	0.00	0.00	0.00	21.45
adult	0.34	0.00	0.00	0.00	2.22	0.54	0.00	0.00	0.00
undetermined larvae	190.47	0.00	0.00	0.00	1,523.76	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-33 (page 2 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 1) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.57	0.00	0.00	0.30	4.28	0.00	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	6,552.68	0.00	1,150.28	4,411.03	0.00	224.44	2,967.63	42,114.82	1,553.23
yolk sac larvae	0.49	0.00	0.00	0.00	0.00	0.00	0.00	3.95	0.00
post-yolk sac larvae	57.75	0.00	0.00	0.00	0.28	11.48	84.89	127.99	237.36
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.28	0.00	0.00	0.00	0.00	0.00	0.00	2.20	0.00
<b>White perch</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>78,898</b>	<b>0</b>	<b>1,159</b>	<b>5,563</b>	<b>123,409</b>	<b>367,821</b>	<b>21,041</b>	<b>78,480</b>	<b>33,713</b>
<b>Gammarus spp.</b>	442	162	25	187	505	317	243	1,484	616
<b>Neomysis americana</b>	49,867	19,367	6,553	78,625	31,801	64,057	60,231	85,590	52,714

Note: Event 1 occurred April 1-15.

Event 2 occurred April 19-27.

Event 3 occurred May 3-7.

Event 4 occurred May 13-20.

Event 5 occurred June 1-8.

Event 6 occurred June 14-25.

Event 7 occurred July 1-10.

Event 8 occurred July 19-28.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-34 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 2) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	31.72	0.00	0.00	253.30	0.42	0.00	0.00	0.00	0.00
yolk sac larvae	0.05	0.00	0.00	0.00	0.00	0.43	0.00	0.00	0.00
post-yolk sac larvae	0.79	0.00	0.00	0.00	4.18	2.13	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.16	0.00	0.00	0.00	1.28	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.05	0.00	0.00	0.00	0.42	0.00	0.00	0.00	0.00
post-yolk sac larvae	2.46	0.00	0.00	0.00	13.36	3.02	2.31	0.97	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.06	0.44	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	30,213.70	0.00	0.00	4.84	79,149.75	109,967.01	9,359.32	27,882.37	15,346.29
yolk sac larvae	0.65	0.00	0.00	0.00	0.00	1.35	0.00	3.86	0.00
post-yolk sac larvae	3,471.64	0.00	0.00	0.00	0.87	604.02	13,306.27	6,366.00	7,495.94
juvenile	5.53	0.00	0.00	0.00	0.00	0.00	0.00	0.00	44.25
adult	5.76	0.00	0.50	12.17	31.20	0.89	0.00	0.43	0.87
undetermined larvae	0.68	0.00	0.00	0.00	5.42	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.05	0.00	0.00	0.00	0.42	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-34 (page 2 of 2)  
PSEG Estuary Enhancement Program  
Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 2) for each life stage  
of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	878.09	0.00	375.18	986.42	0.00	0.00	295.02	3,706.62	1,661.51
yolk sac larvae	0.48	0.00	0.00	0.00	0.00	0.00	0.00	3.86	0.00
post-yolk sac larvae	117.76	0.00	0.00	0.00	6.45	13.64	429.09	295.89	197.05
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>34,730</b>	<b>0</b>	<b>376</b>	<b>1,257</b>	<b>79,214</b>	<b>110,592</b>	<b>23,392</b>	<b>38,260</b>	<b>24,746</b>
<b>Gammarus spp.</b>	1,868	488	72	578	421	283	3,378	3,532	6,193
<b>Neomysis americana</b>	77,354	73,687	36,658	287,156	71,877	68,912	17,474	7,725	55,345

Note: Event 1 occurred April 1-15.  
Event 2 occurred April 19-27.  
Event 3 occurred May 3-7.  
Event 4 occurred May 13-20.  
Event 5 occurred June 1-8.  
Event 6 occurred June 14-25.  
Event 7 occurred July 1-10.  
Event 8 occurred July 19-28.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.



Table 4-35 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 3) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	143.34	0.00	0.00	161.16	25.50	960.03	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	2.25	0.00	0.00	0.82	14.71	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.11	0.00	0.00	0.00	0.00	0.84	0.00	0.00	0.00
post-yolk sac larvae	4.48	0.00	0.00	3.23	6.08	17.18	0.00	9.32	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	18,541.81	0.00	0.00	0.83	9,774.21	131,195.29	6,488.37	783.55	92.22
yolk sac larvae	3.75	0.00	0.00	0.00	0.00	23.80	6.19	0.00	0.00
post-yolk sac larvae	5,708.95	0.00	0.00	0.00	0.00	3,687.23	15,585.06	13,303.00	13,096.31
juvenile	58.72	0.00	0.00	0.00	0.00	0.00	0.00	112.65	357.08
adult	3.54	0.00	0.00	2.04	24.57	1.71	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-35 (page 2 of 2)  
PSEG Estuary Enhancement Program  
Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 3) for each life stage  
of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.05	0.00	0.00	0.00	0.40	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	3,771.01	0.00	1,820.92	534.37	102.45	21,765.37	5,242.59	418.33	284.07
yolk sac larvae	4.45	0.00	0.00	0.00	0.36	35.22	0.00	0.00	0.00
post-yolk sac larvae	214.59	0.00	0.00	0.00	0.00	706.79	723.53	130.09	156.31
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>28,457</b>	<b>0</b>	<b>1,821</b>	<b>702</b>	<b>9,948</b>	<b>158,393</b>	<b>28,046</b>	<b>14,757</b>	<b>13,986</b>
<b>Gammarus spp.</b>	338	67	146	104	113	25	88	949	1,209
<b>Neomysis americana</b>	59,213	10,277	23,256	206,814	29,424	20,511	42,503	120,588	20,332

Note: Event 1 occurred April 1-15.  
Event 2 occurred April 19-27.  
Event 3 occurred May 3-7.  
Event 4 occurred May 13-20.  
Event 5 occurred June 1-8.  
Event 6 occurred June 14-25.  
Event 7 occurred July 1-10.  
Event 8 occurred July 19-28.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-36 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 4) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	34.59	0.00	0.00	276.69	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	2.26	0.00	0.00	0.00	11.63	0.00	6.47	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	2.38	0.00	0.00	0.41	5.60	3.41	9.59	0.00	0.00
juvenile	1.24	0.00	0.00	0.00	0.00	0.00	6.47	3.02	0.41
adult	1.29	0.00	0.00	0.00	0.00	0.00	0.00	3.21	7.08
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	10,550.34	0.00	0.00	0.00	27,425.85	52,182.40	1,191.87	46.54	3,556.04
yolk sac larvae	1.42	0.00	0.00	0.00	0.00	11.36	0.00	0.00	0.00
post-yolk sac larvae	2,500.80	0.00	0.00	0.00	0.39	2,184.59	11,566.53	3,102.37	3,152.57
juvenile	67.28	0.00	0.00	0.00	0.00	0.00	0.00	184.21	354.07
adult	4.92	0.00	10.65	0.85	25.82	1.21	0.38	0.41	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.05	0.00	0.00	0.00	0.00	0.39	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-36 (page 2 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 4) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.05	0.00	0.00	0.44	0.00	0.00	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	91.54	0.00	81.83	439.20	0.00	0.00	7.16	0.00	204.12
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	27.62	0.00	0.00	0.00	0.44	90.32	102.31	18.55	9.31
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>13,286</b>	<b>0</b>	<b>92</b>	<b>718</b>	<b>27,470</b>	<b>54,474</b>	<b>12,891</b>	<b>3,358</b>	<b>7,284</b>
<b>Gammarus spp.</b>	693	30	226	50	29	73	807	1,020	3,313
<b>Neomysis americana</b>	74,170	49,459	125,445	215,777	87,450	58,332	14,746	24,145	18,005

Note: Event 1 occurred April 1-15.

Event 2 occurred April 19-27.

Event 3 occurred May 3-7.

Event 4 occurred May 13-20.

Event 5 occurred June 1-8.

Event 6 occurred June 14-25.

Event 7 occurred July 1-10.

Event 8 occurred July 19-28.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-37 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 5) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.08	0.00	0.00	0.61	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.59	0.62	0.00	0.58	3.52	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.07	0.00	0.00	0.58	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	7.03	0.00	0.00	13.22	19.89	7.80	10.81	1.15	3.36
juvenile	0.14	0.00	0.00	0.00	0.00	0.00	0.58	0.56	0.00
adult	0.78	0.00	0.62	0.00	0.00	1.14	0.60	1.72	2.18
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	2,591.10	0.00	0.00	0.00	15,784.31	2,121.65	2,269.28	165.77	387.78
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	859.89	0.00	0.00	0.00	0.60	1,539.40	2,731.23	1,047.13	1,560.74
juvenile	95.55	0.00	0.00	0.00	0.00	0.00	0.00	161.80	602.58
adult	13.73	0.62	15.07	4.98	42.76	2.48	7.74	34.51	1.68
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-37 (page 2 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 5) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.23	1.85	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.08	0.00	0.60	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	29.42	0.00	0.00	0.00	0.00	188.26	16.69	7.23	23.22
juvenile	0.25	0.00	0.00	0.00	0.00	0.00	0.00	2.02	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>3,599</b>	<b>3</b>	<b>16</b>	<b>20</b>	<b>15,851</b>	<b>3,861</b>	<b>5,037</b>	<b>1,422</b>	<b>2,582</b>
<b>Gammarus spp.</b>	210	47	187	101	21	544	233	363	187
<b>Neomysis americana</b>	127,455	280,538	485,140	63,434	73,408	29,109	57,520	20,853	9,637

Note: Event 1 occurred April 1-15.

Event 2 occurred April 19-27.

Event 3 occurred May 3-7.

Event 4 occurred May 13-20.

Event 5 occurred June 1-8.

Event 6 occurred June 14-25.

Event 7 occurred July 1-10.

Event 8 occurred July 19-28.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-38 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 6) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	2.06	1.11	0.00	0.00	15.38	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	7.32	0.00	0.00	0.00	35.61	15.64	0.00	7.31	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.26	0.00	0.00	0.00	1.16	0.00	0.00	0.89	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	457.07	0.00	0.00	0.00	548.31	2,144.30	296.79	517.13	150.05
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	277.93	0.00	0.00	0.00	3.49	1,304.14	622.93	209.51	83.40
juvenile	16.52	0.00	0.00	0.00	0.00	0.00	0.00	40.21	91.95
adult	8.24	0.00	2.06	16.79	25.13	5.67	10.44	1.84	3.99
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-38 (page 2 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 6) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.30	0.00	0.00	0.00	2.36	0.00	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.12	0.00	0.00	0.00	0.98	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.15	0.00	0.00	0.00	1.16	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.15	0.00	0.00	0.00	0.00	0.00	1.18	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	16.52	0.00	0.00	0.00	1.16	37.81	72.31	2.70	18.22
juvenile	0.69	0.00	0.00	0.00	0.00	1.15	0.00	2.19	2.15
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.15	0.00	0.00	0.00	1.18	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.26	0.00	2.07	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>788</b>	<b>1</b>	<b>4</b>	<b>17</b>	<b>636</b>	<b>3,509</b>	<b>1,004</b>	<b>782</b>	<b>350</b>
<b>Gammarus spp.</b>	4,007	439	20,953	1,410	2,906	1,203	1,870	396	2,881
<b>Neomysis americana</b>	73,868	327,875	130,101	17,576	50,979	14,387	4,229	12,641	33,159

Note: Event 1 occurred April 1-15.

Event 2 occurred April 19-27.

Event 3 occurred May 3-7.

Event 4 occurred May 13-20.

Event 5 occurred June 1-8.

Event 6 occurred June 14-25.

Event 7 occurred July 1-10.

Event 8 occurred July 19-28.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.



Table 4-39 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 7) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.15	0.00	1.17	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	1.50	0.00	0.00	0.00	12.01	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	2.68	0.00	0.00	0.00	6.05	1.21	0.00	14.15	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.43	0.00	0.00	0.00	1.08	0.00	2.40	0.00	0.00
<b>Bay anchovy</b>									
egg	1.95	0.00	0.00	0.00	10.82	4.74	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	160.95	0.00	0.00	0.00	0.00	296.11	623.87	236.28	131.37
juvenile	10.20	0.00	0.00	0.00	0.00	0.00	0.00	21.07	60.57
adult	1.73	0.00	0.00	3.41	6.94	0.00	2.28	1.24	0.00
undetermined larvae	4.22	0.00	0.00	0.00	0.00	33.80	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-39 (page 2 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 7) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.13	0.00	0.00	0.00	0.00	0.00	1.07	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	4.37	0.00	0.00	0.00	34.93	0.00	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	115.68	0.00	0.00	10.49	893.27	21.69	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	207.86	0.00	1,661.87	1.03	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.15	0.00	0.00	1.21	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	70.39	0.00	0.00	0.00	469.63	86.88	6.62	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	54.00	0.00	0.00	0.00	0.00	89.86	320.14	11.95	10.05
juvenile	15.98	0.00	0.00	0.00	0.00	2.74	121.84	1.24	2.02
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	0.96	0.00	4.13	0.00	0.00	3.58	0.00	0.00	0.00
yolk sac larvae	1.46	0.00	11.69	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	6.72	0.00	0.00	0.00	53.74	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.73	1.11	3.51	1.21	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>662</b>	<b>1</b>	<b>1,682</b>	<b>17</b>	<b>1,488</b>	<b>541</b>	<b>1,078</b>	<b>286</b>	<b>204</b>
<b>Gammarus spp.</b>	88,366	20,702	104,348	80,379	295,645	19,429	27,749	39,543	119,132
<b>Neomysis americana</b>	30,200	43,982	68	303	613	27,333	25,747	104,242	39,310

Note: Event 1 occurred April 1-15.

Event 2 occurred April 19-27.

Event 3 occurred May 3-7.

Event 4 occurred May 13-20.

Event 5 occurred June 1-8.

Event 6 occurred June 14-25.

Event 7 occurred July 1-10.

Event 8 occurred July 19-28.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-40 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 8) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	14.07	0.00	0.00	0.00	112.53	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.57	0.00	0.00	0.00	2.41	0.00	2.11	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	1.63	0.00	0.00	0.00	0.00	5.97	0.00	4.75	2.30
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	72.76	0.00	0.00	0.00	0.00	266.91	128.51	138.34	48.35
juvenile	4.51	0.00	0.00	0.00	0.00	0.00	0.00	11.08	25.02
adult	0.56	0.00	0.00	0.00	0.00	4.46	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.50	0.00	0.00	0.00	0.00	3.98	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-40 (page 2 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 8) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	4.97	0.00	0.00	0.00	37.51	0.00	0.00	2.22	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	69.81	0.00	0.00	0.00	558.51	0.00	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	273.54	0.00	0.00	930.43	1,222.13	35.73	0.00	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	384.47	0.00	3,063.62	12.10	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.59	0.00	0.00	0.00	4.69	0.00	0.00	0.00	0.00
post-yolk sac larvae	57.36	0.00	0.00	0.00	292.08	148.00	18.80	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	4.17	0.00	0.00	0.00	0.00	1.99	29.83	1.58	0.00
juvenile	1.02	0.00	0.00	0.00	0.00	0.00	6.61	1.58	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	14.17	0.00	110.94	2.42	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	66.96	0.00	0.00	19.36	463.98	45.20	7.15	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.31	0.00	0.00	0.00	0.00	2.47	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>972</b>	<b>0</b>	<b>3,175</b>	<b>964</b>	<b>2,694</b>	<b>515</b>	<b>193</b>	<b>160</b>	<b>76</b>
<b>Gammarus spp.</b>	267,270	2,330	5,104	111,964	1,566,651	45,625	101,271	153,411	151,805
<b>Neomysis americana</b>	7,137	2	0	5	0	20,097	1,313	35,672	10

Note: Event 1 occurred April 1-15.

Event 2 occurred April 19-27.

Event 3 occurred May 3-7.

Event 4 occurred May 13-20.

Event 5 occurred June 1-8.

Event 6 occurred June 14-25.

Event 7 occurred July 1-10.

Event 8 occurred July 19-28.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-41 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 9) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	58.57	0.00	0.00	146.62	321.91	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	0.30	0.00	2.42	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	1.69	0.00	0.81	10.89	1.83	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.40	0.00	0.00	0.00	0.00	2.36	0.00	0.82	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.11	0.00	0.00	0.91	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.85	0.00	0.00	0.00	6.80	0.00	0.00	0.00	0.00
juvenile	0.30	0.00	0.00	0.00	0.00	0.84	0.82	0.00	0.76
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.33	0.00	0.00	0.00	0.00	1.94	0.00	0.70	0.00
juvenile	0.31	0.00	0.00	0.00	0.00	1.60	0.86	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.10	0.00	0.83	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	34.93	0.00	0.00	0.00	0.00	0.00	72.14	139.83	67.45
juvenile	6.14	0.00	0.00	0.00	0.00	0.00	0.00	19.40	29.74
adult	0.20	0.00	0.00	0.00	0.00	1.58	0.00	0.00	0.00
undetermined larvae	1.24	0.00	0.00	0.00	0.00	0.00	9.90	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	1.15	0.00	0.00	0.00	0.00	4.47	1.60	0.83	2.27
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-41 (page 2 of 2)  
PSEG Estuary Enhancement Program  
Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 9) for each life stage  
of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.20	0.00	1.61	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	57.79	0.00	0.00	2.72	458.93	0.00	0.00	0.69	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	59.57	0.00	0.00	177.88	289.26	9.39	0.00	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	78.07	0.00	0.00	51.29	62.46	486.19	24.59	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	4.96	0.00	0.00	33.87	5.83	0.00	0.00	0.00	0.00
yolk sac larvae	17.82	0.00	0.00	24.07	118.47	0.00	0.00	0.00	0.00
post-yolk sac larvae	68.68	0.00	0.00	198.13	33.70	288.57	28.34	0.69	0.00
juvenile	0.36	0.00	0.00	0.00	0.00	0.00	0.80	2.08	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	12.27	1.63	66.82	28.83	0.92	0.00	0.00	0.00	0.00
yolk sac larvae	13.32	0.00	7.54	26.98	72.08	0.00	0.00	0.00	0.00
post-yolk sac larvae	229.24	0.00	0.00	410.19	646.44	745.08	31.31	0.00	0.90
juvenile	1.76	0.00	0.00	0.00	0.00	1.67	2.51	7.61	2.26
adult	0.44	0.00	0.00	0.90	0.00	0.97	1.64	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>651</b>	<b>2</b>	<b>80</b>	<b>1,113</b>	<b>2,019</b>	<b>1,545</b>	<b>174</b>	<b>173</b>	<b>103</b>
<b>Gammarus spp.</b>	227,946	371	2,264	146,756	68,591	375,855	996,070	129,995	103,663
<b>Neomysis americana</b>	8	0	0	4	3	24	1	31	0

Note: Event 1 occurred April 1-15.  
Event 2 occurred April 19-27.  
Event 3 occurred May 3-7.  
Event 4 occurred May 13-20.  
Event 5 occurred June 1-8.  
Event 6 occurred June 14-25.  
Event 7 occurred July 1-10.  
Event 8 occurred July 19-28.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-42 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 10) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	95.33	0.00	6.32	232.24	444.68	77.92	1.50	0.00	0.00
juvenile	0.66	0.00	0.00	0.00	0.00	0.00	1.50	3.79	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	11.78	0.97	4.27	5.55	83.47	0.00	0.00	0.00	0.00
yolk sac larvae	10.97	0.00	10.96	69.37	7.47	0.00	0.00	0.00	0.00
post-yolk sac larvae	2.96	0.00	0.00	0.00	0.00	22.16	1.55	0.00	0.00
juvenile	0.34	0.00	0.00	0.00	0.00	2.69	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.82	0.00	0.00	4.64	1.92	0.00	0.00	0.00	0.00
post-yolk sac larvae	10.02	0.00	0.00	1.83	52.76	25.55	0.00	0.00	0.00
juvenile	1.52	0.00	0.00	0.00	0.00	9.60	1.61	0.91	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.78	0.00	0.00	6.22	0.00	0.00	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.12	0.00	0.00	0.00	0.00	0.00	0.00	0.94	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	13.70	0.00	0.00	0.00	0.00	0.00	31.82	74.36	3.39
juvenile	1.28	0.00	0.00	0.00	0.00	0.00	0.00	8.51	1.70
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	1.11	0.00	0.00	0.00	0.00	4.07	4.84	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-42 (page 2 of 2)  
PSEG Estuary Enhancement Program  
Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 10) for each life stage  
of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	27.55	0.00	0.00	0.00	162.32	58.05	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	211.37	0.00	0.00	78.85	1,230.10	361.52	3.11	17.35	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.11	0.00	0.00	0.00	0.00	0.00	0.86	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	75.29	0.00	0.00	3.60	207.55	364.80	26.40	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	1.35	0.00	2.69	5.43	2.67	0.00	0.00	0.00	0.00
yolk sac larvae	2.44	0.00	0.00	0.93	18.60	0.00	0.00	0.00	0.00
post-yolk sac larvae	72.44	0.00	0.00	0.00	18.43	547.13	12.06	1.92	0.00
juvenile	0.42	0.00	0.00	0.00	0.00	0.00	1.67	0.94	0.74
adult	0.09	0.00	0.00	0.00	0.00	0.00	0.75	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	18.04	0.00	56.78	17.66	69.06	0.84	0.00	0.00	0.00
yolk sac larvae	29.79	0.00	9.45	86.74	142.16	0.00	0.00	0.00	0.00
post-yolk sac larvae	281.37	0.00	0.00	123.09	928.74	1,164.68	34.43	0.00	0.00
juvenile	1.69	0.00	0.00	0.00	0.00	0.00	4.23	9.33	0.00
adult	0.34	0.00	1.83	0.00	0.00	0.90	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>874</b>	<b>1</b>	<b>92</b>	<b>636</b>	<b>3,370</b>	<b>2,640</b>	<b>126</b>	<b>118</b>	<b>6</b>
<b>Gammarus spp.</b>	346,700	153	910	5,471	29,167	1,406,435	612,808	357,946	360,711
<b>Neomysis americana</b>	2	0	0	19	0	0	0	0	0

Note: Event 1 occurred April 1-15.  
Event 2 occurred April 19-27.  
Event 3 occurred May 3-7.  
Event 4 occurred May 13-20.  
Event 5 occurred June 1-8.  
Event 6 occurred June 14-25.  
Event 7 occurred July 1-10.  
Event 8 occurred July 19-28.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.



Table 4-43 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 11) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	345.47	0.00	0.00	530.14	1,991.79	232.83	9.00	0.00	0.00
juvenile	0.29	0.00	0.00	0.00	0.00	0.00	2.29	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	50.42	0.00	13.12	71.39	318.88	0.00	0.00	0.00	0.00
yolk sac larvae	8.45	0.00	20.78	35.44	11.39	0.00	0.00	0.00	0.00
post-yolk sac larvae	5.97	0.00	0.00	0.00	0.00	36.39	8.64	2.70	0.00
juvenile	0.41	0.00	0.00	0.00	0.00	3.30	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	0.53	0.00	0.94	3.28	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	23.94	0.00	0.00	39.02	152.46	0.00	0.00	0.00	0.00
post-yolk sac larvae	44.95	0.00	0.00	54.13	189.01	114.96	1.53	0.00	0.00
juvenile	3.68	0.00	0.00	0.00	0.00	26.44	3.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	2.03	0.00	0.00	16.25	0.00	0.00	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.44	0.00	0.00	0.00	0.00	0.00	0.00	3.56	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	19.58	0.00	0.00	0.00	0.00	0.00	7.80	142.96	5.91
juvenile	0.46	0.00	0.00	0.00	0.00	0.00	0.00	3.68	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.19	0.00	0.00	0.00	0.00	0.00	1.51	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	1.05	0.00	0.00	0.00	0.00	1.73	2.35	1.79	2.56
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-43 (page 2 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 11) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	89.55	0.00	0.00	1.72	672.24	42.43	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	365.15	0.00	0.00	423.98	1,956.91	522.46	17.85	0.00	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	1.30	0.00	0.00	0.00	0.00	10.36	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	105.71	0.00	0.00	18.92	347.43	124.58	354.75	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	3.68	0.00	2.80	0.00	26.67	0.00	0.00	0.00	0.00
yolk sac larvae	6.59	0.00	0.00	0.00	52.75	0.00	0.00	0.00	0.00
post-yolk sac larvae	8.89	0.00	0.00	0.00	29.95	37.42	3.76	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	125.17	0.00	213.79	60.17	702.39	25.04	0.00	0.00	0.00
yolk sac larvae	38.30	0.00	11.63	56.68	237.19	0.86	0.00	0.00	0.00
post-yolk sac larvae	749.09	0.00	0.00	144.19	2,337.82	3,217.20	290.97	1.72	0.82
juvenile	0.43	0.00	0.00	0.00	0.00	2.53	0.88	0.00	0.00
adult	0.11	0.00	0.00	0.00	0.00	0.00	0.00	0.85	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>2,002</b>	<b>0</b>	<b>263</b>	<b>1,455</b>	<b>9,027</b>	<b>4,399</b>	<b>704</b>	<b>157</b>	<b>9</b>
<b>Gammarus spp.</b>	275,895	24	94	2,134	26,508	879,973	316,597	607,005	374,829
<b>Neomysis americana</b>	0	0	0	2	0	0	0	0	0

Note: Event 1 occurred April 1-15.

Event 2 occurred April 19-27.

Event 3 occurred May 3-7.

Event 4 occurred May 13-20.

Event 5 occurred June 1-8.

Event 6 occurred June 14-25.

Event 7 occurred July 1-10.

Event 8 occurred July 19-28.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-44 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 12) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	329.30	0.00	0.00	455.82	1,847.67	310.44	20.43	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	104.06	8.94	95.58	618.37	109.62	0.00	0.00	0.00	0.00
yolk sac larvae	11.76	0.00	34.04	44.45	15.55	0.00	0.00	0.00	0.00
post-yolk sac larvae	3.62	0.00	0.00	0.00	0.00	0.00	26.43	2.56	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	56.27	0.00	360.78	75.11	14.24	0.00	0.00	0.00	0.00
yolk sac larvae	82.71	0.00	14.34	238.18	409.14	0.00	0.00	0.00	0.00
post-yolk sac larvae	377.74	0.00	0.00	404.67	2,206.86	398.93	9.74	0.84	0.86
juvenile	1.98	0.00	0.00	0.00	0.00	15.83	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	16.80	0.00	0.00	134.39	0.00	0.00	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.10	0.00	0.00	0.00	0.00	0.00	0.00	0.78	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	4.36	0.00	0.00	0.00	0.00	0.00	4.38	22.70	7.76
juvenile	0.59	0.00	0.00	0.00	0.00	0.00	0.00	4.73	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	1.13	0.00	0.00	0.00	0.00	0.00	0.00	5.39	3.61
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-44 (page 2 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 12) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	43.47	0.00	0.00	0.00	282.41	23.84	41.48	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	722.74	0.00	3.17	355.47	5,043.41	315.83	59.75	4.30	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	187.00	0.00	0.00	0.00	765.54	562.43	168.06	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	7.76	0.00	0.87	0.00	61.20	0.00	0.00	0.00	0.00
yolk sac larvae	2.92	0.00	0.00	0.00	23.32	0.00	0.00	0.00	0.00
post-yolk sac larvae	8.72	0.00	0.00	0.00	7.84	53.48	8.44	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.20	0.00	0.00	0.00	0.00	0.00	1.60	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	64.09	7.61	127.11	266.03	111.99	0.00	0.00	0.00	0.00
yolk sac larvae	127.49	0.00	57.52	172.55	789.85	0.00	0.00	0.00	0.00
post-yolk sac larvae	1,712.93	0.00	0.00	55.28	5,862.90	6,929.56	850.37	5.34	0.00
juvenile	0.11	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.92
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>3,868</b>	<b>17</b>	<b>693</b>	<b>2,820</b>	<b>17,552</b>	<b>8,610</b>	<b>1,191</b>	<b>47</b>	<b>13</b>
<b>Gammarus spp.</b>	150,966	13	66	1,728	16,547	205,377	212,560	375,652	395,787
<b>Neomysis americana</b>	0	0	0	0	0	0	0	0	0

Note: Event 1 occurred April 1-15.

Event 2 occurred April 19-27.

Event 3 occurred May 3-7.

Event 4 occurred May 13-20.

Event 5 occurred June 1-8.

Event 6 occurred June 14-25.

Event 7 occurred July 1-10.

Event 8 occurred July 19-28.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-45 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 13) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	383.80	0.00	0.86	55.21	932.75	1,744.32	337.25	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	38.58	2.48	3.76	300.77	1.61	0.00	0.00	0.00	0.00
yolk sac larvae	7.65	0.00	9.36	25.10	26.76	0.00	0.00	0.00	0.00
post-yolk sac larvae	3.12	0.00	0.00	0.00	0.00	0.00	18.62	6.31	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	20.69	0.00	95.56	62.69	7.28	0.00	0.00	0.00	0.00
yolk sac larvae	43.26	0.00	0.79	248.60	90.41	6.31	0.00	0.00	0.00
post-yolk sac larvae	222.60	0.00	0.00	199.67	423.55	1,124.90	31.74	0.95	0.00
juvenile	2.49	0.00	0.00	0.00	0.00	15.96	3.10	0.87	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	69.93	0.00	0.00	184.71	0.00	374.74	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	2.23	0.00	0.00	0.00	0.00	0.00	0.64	14.08	3.11
juvenile	0.21	0.00	0.00	0.00	0.00	0.00	0.00	0.86	0.78
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.42	0.00	0.00	0.00	0.00	0.00	0.00	0.00	3.34
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-45 (page 2 of 2)  
PSEG Estuary Enhancement Program  
Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 13) for each life stage  
of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	100.31	0.00	0.00	1.00	0.00	552.37	68.31	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	295.21	0.00	0.00	33.33	1,019.08	1,087.04	216.78	5.45	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.10	0.81	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	47.75	0.00	0.00	0.00	5.01	344.11	31.97	0.95	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	0.67	0.00	4.44	0.00	0.90	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	1.13	0.00	0.00	0.00	0.00	0.00	9.02	0.00	0.00
juvenile	0.12	0.00	0.00	0.00	0.00	0.00	0.00	0.95	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	36.08	1.65	62.56	220.88	3.52	0.00	0.00	0.00	0.00
yolk sac larvae	123.57	0.00	10.45	88.57	875.27	14.28	0.00	0.00	0.00
post-yolk sac larvae	462.94	0.00	0.00	3.98	18.50	3,272.34	408.70	0.00	0.00
juvenile	0.12	0.00	0.00	0.00	0.00	0.00	0.00	0.95	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>1,863</b>	<b>5</b>	<b>188</b>	<b>1,425</b>	<b>3,405</b>	<b>8,536</b>	<b>1,126</b>	<b>31</b>	<b>7</b>
<b>Gammarus spp.</b>	194,590	4	42	683	6,413	53,148	72,689	407,494	1,016,248
<b>Neomysis americana</b>	0	0	0	0	0	0	0	1	0

Note: Event 1 occurred April 1-15.  
Event 2 occurred April 19-27.  
Event 3 occurred May 3-7.  
Event 4 occurred May 13-20.  
Event 5 occurred June 1-8.  
Event 6 occurred June 14-25.  
Event 7 occurred July 1-10.  
Event 8 occurred July 19-28.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.

Table 4-46 (page 1 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 14) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Alewife</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	24.21	0.00	0.00	20.32	34.10	104.82	29.61	0.00	0.00
juvenile	0.10	0.00	0.00	0.00	0.00	0.00	0.77	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Alosa spp.</b>									
egg	7.65	21.84	7.53	33.89	0.77	0.00	0.00	0.00	0.00
yolk sac larvae	18.63	0.00	0.96	47.47	96.88	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.38	0.00	0.00	0.00	0.00	0.00	2.97	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>American shad</b>									
egg	2.54	0.00	17.25	2.58	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	3.48	0.00	0.00	22.11	5.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	84.21	0.00	0.00	5.13	14.86	605.23	31.63	0.00	0.00
juvenile	1.89	0.00	0.00	0.00	0.00	10.95	3.05	0.71	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.60	0.00	0.00	0.00	4.71	0.00	0.00	0.00	0.00
<b>Atlantic croaker</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic menhaden</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Atlantic silverside</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bay anchovy</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.54	0.00	0.00	0.00	0.00	0.00	0.00	4.23	0.00
juvenile	0.10	0.00	0.00	0.00	0.00	0.00	0.00	0.80	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Blueback herring</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	1.00	0.00	0.00	0.00	0.00	0.00	0.00	7.06	0.77
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Bluefish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 4-46 (page 2 of 2)

PSEG Estuary Enhancement Program  
 Mean density (#/1000 m<sup>3</sup>) by sampling event (collected in Zone 14) for each life stage  
 of each target species caught during the Ichthyoplankton Effort, April-July 2004

Species (Taxon) by Life Stage	All Events*	April		May		June		July	
		Event 1	Event 2	Event 3	Event 4	Event 5	Event 6	Event 7	Event 8
<b>Clupeidae</b>									
egg	0.21	0.00	0.00	0.00	0.00	0.00	0.00	1.60	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	2.99	0.00	0.00	2.54	0.83	16.17	3.80	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	37.03	0.00	0.00	21.99	66.95	138.68	59.82	1.42	0.00
<b>Morone spp.</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	3.46	0.00	0.00	0.00	0.00	4.59	22.43	0.00	0.00
<b>Spot</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Striped bass</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.19	0.00	0.00	0.00	0.00	0.70	0.77	0.00	0.00
juvenile	0.09	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.74
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Weakfish</b>									
egg	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
post-yolk sac larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
juvenile	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
adult	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>White perch</b>									
egg	26.63	1.03	82.12	114.31	8.35	2.09	0.00	0.00	0.00
yolk sac larvae	154.12	0.00	0.94	422.81	773.60	4.76	0.00	0.00	0.00
post-yolk sac larvae	13.19	0.00	0.00	0.00	0.00	92.59	8.01	1.51	0.77
juvenile	0.45	0.00	0.00	0.00	0.00	0.00	0.00	3.55	0.00
adult	0.10	0.00	0.00	0.00	0.00	0.00	0.00	0.75	0.00
undetermined larvae	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<b>Total IP</b>	<b>384</b>	<b>23</b>	<b>109</b>	<b>693</b>	<b>1,006</b>	<b>981</b>	<b>163</b>	<b>22</b>	<b>2</b>
<b>Gammarus spp.</b>	220,169	12	203	373	1,408	7,988	28,266	477,485	1,201,589
<b>Neomysis americana</b>	1	6	0	1	0	0	1	1	2

Note: Event 1 occurred April 1-15.

Event 2 occurred April 19-27.

Event 3 occurred May 3-7.

Event 4 occurred May 13-20.

Event 5 occurred June 1-8.

Event 6 occurred June 14-25.

Event 7 occurred July 1-10.

Event 8 occurred July 19-28.

\*The "All Events" column contains the mean density for all events combined. It is not the total across each row.