New Jersey Department of Health Children's Oral Health Education Program



Tooth Eruption

Primary Teeth	Upper Teeth	Erupt	Shed
EXE	Central incisor	8-12 mos.	6-7 yrs.
	Lateral incisor	9-13 mos.	7-8 yrs.
	Canine (cuspid)	16-22 mos.	10-12 yrs.
(z)	First molar	13-19 mos.	9-11 yrs.
(F)	Second molar	25-33 mos.	10-12 yrs.
\bigcirc	Lower Teeth	Erupt	Shed
	Second molar	23-31 mos.	10-12 yrs.
(\}\ / 2	First molar	14-18 mos.	9-11 yrs.
A	Canine (cuspid)	17-23 mos.	9-12 yrs.
	Lateral incisor	10-16 mos.	7-8 yrs.
	Central incisor	6-10 mos.	6-7 yrs.

Permanent Teeth	Upper Teeth	Erupt
-ECB	Central incisor	7-8 yrs.
	Lateral incisor	8-9 yrs.
	Canine (cuspid)	11-12 yrs.
	First premolar (first bicuspid)	10-11 yrs.
	Second premolar (second bicuspid)	10-12 yrs.
	First molar	6-7 yrs.
(₹) (₹) -	Second molar	12-13 yrs.
(T)	Third molar (wisdom tooth)	17-21 yrs.
	Lower Teeth	Erupt
	Third molar (wisdom tooth)	17-21 yrs.
(±) (±)	Second molar	11-13 yrs.
(4)	First molar	6-7 yrs.
	Second premolar (second bicuspid)	11-12 yrs.
	First premolar (first bicuspid)	10-12 yrs.
Share with	Canine (cuspid)	9-10 yrs.
	Lateral incisor	7-8 yrs.
	Central incisor	6-7 yrs.

Source: The American Dental Association