





Pascack Brook - FIBI059



-  FIBI Sampling Location
-  Small Streams (1st and 2nd Order)
-  Large Streams (3rd Order and Above)



SUMMARY OF RESULTS

FIBI059 - Pascack River



1. Stream Name:	Pascack River
2. Sampling Date:	07-30-2003
3. Sampling Location:	Emerson Rd
4. Municipality	Emerson
5. County:	Bergen
6. Watershed Management Area:	5
7. Contributing Drainage Area:	29 Square Miles
8. Electrofishing Gear:	Barge
9. FIBI Score and Rating:	36 - Fair
10. Habitat Score and Rating:	120 - Suboptimal
11. Fishable Species Present:	Yes
12. Relevant AMNET ¹ Station Data:	
Proximity of FIBI station to AMNET station:	0.94 mi downstream AN0207
AMNET Rating:	Round 1 - Moderate Round 2 – Non-impaired Round 3 - Moderate
13. Stream Chemistries:	
Dissolved Oxygen	7.6 mg/L
Temperature.	20.2 °C
pH	8.44
Conductivity	518 µmhos/cm
14. Number of Fish With Anomalies:	0
15. Length of Stream Segment Sampled	150 Meters
16. Water Clarity:	Slightly Turbid
17. Average Forest Open Canopy:	70.5%
18. Discharge:	58.7 ft. ³ /sec
19. Substrate:	65% Gravel and Sand, 5% Boulder, 20% Clay, 10% Silt
20. Habitat:	0% Riffle, 90% Run, 10% Pool
21. Snags	Yes
22. Periphyton	Slight
23. Submerged Aquatic Vegetation	No
24. Other observations:	
25. Number of Fish Species Identified:	10
26. Total Number of Fish Collected:	99

¹ AMNET is the acronym for the DEP's ambient benthic macroinvertebrate monitoring network – a series of 820 monitoring stations located throughout the state's waterways that collects data on the health of bottom dwelling stream fauna which in turn is used to assess general water quality.

FIBI059
PASCACK RIVER
EMERSON ROAD
WESTWOOD BORO, BERGEN COUNTY



Emerson Road

Kaufman Road



LEGEND

- START
- FINISH
- SEGMENT SAMPLED
- ↑ DIRECTION OF FLOW

FIBI059 - Pascack River @ Emerson Rd Excellent Good **Fair** Poor
Date Sampled - 7/30/2003

	Score
# of Fish Species	5
# of Benthic Insectivorous Species (BI)	3
# of Trout and Centrarchid Species (trout, bass, sunfish, crappie)	5
# of Intolerant Species (IS)	1
Proportion of Individuals as White Suckers	5
Proportion of Individuals as Generalists (carp, creek chub, banded killifish, goldfish, fathead minnow, green sunfish)	5
Proportion of Individuals as Insectivorous Cyprinids (I and BI)	1
Proportion of Individuals as Trout *whichever gives better score OR Proportion of Individuals as Piscivores (Excluding American Eel)*	3
Number of Individuals in Sample	3
Proportion of Individuals w/disease/anomalies (excluding blackspot)	5
Total	36

Stream Rating

45-50	Excellent
37-44	Good
29-36	Fair
10-28	Poor

HABITAT ASSESSMENT FOR *HIGH GRADIENT STREAMS* Pascack River (FIB1059) – 7/30/03

	Condition Category																			
	Optimal					Suboptimal					Marginal					Poor				
1. Epifaunal Substrate /Available Cover Greater than 70% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are <u>not</u> new fall and <u>not</u> transient). SCORE 12	20 19 18 17 16					15 14 13 12 11					10 9 8 7 6					5 4 3 2 1 0				
2. Embeddedness Gravel, cobble, and boulder particles are 0-25% surrounded by fine sediment. Layering of cobble provides diversity of niche space SCORE 15	20 19 18 17 16					15 14 13 12 11					10 9 8 7 6					5 4 3 2 1 0				
3. Velocity/Depth Regimes All 4 velocity/depth regimes present (slow-deep, slow-shallow, fast-deep, fast-shallow). (slow is <0.3 m/s, deep is >0.5 m) SCORE 8	20 19 18 17 16					15 14 13 12 11					10 9 8 7 6					5 4 3 2 1 0				
4. Sediment Deposition Little or no enlargement of islands or point bars and less than 5% (<20% for low-gradient streams) of the bottom affected by sediment deposition. SCORE 15	20 19 18 17 16					15 14 13 12 11					10 9 8 7 6					5 4 3 2 1 0				
5. Channel Flow Status Water reaches base of both lower banks, and minimal amount of channel substrate is exposed. SCORE 17	20 19 18 17 16					15 14 13 12 11					10 9 8 7 6					5 4 3 2 1 0				
6. Channel Alteration Channelization or dredging absent or minimal; stream with normal pattern. SCORE 16	20 19 18 17 16					15 14 13 12 11					10 9 8 7 6					5 4 3 2 1 0				
7. Frequency of Riffles (or bends) Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important. SCORE 5	20 19 18 17 16					15 14 13 12 11					10 9 8 7 6					5 4 3 2 1 0				
8. Bank Stability (score each bank) Note: determine left or right side by facing downstream. SCORE <u>6</u> (LB) SCORE <u>5</u> (RB)	Left 10 9 Right 10 9					8 7 6 8 7 6					5 4 3 5 4 3					2 1 0 2 1 0				
9. Bank Vegetative Protection (score each bank) SCORE <u>7</u> (LB) SCORE <u>5</u> (RB)	Left 10 9 Right 10 9					8 7 6 8 7 6					5 4 3 5 4 3					2 1 0 2 1 0				
10. Riparian Vegetative Zone Width (score each bank riparian zone) SCORE <u>7</u> (LB) SCORE <u>2</u> (RB)	Left 10 9 Right 10 9					8 7 6 8 7 6					5 4 3 5 4 3					2 1 0 2 1 0				

HABITAT SCORE

120

HABITAT SCORES	VALUE
OPTIMAL	160 – 200
SUB-OPTIMAL	110 – 159
MARGINAL	60 – 109
POOR	< 60

FIBI059 07-30-2003

Pascack River

LISTED IN ORDER OF ABUNDANCE FOUND

COMMON NAME	SCIENTIFIC NAME	# FOUND	SIZE RANGE (INCHES)
Tesselated darter	<i>Etheostoma olmstedi</i>	27	
Yellow perch	<i>Perca flavescens</i>	19	1.8 - 6.3
Pumpkinseed sunfish	<i>Lepomis gibbosus</i>	17	2.8 - 4.1
Largemouth bass	<i>Micropterus salmoides</i>	13	1.8 - 3.1
Bluegill sunfish	<i>Lepomis macrochirus</i>	8	4.5
Yellow bullhead	<i>Ameiurus natalis</i>	7	3.7 - 8.1
White sucker	<i>Catostomus commersoni</i>	4	
Green sunfish	<i>Lepomis cyanellus</i>	2	4.5 - 4.7
Redbreast sunfish	<i>Lepomis auritus</i>	1	3.9
Mummichog	<i>Fundulus heteroclitus</i>	1	

* Regulated as a fishable species under current New Jersey Fish and Wildlife codes

Species Identified at Pascack River (FIBI059)
(Not to Scale)

John Scarola



Yellow Bullhead

AFS



Largemouth Bass

John Scarola



Redbreast Sunfish

John Scarola



Yellow Perch

John Scarola



Mummichog


Konrad Schmidt



Green Sunfish

Species Identified at Pascack River (FIBI059)
(Not to Scale)

John Scarola



Tessellated Darter

John Scarola



Bluegill

John Scarola



White Sucker

John Scarola



Pumpkinseed