


Whippany River

FIBI074



FIBI Ratings

- Excellent (Green circle)
- Fair (Orange circle)
- Good (Yellow circle)
- Poor (Red circle)

- 1st & 2nd Order Streams (Dashed blue line)
- 3rd Order and Higher Streams (Solid blue line)



0 1 2 4 Miles



SUMMARY OF RESULTS – FIBI074





| | |
|---|--|
| 1. Stream Name: | Whippany River |
| 2. Sampling Date: | 08/06/2008 |
| 3. Sampling Location: | Whitehead Road |
| 4. Municipality: | Morris Twp. |
| 5. County: | Morris |
| 6. Watershed Management Area: | 5 |
| 7. Contributing Drainage Area (Sq. Mi.): | 8.4 |
| 8. Electrofishing Gear: | 2 Backpacks |
| 9. FIBI Score and Rating: | Round 1* Good (42); Round 2 Good (42) |
| 10. Habitat Score and Rating: | Round 1 Sub-Optimal (137); Round 2 Sub-Optimal (154) |
| 11. Fishable Species Present: | Yes |
| 12. Relevant AMNET ¹ Station Data: | |
| Proximity of FIBI station to AMNET station: | AN0233 |
| AMNET Rating: | R2 – Fair, R3 - Good |
| 13. Stream Chemistries: | |
| Dissolved Oxygen (mg/l) | 8.20 |
| Temperature °C. | 20.55 |
| pH | 5.99 |
| Conductivity (µmhos/cm) | 333 |
| 14. Length of Stream Sampled: | 150m |
| 15. Water Clarity: | Slightly Turbid |
| 16. Average Open Forest Canopy: | 9.6% |
| 17. Discharge: | 4.2 cfs |
| 18. Substrate: | 65% Gravel/Sand, 5% Cobble, 20% Mud, 10% Silt |
| 19. Habitat: | 10% Riffle, 55% Run, 35% Pool |
| 20. Snags: | Yes |
| 21. Periphyton: | None |
| 22. Submerged Aquatic Vegetation: | Yes |
| 23. Outfalls: | None |
| 24. Number of Fish Species Identified: | 14 |
| 25. Total Number of Fish Collected: | 327 |
| 26. Number of Fish With Anomalies: | 7 |
| 27. Other Observations: | |

¹ AMNET is the acronym for the DEP's ambient benthic macroinvertebrate monitoring network – a series of 820 monitoring stations located throughout the state's waterways that collects data on the health of bottom dwelling stream fauna which in turn is used to assess general water quality

* Round 1 data was scored prior to the FIBI metric recalibration.

FIBI074-R2

WHIPPANY RIVER
WHITEHEAD ROAD
MORRIS TWP.
MORRIS



**FIBI074-Whippany River @ Whitehead Rd.
Date Sampled - 8/06/2008**

Excellent **Good** Fair Poor

| | Score |
|--|-------------------------------|
| # of Fish Species | 5 |
| # of Benthic Insectivorous Species (BI) (excluding White Suckers and Bullheads) | 3 |
| # of Trout and Centrarchid Species (excluding Green Sunfish and Bluegill) | 5 |
| # of Intolerant Species (IS) | 5 |
| Proportion of Tolerant Individuals | 5 |
| Proportion of Individuals as Generalists | 5 |
| Proportion of Individuals as Insectivorous Cyprinids | 1 |
| Proportion of Individuals as Trout | *whichever gives better score |
| OR | |
| Proportion of Individuals as Piscivores (excluding American Eel)* | 5 |
| # of Individuals in Sample (excluding Tolerant Species) | 5 |
| Proportion of Individuals w/disease/anomalies (excluding blackspot) | 3 |
| Total | 42 |

| Stream Rating | |
|----------------------|-----------|
| 45-50 | Excellent |
| 37-44 | Good |
| 29-36 | Fair |
| 10-28 | Poor |

HABITAT ASSESSMENT FOR HIGH GRADIENT STREAMS Whippany R. (FIBI074) - 8/6/2008

| | Condition Category | | | | | | | | | | | | | | | | | | | | |
|--|---|----|----|----|----|--|----|----|----|----|---|---|---|---|---|---|---|---|---|---|---|
| | Optimal | | | | | Suboptimal | | | | | Marginal | | | | | Poor | | | | | |
| 1. Epifaunal Substrate /Available Cover | Greater than 70% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are <u>not</u> new fall and <u>not</u> transient). | | | | | 40-70% mix of stable habitat; well-suited for full colonization potential; adequate habitat for maintenance of populations; presence of additional substrate in the form of newfall, but not yet prepared for colonization (may rate at high end of scale). | | | | | 20-40% mix of stable habitat; habitat availability less than desirable; substrate frequently disturbed or removed. | | | | | Less than 20% stable habitat; lack of habitat is obvious; substrate unstable or lacking. | | | | | |
| SCORE 18 | 20 | 19 | 18 | 17 | 16 | 15 | 14 | 13 | 12 | 11 | 10 | 9 | 8 | 7 | 6 | 5 | 4 | 3 | 2 | 1 | 0 |
| 2. Embeddedness | Gravel, cobble, and boulder particles are 0-25% surrounded by fine sediment. Layering of cobble provides diversity of niche space | | | | | Gravel, cobble, and boulder particles are 25-50% surrounded by fine sediment. | | | | | Gravel, cobble, and boulder particles are 50-75% surrounded by fine sediment. | | | | | Gravel, cobble, and boulder particles are more than 75% surrounded by fine sediment. | | | | | |
| SCORE 11 | 20 | 19 | 18 | 17 | 16 | 15 | 14 | 13 | 12 | 11 | 10 | 9 | 8 | 7 | 6 | 5 | 4 | 3 | 2 | 1 | 0 |
| 3. Velocity/Depth Regimes | All 4 velocity/depth regimes present (slow-deep, slow-shallow, fast-deep, fast-shallow). (slow is <0.3 m/s, deep is >0.5 m) | | | | | Only 3 of the 4 regimes present (if fast-shallow is missing, score lower than if missing other regimes). | | | | | Only 2 of the 4 habitat regimes present (if fast-shallow or slow-shallow are missing, score low). | | | | | Dominated by 1 velocity / depth regime (usually slow-deep). | | | | | |
| SCORE 19 | 20 | 19 | 18 | 17 | 16 | 15 | 14 | 13 | 12 | 11 | 10 | 9 | 8 | 7 | 6 | 5 | 4 | 3 | 2 | 1 | 0 |
| 4. Sediment Deposition | Little or no enlargement of islands or point bars and less than 5% (<20% for low-gradient streams) of the bottom affected by sediment deposition. | | | | | Some new increase in bar formation, mostly from gravel, sand or fine sediment; 5-30% (20-50% for low-gradient) of the bottom affected; slight deposition in pools. | | | | | Moderate deposition of new gravel, sand or fine sediment on old and new bars; 30-50% (50-80% for low-gradient) of the bottom affected; sediment deposits at obstructions, constrictions, and bends; moderate deposition of pools prevalent. | | | | | Heavy deposits of fine material, increased bar development; more than 50% (80% for low-gradient) of the bottom changing frequently; pools almost absent due to substantial sediment deposition. | | | | | |
| SCORE 10 | 20 | 19 | 18 | 17 | 16 | 15 | 14 | 13 | 12 | 11 | 10 | 9 | 8 | 7 | 6 | 5 | 4 | 3 | 2 | 1 | 0 |
| 5. Channel Flow Status | Water reaches base of both lower banks, and minimal amount of channel substrate is exposed. | | | | | Water fills >75% of the available channel; or <25% of channel substrate is exposed. | | | | | Water fills 25-75% of the available channel, and/or riffle substrates are mostly exposed. | | | | | Very little water in channel and mostly present as standing pools. | | | | | |
| SCORE 18 | 20 | 19 | 18 | 17 | 16 | 15 | 14 | 13 | 12 | 11 | 10 | 9 | 8 | 7 | 6 | 5 | 4 | 3 | 2 | 1 | 0 |
| 6. Channel Alteration | Channelization or dredging absent or minimal; stream with normal pattern. | | | | | Some channelization present, usually in areas of bridge abutments; evidence of past channelization, i.e., dredging, (greater than past 20 yr) may be present, but recent channelization is not present. | | | | | Channelization may be extensive; embankments or shoring structures present on both banks; and 40 to 80% of stream reach channelized and disrupted. | | | | | Banks shored with gabion or cement; over 80% of the stream reach channelized and disrupted. In stream habitat greatly altered or removed entirely. | | | | | |
| SCORE 20 | 20 | 19 | 18 | 17 | 16 | 15 | 14 | 13 | 12 | 11 | 10 | 9 | 8 | 7 | 6 | 5 | 4 | 3 | 2 | 1 | 0 |
| 7. Frequency of Riffles (or bends) | Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important. | | | | | Occurrence of riffles infrequent; distance between riffles divided by the width of the stream is between 7 to 15. | | | | | Occasional riffle or bend; bottom contours provide some habitat; distance between riffles divided by the width of the stream is between 15 to 25. | | | | | Generally all flat water or shallow riffles; poor habitat; distance between riffles divided by the width of the stream is a ratio of >25. | | | | | |
| SCORE 5 | 20 | 19 | 18 | 17 | 16 | 15 | 14 | 13 | 12 | 11 | 10 | 9 | 8 | 7 | 6 | 5 | 4 | 3 | 2 | 1 | 0 |
| 8. Bank Stability (score each bank) Note: determine left or right side by facing downstream. | Banks stable; evidence of erosion or bank failure absent or minimal; little potential for future problems. <5% of bank affected. | | | | | Moderately stable; infrequent, small areas of erosion mostly healed over. 5-30% of bank in reach has areas of erosion. | | | | | Moderately unstable; 30-60% of bank in reach has areas of erosion; high erosion potential during floods. | | | | | Unstable; many eroded areas; "raw" areas frequent along straight sections and bends; obvious bank sloughing; 60-100% of bank has erosional scars. | | | | | |
| SCORE 7 (LB) | Left | 10 | 9 | | | 8 | 7 | 6 | | | 5 | 4 | 3 | | | 2 | 1 | 0 | | | |
| SCORE 5 (RB) | Right | 10 | 9 | | | 8 | 7 | 6 | | | 5 | 4 | 3 | | | 2 | 1 | 0 | | | |
| 9. Bank Vegetative Protection (score each bank) | More than 90% of the streambank surfaces and immediate riparian zone covered by native vegetation, including trees, under story shrubs, or nonwoody macrophytes; vegetative disruption through grazing or mowing minimal or not evident; almost all plants allowed to grow naturally. | | | | | 70-90% of the streambank surfaces covered by native vegetation, but one class of plants is not well-represented; disruption evident but not affecting full plant growth potential to any great extent; more than one-half of the potential plant stubble height remaining. | | | | | 50-70% of the streambank surfaces covered by vegetation; disruption obvious; patches of bare soil or closely cropped vegetation common; less than one-half of the potential plant stubble height remaining. | | | | | Less than 50% of the streambank surfaces covered by vegetation; disruption of streambank vegetation is very high; vegetation has been removed to 5 centimeters or less in average stubble height. | | | | | |
| SCORE 10 (LB) | Left | 10 | 9 | | | 8 | 7 | 6 | | | 5 | 4 | 3 | | | 2 | 1 | 0 | | | |
| SCORE 10 (RB) | Right | 10 | 9 | | | 8 | 7 | 6 | | | 5 | 4 | 3 | | | 2 | 1 | 0 | | | |
| 10. Riparian Vegetative Zone Width (score each bank riparian zone) | Width of riparian zone >18 meters; human activities (i.e., parking lots, roadbeds, clear-cuts, lawns, or crops) have not impacted zone. | | | | | Width of riparian zone 12-18 meters; human activities have impacted zone only minimally. | | | | | Width of riparian zone 6-12 meters; human activities have impacted zone a great deal. | | | | | Width of riparian zone <6 meters; little or no riparian vegetation due to human activities. | | | | | |
| SCORE 10 (LB) | Left | 10 | 9 | | | 8 | 7 | 6 | | | 5 | 4 | 3 | | | 2 | 1 | 0 | | | |
| SCORE 10 (RB) | Right | 10 | 9 | | | 8 | 7 | 6 | | | 5 | 4 | 3 | | | 2 | 1 | 0 | | | |

HABITAT SCORE

153

| HABITAT SCORES | VALUE |
|--------------------|------------------|
| OPTIMAL | 160 - 200 |
| SUB-OPTIMAL | 110 - 159 |
| MARGINAL | 60 - 109 |
| POOR | < 60 |

FIBI074-R2

Whippany River

08/06/2008

| Common Name | Scientific Name | Abundance | Size Range (inches) |
|------------------------|-----------------------------------|-----------|---------------------|
| Tessellated Darter | <i>Etheostoma olmstedii</i> | 153 | - |
| American Brook Lamprey | <i>Lampetra appendix</i> | 86 | - |
| White Sucker | <i>Catostomus commersoni</i> | 23 | - |
| Redfin Pickerel | <i>Esox americanus americanus</i> | 14 | 3.5 - 9.8 |
| Brown Trout | <i>Salmo trutta</i> | 12 | 2.2 - 11.6 |
| Eastern Mudminnow | <i>Umbra pygmaea</i> | 8 | - |
| Common Shiner | <i>Luxilus cornutus</i> | 8 | - |
| Spottail Shiner | <i>Notropis hudsonius</i> | 6 | - |
| Largemouth Bass | <i>Micropterus salmoides</i> | 5 | 2.0 - 5.2 |
| Blacknose Dace | <i>Rhinichthys atratulus</i> | 4 | - |
| Chain Pickerel | <i>Esox niger</i> | 3 | 3.9 - 4.1 |
| Creek Chubsucker | <i>Erimyzon oblongus</i> | 3 | - |
| Brook Trout - stocked | <i>Salvelinus fontinalis</i> | 1 | 11.7- 11.7 |
| Bluegill | <i>Lepomis macrochirus</i> | 1 | 3.8 - 3.8 |

Whippany River - FIBI074



American Brook Lamprey



White Sucker



Tessellated Darter



Eastern Mudminnow



Spottail Shiner



Common Shiner

Whippany River - FIBI074



Blacknose Dace



Chain Pickerel



Bluegill



Brook Trout



Largemouth Bass



Brown Trout

Whippany River - FIBI074



Redfin Pickerel



Creek Chubsucker