

New Jersey Department of Health and Senior Services Children's Oral Health Program **Tooth Eruption**



Primary Teeth	Upper Teeth	Erupt	Shed
	 Central incisor 	8-12 mos.	6-7 yrs.
	 Lateral incisor 	9-13 mos.	7-8 yrs.
	 Canine (cuspid) 	16-22 mos.	10-12 yrs.
ET CA	First molar	13-19 mos.	9-11 yrs.
	Second molar	25-33 mos.	10-12 yrs.
4			10 12 9101
(Λ) (I)	Lower Teeth	Erupt	Shed
	 Second molar 	23-31 mos.	10-12 yrs.
(γ) (γ)	 First molar 	14-18 mos.	9-11 yrs.
	Canine (cuspid)	17-23 mos.	9-12 yrs.
Smith	 Lateral incisor 	10-16 mos.	7-8 yrs.
- Cut	 Central incisor 	6-10 mos.	6-7 yrs.
Permanent Teeth	Upper Teeth		Erupt
CER DO	 Central incisor 		7-8 yrs.
CM2-	 Lateral incisor 		8-9 yrs.
	 Canine (cuspid) 		11-12 yrs.
	 First premolar (first bicuspid) 		10-11 yrs.
	Second premolar (second bicuspid)		10-12 yrs.
	- Electron 1 and		
	First molar		6-7 yrs.
	Second molar		12-13 yrs.
		h)	
	Second molar	h)	12-13 yrs.
	Second molarThird molar (wisdom toot		12-13 yrs. 17-21 yrs.
	 Second molar Third molar (wisdom toot <i>Lower Teeth</i> 		12-13 yrs. 17-21 yrs. Erupt
E E E E E E E E E E E E E E E E E E E	Second molar Third molar (wisdom toot <i>Lower Teeth</i> Third molar (wisdom tooth		12-13 yrs. 17-21 yrs. Erupt 17-21 yrs.
E A A A A A A A A A A A A A A A A A A A	Second molar Third molar (wisdom toot <i>Lower Teeth</i> Third molar (wisdom tooth Second molar)	12-13 yrs. 17-21 yrs. Erupt 17-21 yrs. 11-13 yrs.
E A A A A A A A A A A A A A A A A A A A	Second molar Third molar (wisdom toot <i>Lower Teeth</i> Third molar (wisdom tooth Second molar First molar) vicuspid)	12-13 yrs. 17-21 yrs. Erupt 17-21 yrs. 11-13 yrs. 6-7 yrs.
E A A A A A A A A A A A A A A A A A A A	Second molar Third molar (wisdom toot <i>Lower Teeth</i> Third molar (wisdom tooth Second molar First molar Second premolar (second b) vicuspid)	12-13 yrs. 17-21 yrs. Erupt 17-21 yrs. 11-13 yrs. 6-7 yrs. 11-12 yrs.
K X X X X X X X X X X X X X X	Second molar Third molar (wisdom toot <i>Lower Teeth</i> Third molar (wisdom tooth Second molar First molar Second premolar (second b First premolar (first bicusp) vicuspid)	12-13 yrs. 17-21 yrs. Erupt 17-21 yrs. 11-13 yrs. 6-7 yrs. 11-12 yrs. 10-12 yrs.

Source: The American Dental Association



New Jersey Department of Health and Senior Services Children's Oral Health Program **Tooth Eruption**



Drimory Tooth	Upper Teeth	Erupt	Shed
Primary Teeth	Central incisor	8-12 mos.	6-7 yrs.
and	 Lateral incisor 	9-13 mos.	7-8 yrs.
	Canine (cuspid)	16-22 mos.	10-12 yrs.
ET C	First molar	13-19 mos.	9-11 yrs.
	Second molar	25-33 mos.	10-12 yrs.
(4)			
$(A) \qquad (I)$	Lower Teeth	Erupt	Shed
	 Second molar 	23-31 mos.	10-12 yrs.
(γ) (γ)	 First molar 	14-18 mos.	9-11 yrs.
	- Canine (cuspid)	17-23 mos.	9-12 yrs.
	 Lateral incisor 	10-16 mos.	7-8 yrs.
	Central incisor	6-10 mos.	6-7 yrs.
Permanent Teeth	Upper Teeth		Erupt
ED	 Central incisor 		7-8 yrs.
AND_	 Lateral incisor 		8-9 yrs.
	 Canine (cuspid) 		11-12 yrs.
	 First premolar (first bicusp 	id)	10-11 yrs.
	 Second premolar (second b) 	picuspid)	10-12 yrs.
	 First molar 		6-7 yrs.
	 Second molar 		12-13 yrs.
(A) (F)	Third molar (wisdom tooth	n)	17-21 yrs.
	Lower Teeth		Erupt
	• Third molar (wisdom tooth))	17-21 yrs.
	Second molar		11-13 yrs.
\mathbf{F} \mathbf{F}	First molar		6-7 yrs.
The second second	• Second premolar (second bi	cuspid)	11-12 yrs.
The state	• First premolar (first bicuspi	d)	10-12 yrs.
Charles and	Canine (cuspid)		9-10 yrs.
	Lateral incisor		7-8 yrs.
L	Central incisor		6-7 yrs.