



**LANDSLIDE SUSCEPTIBILITY
OF
PASSAIC COUNTY, NEW JERSEY**

Prepared by Scott D. Stanford, New Jersey Geological Survey
for the
New Jersey State Police, Office of Emergency Management

2004

- None—HAZUS number 0
- Landslide Class A I—strongly cemented rock, slope angle 15-20 degrees (HAZUS number 1)
- Landslide Class A II—strongly cemented rock, slope angle 20-30 degrees (HAZUS number 2)
- Landslide Class A IV—strongly cemented rock, slope angle 30-40 degrees (HAZUS number 5)
- Landslide Class B III—weakly cemented rock and soil, slope angle 10-15 degrees (HAZUS number 3)
- Landslide Class B IV—weakly cemented rock and soil, slope angle 15-20 degrees (HAZUS number 4)
- Landslide Class B V—weakly cemented rock and soil, slope angle 20-30 degrees (HAZUS number 7)

Landslide classes are from the HAZUS User's Manual, Table 9.2 (National Institute of Building Sciences, 1997). Slope angles were measured from the following U. S. Geological Survey 7.5 minute quadrangles: Ramsey, Hackensack, Paterson, Weehawken (all with 10 foot contour interval), and Orange, Wanaque, Newfoundland, Wawayanda, Greenwood Lake, Pompton Plain and Slootsburg (20 foot contour interval). Slope materials were determined from field mapping (Stanford, 1991, 1992a, 1992b, 1993a, 1993b, 1993c, 2001, 2003; Stone and others, 2002).

REFERENCES CITED

National Institute of Building Sciences, 1997, HAZUS user's manual: Washington, D. C., National Institute of Building Sciences Publication 5200.

Stanford, S. D., 1991, Surficial geology of the Newfoundland quadrangle, Passaic, Morris, and Sussex counties, New Jersey: N. J. Geological Survey Geologic Map Series 91-3, scale 1:24,000.

Stanford, S. D., 1992a, Surficial geology of the Greenwood Lake and Slootsburg quadrangles, Passaic, Bergen, and Sussex counties, New Jersey: N. J. Geological Survey Open File Map 6, scale 1:24,000.

Stanford, S. D., 1992b, Surficial geology of the Wawayanda and Pine Island quadrangles, Sussex and Passaic counties, New Jersey: N. J. Geological Survey Open File Map 9, scale 1:24,000.

Stanford, S. D., 1993a, Surficial geology of the Weehawken and Central Park quadrangles, Bergen, Hudson, and Passaic counties, New Jersey: N. J. Geological Survey Geologic Map Series 93-1, scale 1:24,000.

Stanford, S. D., 1993b, Surficial geology of the Hackensack quadrangle, Bergen and Passaic counties, New Jersey: N. J. Geological Survey Open File Map 14, scale 1:24,000.

Stanford, S. D., 1993c, Surficial geology of the Wanaque quadrangle, Bergen, Morris, and Passaic counties, New Jersey: N. J. Geological Survey Geologic Map Series 93-4, scale 1:24,000.

Stanford, S. D., 2001, Surficial geology of the Orange quadrangle, Essex, Passaic, Hudson, and Bergen counties, New Jersey: N. J. Geological Survey Open File Map 41, scale 1:24,000.

Stanford, S. D., 2003, Surficial geology of the Paterson quadrangle, Passaic, Bergen, and Essex counties, New Jersey: N. J. Geological Survey Open File Map 54, scale 1:24,000.

Stone, B. D., Stanford, S. D., Witte, R. W., 2002, Surficial geologic map of northern New Jersey: U. S. Geological Survey Miscellaneous Investigations Series Map 1-2540-C, scale 1:100,000.

SCALE 1:48000

